

Comparison of Swarming Tendency and Defensive Behavior of Yığılca Local and Other Commonly Used Honeybee Genotypes in Turkey ^[1]

Ayhan GÖSTERİT *  Yakup ÇIKILI ** Meral KEKEÇOĞLU ***

[1] This study was supported by grant TUBITAK TOVAG-110 O 432 from the Scientific and Technological Research Council of Turkey

* Süleyman Demirel University, Faculty of Agriculture, Department of Animal Science, TR-32260 Isparta - TURKEY

** Düzce University, Çilimli Vocational School, TR-81750 Düzce - TURKEY

*** Düzce University, Faculty of Science, Department of Biology, TR-81620 Düzce - TURKEY

Makale Kodu (Article Code): KVFD-2011-5996

Summary

In this study, swarming tendency and defensive behavior of Yığılca local honeybee were determined and compared with other commonly used honeybee genotypes in Turkey. Colonies were headed by naturally mated queens and 10 colonies of Yığılca local honeybee, 12 colonies of *Apis mellifera caucasica* cross and 12 colonies of *Apis mellifera anatoliaca* cross were used in the experiment. In swarming season, Yığılca honeybee colonies constructed more queen cells (49.86 ± 18.00) than both *A.m. anatoliaca* cross (13.00 ± 7.00) and *A.m. caucasica* (8.00 ± 1.15) cross colonies. Similarly, according to results of sting test, the highest number of stings was determined in Yığılca honeybee colonies (18.38 ± 4.24), followed by *A.m. anatoliaca* (5.50 ± 2.15) and *A.m. caucasica* (3.75 ± 0.62) crosses. Results showed that Yığılca local honeybee genotype has a more swarming tendency and is more defensive than *A.m. anatoliaca* and *A.m. caucasica* crosses.

Keywords: Yığılca honeybee, Swarming, Defensive behavior, Genetic conservation

Yığılca Yerel Bal Arısının Oğul Verme Eğilimi ve Savunma Davranışı Bakımından Türkiye'de Yaygın Olarak Kullanılan Diğer Genotipler ile Karşılaştırılması

Özet

Bu çalışmada, Yığılca yerel bal arısının oğul verme eğilimi ve savunma davranışının belirlenmesi ve Türkiye'de yaygın olarak kullanılan diğer bal arısı genotipleri ile karşılaştırılması amaçlanmıştır. Araştırmada, doğal çiftleşmiş ana arılar ile oluşturulan Yığılca bal arısı genotipinden 10 koloni, Anadolu ve Kafkas ırkı melezlerinden ise 12'şer koloni olmak üzere toplam 34 adet bal arısı kolonisi kullanılmıştır. Oğul verme mevsimi süresince Yığılca bal arısı kolonilerinin (49.86 ± 18.00) hem Anadolu ırkı melezi kolonilerinden (13.00 ± 7.00) hem de Kafkas ırkı melezi kolonilerinden (8.00 ± 1.15) daha fazla ana arı yüksüğü yaptıkları belirlenmiştir. Benzer şekilde sokma testi sonuçlarına göre, en yüksek iğne sayısı Yığılca bal arısı kolonilerinde (18.38 ± 4.24) belirlenirken, bu grubu Anadolu ırkı melezi (5.50 ± 2.15) ve Kafkas ırkı melezi (3.75 ± 0.62) koloniler takip etmiştir. Araştırma sonuçları Yığılca yerel bal arısı genotipinin Anadolu ve Kafkas ırkı melezlerine göre daha fazla oğul verme eğilimi gösterdiği ve daha hırçın olduğunu ortaya koymuştur.

Anahtar sözcükler: Yığılca bal arısı, Oğul, Savunma davranışı, Genetik koruma

INTRODUCTION

The honeybee (*Apis mellifera* L.), the most economically important beneficial insect, has proven to be highly adaptive to a wide variety of ecosystems in its native range of Africa, Europe, and Central and Western Asia. Across this

range, numerous subspecies and ecotypes of *A. mellifera* have been described based upon behavior, morphology and molecular evidence. While some subspecies inhabit large geographic areas, some subspecies and all ecotypes



İletişim (Correspondence)



+90 246 2114653



ayhangosterit@sdu.edu.tr

inhabit relatively small geographic areas with smaller population size ¹⁻⁵. Specific behavioral and phenological adaptations to local environment conditions are known for some honeybee populations ^{1,6,7}. The survival of these locally adapted populations or ecotypes results from a number of traits conferred selective advantage to the population within an ecologically distinct area ⁸. Each honeybee genotypes that genetically different have their own peculiar behavior traits. Within the endemic range of the honeybee, the variation between the behavioral traits, such as swarming and defensive, provides the basis for subspecific classification ^{9,10}.

Several honeybee races and native ecotypes have been adapted to different ecological regions of Turkey. They have some differences in their morphology, behavior and physiology according to the environmental conditions. Based on multivariate analysis of morphometric data, Ruttner ¹ suggested that three subspecies exist in Turkey: *A.m. caucasica* in the northeast, *A.m. meda* in the southeast and *A.m. anatoliaca* throughout the rest of the country including Thrace (European part of Turkey). Also some researches studied the honeybee populations using different approach and reported that there are another *A. mellifera* subspecies in Turkey: *A.m. carnica* in Thrace and *A.m. syriaca* in the southeast ¹¹⁻¹⁴.

A.m. anatoliaca and *A.m. caucasica* are used intensively for commercial queen bee rearing in Turkey. For this reason, it is presumed that these two honeybee genotypes and their crosses constitute the majority of honeybee population in the country. Gentle and productive *A.m. caucasica* also has been used in beekeeping in many places around the world especially in higher elevations ^{15,16}. The Anatolian honeybees, *A.m. anatoliaca*, have many ecotypes and local populations that differ from each other morphologically, physiologically and behaviorally ¹⁷. One of the local populations inhabits in Yiğilca, a small geographic area in Western Black Sea Region in Turkey. Morphometric and genetic studies reported that Yiğilca local honeybee population differed from other populations of *A.m. anatoliaca* ¹⁸⁻²⁰. Despite the wide spread efforts for genetic conservation of this population and great demand for queen and colony from beekeepers, there is no data about its behavioral, colony development and production characteristics. Therefore, the aim of the present study was to determine the swarming tendency and defensive behavior of Yiğilca honeybee and compare with *A.m. caucasica* and *A.m. anatoliaca* crosses which are the most common genotypes in honeybee population of Turkey.

MATERIAL and METHODS

The experiment was carried out in Düzce, Turkey. Thirty four colonies headed by naturally mated queens were used in the experiment; 10 colonies of Yiğilca local honeybee, 12 colonies of *A.m. caucasica* and 12 colonies of *A.m. anatoliaca*.

Queens of *A.m. caucasica* and *A.m. anatoliaca* were reared in apiaries where queens are reared commercially at Central Anatolia Region. Yiğilca honeybee queens were reared from original breeder colonies of this genotype in Yiğilca (Düzce), according to Laidlaw ²¹. In all groups, queens were reared from different breeder colonies selected by randomly for representation of the population. Each mated queen was introduced into 6-frame colonies. These colonies were managed the same way for a period of 60 days prior to the beginning of the experiments to allow time for workers in the colony to be replaced by daughters of the new queens ²². The experimental colonies were equalized with regard to adult bee, brood and food stocks. They were kept in Langstroth hives in the same apiary and evaluated for swarming tendency and defensive behavior for a reproduction season.

In swarming season from April to July, all colonies in the genotype groups were controlled every week under normal colony management conditions. During the observation period, the numbers of colony had a swarming tendency and the numbers of queen cell in these colonies were recorded to assay the swarming tendency of genotypes. All sealed and unsealed queen cells were counted and destroyed in queenright colonies ⁶. To quantify defensive behavior, black coloured suede target attached to the end of a one meter stick was waved for one minute by hand in a rhythmic way approximately 10 cm from the entrance of the hive to stimulate the defensive response of colonies. The numbers of the stings in the target were recorded in order to measure the defensive trait of each colony. This process was repeated 8 times at different dates and two colonies which were equal with regard to worker population from each genotype were tested simultaneously in each repetition ^{6,23}. For statistical analysis, SPSS statistical program was used. Chi-square test was performed to analyze the data of swarming tendency. Swarming tendency of genotype groups was also compared by t-test. Defensive behavior trait data of the three genotype groups were analyzed by ANOVA and the means were compared using the Duncan multiple comparison test.

RESULTS

The swarming tendency and the number of queen cells constructed in groups were given in [Table 1](#). Results of χ^2 analyses showed that there was significant difference in the swarming tendency between the genotype groups. According to the results, Yiğilca local honeybee colonies constructed more queen cells than other genotypes ($P < 0.05$). In swarming season from April to July, 2 of 11 colonies (18.18%) in Anatolian group, 3 of 10 colonies (30.00%) in Caucasian group and 7 of 9 colonies (77.77%) in Yiğilca group constructed queen cells known as swarm marker.

According to results of sting test, defensive behavior of genotypes (the numbers of sting) were significantly

Table 1. Swarming tendency and number of queen cells in genotype groups (n = number of colonies had a swarming tendency)**Tablo 1.** Genotip gruplarında oğul verme eğilimi ve ana arı yüksek sayıları (n= oğul verme eğilimi gösteren koloni sayısı)

Genotypes	Number of Queen Cells				Swarming Tendency (%)
	n	X±Sx	Min	Max	
<i>A.m. anatoliaca</i>	2	13.00±7.00	6	20	18.18b
<i>A.m. caucasica</i>	3	8.00±1.15	6	10	30.00b
Yığılca Local Honeybee	7	49.86±18.00	2	136	77.77a

Means followed by different letters (a,b) in the same column are different (P<0.05)

Table 2. Number of stings (results of sting test) in genotype groups (n = number of colonies)**Tablo 2.** Genotip gruplarında belirlenen iğne sayıları (n = koloni sayısı)

Genotypes	n	X±Sx	Min	Max
<i>A.m. anatoliaca</i>	6	5.50±2.15b	1	18
<i>A.m. caucasica</i>	5	3.75±0.62b	2	6
Yığılca Local Honeybee	4	18.38±4.24a	6	40

Means followed by different letters (a,b) in the same column are different (P<0.01)

different each other (P<0.01). In the present study, *A.m. caucasica* was the gentlest genotype and Yığılca local honeybee was the most defensive genotype. Although Yığılca local honeybee was significantly different from *A.m. caucasica* and *A.m. anatoliaca* in terms of defensive behavior, there was no significant difference between the *A.m. caucasica* and *A.m. anatoliaca* (Table 2).

DISCUSSION

The honeybee population show a great genetic variation and this variation provide some advantages to improve the honeybee culture in Turkey²⁴. Several honeybee subspecies and ecotypes have been adapted to different ecological regions^{1,12,14,17}. *A.m. anatoliaca* and *A.m. caucasica* subspecies, and their reciprocal crosses constitute the majority of honeybee population in the country. There are several local populations of the *A.m. anatoliaca*, such as Muğla bees, Yığılca bees and Giresun bees. However, these local genotypes were not isolated in their geographical range, less well defined and need further investigation⁵.

The survival of local or native honeybee populations results from a number of traits commonly perceived as adaptive, many of which are related to reproductive swarming and defensive behavior^{6,10,25}. This present study clearly showed that Yığılca genotype is more inclined to swarm and is more defensive than *A.m. anatoliaca* and *A.m. caucasica* crosses. Yığılca honeybee colonies constructed more queen cells (49.86±18.00) than both *A.m. anatoliaca* (13.00±7.00) and *A.m. caucasica* (8.00±1.15) crosses. Genç et al.⁶ reported that Erzurum honeybee genotype had more defensive behavior but had not more swarming tendency than *A.m. anatoliaca* and *A.m. caucasica*. Similarly, Yücel and Kösoğlu²⁵ found that Muğla ecotype showed better performance for adaptation to environmental conditions and had more defensive behavior than Italian

cross. The variation in swarming tendency and defensive behavior can be used for classification of honeybee population^{9,10}. Morphometric and genetic studies also suggested that endemic honeybee in Yığılca province of Düzce distinct from the other population and the conservation of this genotype in its native range may be worthwhile¹⁸⁻²⁰.

The importation of foreign queens and the practice of moving hives several times in a year are factors that can affect the genetic structure of a local honeybee population through genetic introgression²⁶. It is known that the migratory beekeeping is practiced extensively in Turkey. However, local beekeepers claimed that neither foreign honeybee colonies nor queens were introduced into Yığılca location and colonies have been managed traditionally. Therefore, it should be considered that endemic honeybee population may exist in Yığılca location. In many countries of Europe, there is a kind of "certification" for local honeybee populations and official conservation programme have been managed⁵. Although conservation of Yığılca local honeybee is also proposed, its adaptive characteristics have not been studied making it difficult to rapidly implement conservation efforts. Identify the other characteristics of this local genotype in both their native location and other region, for this reason, need to be examined in future studies.

ACKNOWLEDGEMENTS

This study was supported by grant TUBITAK TOVAG-1100432 from the Scientific and Technological Research Council of Turkey (Project coordinator: Dr. Ayhan GÖSTERİT).

REFERENCES

1. Ruttner F: Biogeography and taxonomy of honeybees. Springer-Verlag, Berlin, 1988.

- 2. Sheppard WS, Arias MC, Grech A, Meixner MD:** *Apis mellifera ruttneri*, a new honey bee subspecies from Malta. *Apidologie*, 28, 287-293, 1997.
- 3. Sheppard WS, Meixner MD:** *Apis mellifera pomonella*, a new honey bee subspecies from Central Asia. *Apidologie*, 34, 367-375, 2003.
- 4. Strange JP, Garnery L, Sheppard WS:** Morphological and molecular characterization of the Landes honey bee (*Apis mellifera* L.) ecotype for genetic conservation. *J Insect Conserv*, 12, 527-537, 2008.
- 5. Bouga M, Alaux C, Bienkowska M, Büchler R, Carreck NL, Cauia E, Chlebo R, Dahle B, Dall'Olio R, De la Rúa P, Gregorc A, Ivanova E, Kence A, Kence M, Kezic N, Kiprijanovska H, Kozmus P, Kryger P, Le Conte Y, Lodesani M, Murilhas AM, Siceanu A, Soland G, Uzunov A, Wilde J:** A review of methods for discrimination of honey bee populations as applied to European beekeeping. *J Apicult Res*, 50 (1): 51-84, 2011.
- 6. Genç F, Dülger C, Kutluca S, Dodoloğlu A:** Comparison of some behavioural characteristics of Caucasian, Central Anatolian and Erzurum honeybee (*Apis mellifera* L.) genotypes in the conditions of Erzurum. *Turk J Vet Anim Sci*, 23, 651-656, 1999.
- 7. Andere C, Palacio MA, Rodriguez EM, Figini E, Dominguez MT, Bedascarrasbure E:** Evaluation of the defensive behavior of two honeybee ecotypes using a laboratory test. *Genet Mol Biol*, 25 (1): 57-60, 2002.
- 8. Strange JP, Garnery L, Sheppard WS:** Persistence of the Landes ecotype of *Apis mellifera mellifera* in southwest France: Confirmation of a locally adaptive annual brood cycle trait. *Apidologie*, 38, 259-267, 2007.
- 9. Guzman-Novoa E, Hunt GJ, Uribe-Rubio JL, Prieto-Merlos D:** Genotypic effects of honey bee (*Apis mellifera*) defensive behavior at the individual and colony levels: the relationship of guarding, pursuing and stinging. *Apidologie*, 35, 15-24, 2004.
- 10. Villa JD:** Swarming behavior of honey bees (Hymenoptera: Apidae) in Southeastern Louisiana. *Ann Entomol Soc Am*, 97 (1): 111-116, 2004.
- 11. Kandemir I, Kence A:** Allozyme variability in a Central Anatolian honeybee (*Apis mellifera* L.) population. *Apidologie*, 26, 503-10, 1995.
- 12. Smith DR, Slaymaker A, Palmer M, Kaftanoglu O:** Turkish honey bees belong to the east Mediterranean mitochondrial lineage. *Apidologie*, 28, 269-274, 1997.
- 13. Kandemir I, Kence M, Kence A:** Genetic and morphometric variation in honeybee (*Apis mellifera* L.) populations of Turkey. *Apidologie*, 31, 343-56, 2000.
- 14. Palmer MN, Smith DR, Kaftanoglu O:** Turkish honeybees: Genetic variation and evidence for a fourth lineage of *Apis mellifera* mtDNA. *J Hered*, 91, 42-46, 2000.
- 15. Adl MBF, Gençer HV, Firatlı Ç, Bahreini R:** Morphometric characterization of Iranian (*Apis mellifera meda*), Central Anatolian (*Apis mellifera anatoliaca*) and Caucasian (*Apis mellifera caucasica*) honey bee populations. *J Apicult Res Bee World*, 46 (4): 225-231, 2007.
- 16. Güler A:** A morphometrics model for determining the effect of commercial queen bee usage on the native honeybee (*Apis mellifera* L.) population in Turkish province. *Apidologie*, 41, 622-635, 2010.
- 17. Güler A, Kaftanoğlu O:** Türkiye'deki önemli balırsı (*Apis mellifera* L.) ırk ve ekotiplerinin morfolojik özellikleri-I. *Turk J Vet Anim Sci*, 23, 565-570, 1999.
- 18. Kekeçoğlu M:** Honey bee biodiversity in Western Black Sea and evidence for a new honey bee ecotype in Yiğilca province. *Res J Biol Sci*, 2 (1): 73-78, 2009.
- 19. Kekeçoğlu M, Soysal Mİ:** Genetic diversity of bee ecotypes in Turkey and evidence for geographical differences. *Rom Biotech Lett*, 15 (5): 5646-5653, 2010.
- 20. Güler A, Bıyık S, Güler M:** Morphological characteristics of the honey bee (*Apis mellifera*) population of the Western Black Sea Region. *7th National Animal Science Congress*, Sep 14-16, Adana, Turkey, 2011.
- 21. Laidlaw HH:** Contemporary queen rearing. A Dadant Publication, Dadant and Sons, Hamilton, Illinois, U.S.A., 1985
- 22. Arechavaleta-Velasco ME, Hunt GJ:** Genotypic variation in the expression of guarding behavior and the role of guards in the defensive response of honey bee colonies. *Apidologie*, 34, 439-447, 2003.
- 23. Guzman-Novoa E, Page RE:** Selective breeding of honeybees (Hymenoptera: Apidae) in Africanized areas. *J Econ Entomol*, 92, 521-525, 1999.
- 24. Akyol E, Şahinler N, Özkök D:** Honeybee (*Apis mellifera*) races, ecotypes and their general characteristics in Turkey. *J Anim Vet Adv*, 5 (9): 771-774, 2006.
- 25. Yücel B, Kösoğlu M:** Comparisons of Muğla ecotype and Italian cross honey bees for some performances in Aegean Region (Turkey). *Kafkas Univ Vet Fak Derg*, 17 (6): 1025-1029, 2011.
- 26. Garnery L, Franck P, Baudry E, Vautrin D, Cornuet JM, Solignac M:** Genetic diversity of the west European honey bee (*Apis mellifera mellifera* and *Apis m. iberica*), II Microsatellite loci. *Genet Sel Evol*, 30, 49-74, 1998.