

# The Effect of Hot-Iron Disbudding on Thiol-Disulphide Homeostasis in Calves

Hasan ERDOĞAN <sup>1,a</sup> İlker ÇAMKERTEN <sup>2,b</sup> Güzin ÇAMKERTEN <sup>3,c</sup> Kerem URAL <sup>1,d</sup>  
Songül ERDOĞAN <sup>1,e</sup> İsmail GÜNAL <sup>4,f</sup> Özcan EREL <sup>5,g</sup>

<sup>1</sup> Adnan Menderes University, Veterinary Faculty, Internal Medicine, TR-09016 Aydın - TURKEY

<sup>2</sup> Aksaray University, Veterinary Faculty, Internal Medicine, TR-68100 Aksaray - TURKEY

<sup>3</sup> Aksaray University, Veterinary Faculty, Biochemistry, TR-68100 Aksaray - TURKEY

<sup>4</sup> Harran University, Veterinary Faculty, Internal Medicine, TR-63000 Şanlıurfa - TURKEY

<sup>5</sup> Yıldırım Beyazıt University, Faculty of Medicine, Biochemistry, TR-06800 Ankara - TURKEY

<sup>a</sup> ORCID: 0000-0001-5141-5108; <sup>b</sup> ORCID: 0000-0002-6952-7703; <sup>c</sup> ORCID: 0000-0003-2732-9490; <sup>d</sup> ORCID: 0000-0003-1867-7143; <sup>e</sup> ORCID: 0000-0002-7833-5519; <sup>f</sup> ORCID: 0000-0003-3679-4132; <sup>g</sup> ORCID: 0000-0002-2996-3236

Article Code: KVFD-2018-20950 Received: 12.09.2018 Accepted: 26.08.2018 Published Online: 26.12.2018

## How to Cite This Article

Erdoğan H, Çamkerten İ, Çamkerten G, Ural K, Erdoğan S, Günal İ, Erel Ö: The effect of hot-iron disbudding on thiol-disulphide homeostasis in calves. *Kafkas Univ Vet Fak Derg*, 2018 (Article in Press). DOI: 10.9775/kvfd.2018.20950

## Abstract

We aimed to examine the effect of hot-iron disbudding on serum thiol-disulphide homeostasis levels as a marker of oxidant stress in relationship with trauma in calves under sedation, local anaesthesia, and the non-steroidal anti-inflammatory drug ketoprofen. A total of 30 Holstein calves were enrolled in the study and allocated into three groups: disbudded following sedation with xylazine (n=10) (group I); disbudded following sedation (xylazine) and local anaesthesia with lidocaine (n=10) (group II); and disbudded after sedation (xylazine), local anaesthesia (lidocaine), and ketoprofen (n=10) (group III). Blood samples were withdrawn before (0. min) and 30, 60, 90, and 120 min after dehorning. Serum native thiols, total thiols, and disulphide levels were detected with a novel assay. Native thiol and total thiol levels were reduced in all groups without any significance during the study period. At the 90th min of the study, native thiol levels in group II were significantly lower than in groups I and III. There were no significant alterations in total thiol levels in both groups. Disulphide levels showed no significant changes in group, time, and group by time interactions, but at the 60th min, groups I and III had the lowest levels. Disulphite/native thiol, disulphite/total thiol, and native thiol/total thiol levels had significant group alterations in the 60th min. The reduction of native thiol and total thiol levels in all groups without significance might be related to the antioxidant activity of plasma; however, it is thought that the pain management procedures should be related to the sensitive oxidative balance by thiols.

**Keywords:** Calf, Disbudding, Sedation, Thiol-disulphide

## Buzağlarda Sıcak Koter ile Boynuzsuzlaştırmanın Tiyol-Disülfid Homeostazı Üzerine Etkisi

### Öz

Bu çalışmada sedasyon, lokal anestezi ve non-steroidal bir ilaç olan ketoprofen uygulanmış sıcak koterizasyon işlemi ile boynuzsuzlaştırılan buzağlarda oksidatif stresin değerlendirilmesinde tiyol-disülfid homeostazına olan etkilerinin belirlenmesi amaçlandı. Çalışmaya toplam 30 adet Holstein buzağı dahil edilerek üç gruba ayrıldı: sedasyon (xylazine) işlemi takiben boynuzsuzlaştırılan (n=10) (grup I); sedasyon (xylazine) ve lidokain ile lokal anestezi işlemi sonrasında boynuzsuzlaştırılan (n=10) (grup II); ve sedasyon (xylazine), lokal anestezi (lidocaine) ve ketoprofen uygulaması sonrasında boynuzsuzlaştırılan (n=10) (grup III). Kan örnekleri, boynuzsuzlaştırmadan önce (0. dak) ve 30, 60, 90 ve 120 dakika sonra olacak şekilde toplandı. Serum natif tiyol, total tiyol ve disülfür seviyeleri yeni bir test ile belirlendi. Çalışma süresince natif tiyol ve total tiyol seviyelerinde istatistiksel anlamlı olmayan azalmalar belirlendi. Araştırmanın 90. dakikasında grup II'de bulunan hayvanların natif tiyol seviyelerinin grup I ve III'e göre istatistiksel anlamlı düşüktü. Total tiyol seviyelerinde ise gruplar arasında istatistiksel anlamlı farklılıkların olmadığı belirlendi. Disülfid seviyelerinde grup, zaman ve grup zaman arasında farklılıklar belirlenmezken, araştırmanın 60. dakikasında grup I ve III'te en düşük seviyelere geldiği belirlendi. Disülfid/natif tiyol, disülfid/total tiyol ve natif tiyol/total tiyol seviyelerinin 60. dakikada gruplar arasında istatistiksel anlamlı değişimler gösterdiği belirlendi. Natif tiyol ve total tiyol seviyelerindeki bu azalmaların plazmadaki antioksidan kapasite ile ilişkili olabileceği, ancak uygulanan ağrı yönetimi prosedürlerinin tiyol ile oluşturulan hassas oksidatif dengeyi etkilediği düşünüldü.

**Anahtar sözcükler:** Buzağı, Boynuzsuzlaştırma, Sedasyon, Tiyol-disülfid



### İletişim (Correspondence)



+90 256 2470700/372 Fax: + 90 256 2470720



hasan.erdogan@adu.edu.tr

## INTRODUCTION

Dehorning is one of the frequently applied practices in livestock and is depended on to keep animals safe from injuries. Dehorning is a stressful and painful process that results in many homeostatic changes in animals [1]. The disbudding procedure is another term for dehorning in calves up to 3 months of age. The disbudding procedure can be applied to calves in different ways, such as hot-iron disbudding, chemical disbudding with caustic pastes, and surgical disbudding using scoop dehorning. In calves up to 8 weeks of age, hot-iron disbudding can be used [2]. However, this provokes third-degree burns in the area where it is performed [3]. Along with hot-iron disbudding, inflammatory changes, severe pain, behavioural/physiological alterations, and acute stress responses exist in calves [2,4-6]. Numerous reports have described behavioural and physiological reactions to disbudding in calves [1,7-9]. Many inflammatory conditions cause an increase in oxidative stress mediators that are provoked by pro-inflammatory cytokines in castrated and dehorned calves [10,11]. Previous studies and the American Veterinary Medical Association indicate the essentiality of pain management by pharmacological agents. These studies point out several methodologies, such as local anaesthesia [4,12,13], non-steroidal anti-inflammatory drugs (NSAIDs) [13-15], and sedatives [16,17].

Oxidative balance is described as the equilibrium among free radical eradication and production. Cell damage initiated by free radicals is limited to oxidative balance, and the imbalanced free radical production derives oxidative stress. Thiol is a novel and substantial antioxidant used to eliminate reactive oxygen via non-enzymatic and enzymatic pathways [18,19]. The plasma thiol pool includes both low molecular weight thiols (e.g. glutathione, cysteine, and homocysteine) and protein thiols. Thiols have an antioxidant role in oxidation reactions by composing disulphide bonds. Dynamic thiol/disulphide homeostasis is crucial for detoxification, apoptosis, and the processes of controlling enzymatic reactions and signalling pathways. Many inflammatory conditions are initiated by anomalous thiol/disulphide levels [20-25]. Korkmaz et al. [9] described the alterations of oxidative stress parameters in calves and mature cows undergoing hot iron dehorning, and defined the amputation process [9,26]. To our knowledge, dynamic thiol/disulphide homeostasis has not been studied previously in calves. Therefore, in the present study, we aimed to evaluate the alterations of thiol/disulphide homeostasis in calves undergoing disbudding with different analgesia and anaesthesia procedures.

## MATERIAL and METHODS

### Experimental Design, Calves, and Treatments

The study included 30 Holstein calves from both sexes (17

male and 13 female) at 10 weeks of age (BW= 85.6±8.9 kg). All calves were assigned to individual pens 7 days prior to the study and weaned at 7 weeks of age. Calves were fed with ad libitum access to water and a calf starter during the entire period. Study procedures were approved by Local Animal Ethic Committee of Adnan Menderes University with a number of 2017-058.

Calves were randomly divided into three groups. Study groups were designed as the treatment procedure: group I (n=10) xylazine group (with disbudding under sedation with an intramuscular injection of xylazine [Xylazinbio®, Interhas, Czech Republic (dose of 0.25 mg/kg)]); group II (n=10) to those of calves administered xylazine and local anaesthesia [with disbudding under xylazine sedation and subcutaneous infiltration of 20 mg of lidocain (Adokain®, Sanovel, Turkey) for horn buds prior to disbudding]; and group III (n=10) received xylazine, local anaesthesia, and subcutaneous injection of meloxicam (Maxicam®, Sanovel, Turkey) with a dose of 0.5 mg/kg before dehorning. All treatment and disbudding procedures were performed by the same researcher. Furthermore, the local anaesthesia procedure was carried out with a ring block and corneal nerve block between the horn bud and lateral canthus of the eye. Calves were deprived from calf starter 12 h prior to the study (in an attempt to decrease the risk of bloat whereas personnel controlled the calves during reanimation).

### Sample Collection and Analysis

Peripheral blood specimens were withdrawn from *Vena jugularis* starting prior to the dehorning and after drug application at 0, 30, 60, 90, and 120 min in lithium heparinised tubes. Immediately after the blood samples were taken, plasma samples were removed using a portable centrifuge in the eppendorf tubes. Plasma samples were stored at -80°C, then moved to the laboratory and analysed.

Total thiol (-S-S- + -SH) includes native and reduced thiol. A novel automatic and spectrophotometric technique established by Erel and Neselioglu [27] was used to determine the thiol/disulphide concentrations. The principle of this method is based on the degradation of dynamic disulphide bonds (-S-S-) to functional thiol groups (-SH) with a sodium borohydride (NaBH<sub>4</sub>) solution. The remaining NaBH<sub>4</sub> residue was totally removed by formaldehyde. Thus, this inhibited extra reduction of 5,5'-dithiobis-2-nitrobenzoic acid (DTNB) along with any disulphide bonds resulting from the reaction with DTNB. The following reaction with the DTNB-modified Ellman reagent was used to detect the amount of total thiol. The disulphide levels were counted automatically as half of the quantity of total thiol and native thiol. Disulphide/total thiol percent ratios, disulphide/native thiol percent ratios, and disulphide/total thiol percent ratios were calculated from the measured disulphide, total thiol, and native thiol parameters.

### Statistical Analyses

All repeated measurements were tabulated as means and standard errors according to descriptive statics. Normality tests were confirmed using the Shapiro-Wilk test. Obtained data were evaluated using both parametric repeated measures of ANOVA and non-parametric and Kruskal Wallis tests for group, time, and group-time interactions. The SPSS 22.0 packet program was used for all tests and  $P < 0.05$  was considered significant.

## RESULTS

There was no statistical difference in native thiol levels in any of the calves in the study groups. Native thiol levels were determined to be statistically lower in group I and III compared to group II at the 90<sup>th</sup> min following disbudding. The native thiol levels decreased in a statistically insignificant manner regarding all the treatment groups depending on time. Total thiol levels were found to be lowest at the 60<sup>th</sup> min of the study, while no significant differences were observed in group, time or group-time interaction in any treatment group. The total thiol levels increased from the 60<sup>th</sup> min of administration to the 90<sup>th</sup> and 120<sup>th</sup> min, but were lower than values measured at min 0. In the disulphide levels, the differences in terms of group, time, and group-time interaction in the application groups were not statistically significant. The disulphide concentrations decreased to the lowest level in groups I and III at the 60<sup>th</sup> min of application but increased in group II compared to min 0. There was no statistical difference in terms of time and group-time interactions in any of the disulphite/native

thiol, disulphite/total thiol, and native thiol/total thiol levels. There were differences at the 60<sup>th</sup> min of application between all groups (Table 1).

## DISCUSSION

In livestock, the pain management of dehorning or disbudding procedures is an important animal welfare issue [28]. In addition to local anaesthesia, NSAID analgesia appears to be generally beneficial, but the lack of specific recommendations for analgesic protocols may reflect the diversity studied in the literature. It is the current recommendation of clinicians and veterinarians that local anaesthetic and NSAIDs in North America can be obtained in full compliance [29,30]. Different methodologies might be used for dehorning (e.g. surgical amputation, chemical methods, or cautery), but disbudding with a cautery is still the most preferred method by livestock producers in the United States, Canada, and North America [29-31]. For managing pain and cortisol spikes in calves after the cautery disbudding process, local anaesthetic agents, non-steroidal anti-inflammatory drugs, and sedatives are used together or solely [30]. Stock et al. [1] reported that the suppression of increases in cortisol levels reduced the pain-related inflammatory response. In our study, the calves undergoing the disbudding process were divided into groups based on commonly used pain management methods.

In a study that evaluated serum oxidant and antioxidant status, the concentrations of nitric oxide (NO) and malondialdehyde (MDA) levels did not reveal any difference

**Table 1.** Time-dependent disulfide concentrations pursuant to treatment groups

Parameter	Group	0. min	30. min	60. min	90. min	120. min	Interactions	P value
Native thiol (µmol/L)	Group I	273.4±13.2	271.5±11.7	249.1±19.6	254.1±16.8	247.6±9.3	Group	0.028
	Group II	240.0±11.6	230.0±11.9	221.4±16.9	224.7±9.2	232.7±14.4	Time	0.211
	Group III	269.6±8.1	254.4±5.4	260.2±2.7	263.3±2.9	251.7±13.2	Group & time	0.779
Total thiol (µmol/L)	Group I	317.2±12.7	318.3±14.8	284.1±24.0	299.4±19.2	290.3±12.2	Group	0.134
	Group II	285.4±14.8	274.6±17.2	269.7±19.1	268.8±8.1	281.6±11.7	Time	0.168
	Group III	313.5±7.2	289.2±9.6	287.8±4.2	292.5±2.8	294.1±16.5	Group & time	0.823
Disulphide (µmol/L)	Group I	21.9±3.7	23.4±3.4	17.5±4.2	22.5±3.3	21.3±2.8	Group	0.106
	Group II	22.7±2.3	22.3±3.0	24.2±2.8	22.0±2.8	24.4±2.1	Time	0.271
	Group III	22.0±1.6	17.4±2.7	13.8±2.1	14.6±1.5	21.2±2.4	Group & time	0.451
Disulphide/native thiol (%)	Group I	7.4±1.4	8.6±1.3	6.9±1.6	9.1±1.3	8.6±1.1	Group	0.014
	Group II	9.4±0.8	9.5±1.0	11.2±1.4	10.1±1.6	11.0±1.5	Time	0.435
	Group III	8.2±0.8	6.8±1.0	5.3±0.8	5.6±0.6	8.4±0.7	Group & time	0.284
Disulphide/total thiol (%)	Group I	6.7±1.1	7.3±0.9	5.8±1.2	7.5±0.9	7.3±0.8	Group	0.017
	Group II	7.9±0.6	7.9±0.7	9.0±1.0	8.2±1.9	8.9±1.0	Time	0.342
	Group III	7.0±0.6	5.9±0.8	4.8±0.7	5.0±0.5	7.1±0.5	Group & time	0.312
Native thiol/total thiol (%)	Group I	86.3±2.3	85.5±1.9	88.4±2.5	84.8±1.8	85.5±1.6	Group	0.019
	Group II	84.2±1.1	84.2±1.5	82.0±1.9	83.6±2.2	82.3±2.0	Time	0.342
	Group III	85.9±1.1	88.2±1.6	90.5±1.3	90.1±1.0	85.7±1.0	Group & time	0.285

between control and Dex (dexketoprofen trometamol) groups. In addition, glutathione (GSH) significantly increased at 15 min after disbudding in the Dex group, and total antioxidant activity did not show any difference between groups. There were no significant alterations in the examined parameters in both groups during the study period<sup>[9]</sup>. Among the adverse consequences of stress, oxidative stress, which is characterised by the accumulation of radical oxygen species (ROS), can affect life<sup>[32]</sup>.

The thiol-disulphide homeostasis situation has important responsibilities in antioxidant protection, apoptosis, signal transduction, detoxification, regulation of enzymatic activity, and cellular signalling mechanisms<sup>[20,21]</sup>. Furthermore, thiol-disulphide homeostasis is increasingly being evaluated in various medical conditions in humans, such as diabetes<sup>[22]</sup>, rheumatoid arthritis<sup>[23]</sup>, cancer<sup>[25]</sup>, multiple sclerosis<sup>[33]</sup>, hepatic disorders<sup>[24]</sup>, and surgery<sup>[34]</sup>. The plasma thiol pool is constituted by cysteine, GSH, homocysteine, and albumin. In expanded oxidative stress conditions, thiol concentrations are decreased to compensate for the reactive oxygen radicals, wherein the sulfhydryl groups of the thiols play an important role<sup>[35]</sup>. Under oxidative stress conditions, thiol molecules engage disulphide bonds, which are reduced back to thiols to tolerate thiol/disulphide homeostasis<sup>[36]</sup>. In our study, thiol-disulphide homeostasis was evaluated in calves undergoing the dehorning process with different pain management regimens. Native thiol and total thiol concentrations were decreased in all groups of calves during the study period. Native thiol concentrations were found to be significantly different at the 90<sup>th</sup> min of the study period between group II and both groups I and III. In contrast, total thiol levels showed no significant alterations in group, time, or group-time interactions, but the lowest levels were examined at the 60<sup>th</sup> min of the study. This might be explained by thiols' negative reduction properties as electron acceptors. Thiol groups interact with oxidants and are neutralised to a less toxic product called disulphide. Total thiol and native thiol concentrations have been shown to be reduced by oxidation<sup>[27]</sup>. In our study, another interesting finding was detected in disulphide levels. There were no significant alterations in disulphide concentrations in both groups. In the 60<sup>th</sup> min of the study, the disulphide concentrations reached minimum levels in groups I and III. However, in group II, the disulphide concentrations were greater than at the beginning of the study. Reductions of thiol concentrations without a rise in disulphide levels might be the outcome of inadequate intake or increased devastation because of its use in other syntheses instead of conversion to disulphide<sup>[37]</sup>. The calculated parameters of disulphite/native thiol, disulphite/total thiol, and native thiol/total thiol levels were significant in group interactions at the 60<sup>th</sup> min of the study. In this study, decreases in native thiol and total thiol concentrations without increases in disulphide levels might be related to nutritional factors instead of oxidative stress. Furthermore, oxidative stress might be

suppressed by pain management strategies. This study is thought to be limited by the fact that pain management cannot be measured by cortisol levels.

In conclusion, to the best of our knowledge, this is the first study to examine thiol disulphide homeostasis in calves undergoing dehorning. Examining thiol/disulphide homeostasis during dehorning with different pain management procedures might be beneficial as an early evaluation test to recognise the best strategies in calves. Further studies are warranted to understand the association between oxidative stress and dehorning.

## DECLARATION OF CONFLICTING INTEREST

All authors have declared to be any financial and personal contest effected this study by other people or organizations.

## REFERENCES

- Stock ML, Baldrige SL, Griffin D, Coetzee JF:** Bovine dehorning: Assessing pain and providing analgesic management. *Vet Clin North Am Food Anim Pract*, 29 (1): 103-133, 2013. DOI: 10.1016/j.cvfa.2012.11.001
- Stafford KJ, Mellor DJ:** Dehorning and disbudding distress and its alleviation in calves. *Vet J*, 169 (3): 337-349, 2005. DOI: 10.1016/j.tvjl.2004.02.005
- Taschke AC, Fölsch DW:** Ethological, physiological and histological aspects of pain and stress in cattle when being dehorned. *Tierarztl Prax*, 25 (1): 19-27, 1997.
- McMeekan CM, Stafford KJ, Mellor DJ, Bruce RA, Ward RN, Gregory NG:** Effects of regional analgesia and/or a non-steroidal anti-inflammatory analgesic on the acute cortisol response to dehorning in calves. *Res Vet Sci*, 64 (2): 147-150, 1998. DOI: 10.1016/S0034-5288(98)90010-8
- Weary DM, Niel L, Flower FC, Fraser D:** Identifying and preventing pain in animals. *Appl Anim Behav Sci*, 100 (1-2): 64-76, 2006. DOI: 10.1016/j.applanim.2006.04.013
- Doherty TJ, Kattesh HG, Adcock RJ, Welborn MG, Saxton AM, Morrow JL, Dailey JW:** Effects of a concentrated lidocaine solution on the acute phase stress response to dehorning in dairy calves. *J Dairy Sci*, 90 (9): 4232-4239, 2007. DOI: 10.3168/jds.2007-0080
- Molony V, Kent JE, Robertson IS:** Assessment of acute and chronic pain after different methods of castration of calves. *Appl Anim Behav Sci*, 46 (1-2): 33-48, 1995. DOI: 10.1016/0168-1591(95)00635-4
- Caray D, Des Roches AD, Frouja S, Andanson S, Veissier I:** Hot-iron disbudding: stress responses and behavior of 1-and 4-week-old calves receiving anti-inflammatory analgesia without or with sedation using xylazine. *Livest Sci*, 179, 22-28, 2015. DOI: 10.1016/j.livsci.2015.05.013
- Korkmaz M, Sarıtaş ZK, Bülbül A, Demirkan I:** Effect of pre-emptive dexketoprofen trometamol on acute cortisol, inflammatory response and oxidative stress to hot-iron disbudding in calves. *Kafkas Univ Vet Fak Derg*, 21 (4): 563-568, 2015. DOI: 10.9775/kvfd.2015.12963
- Earley B, Crowe MA:** Effects of ketoprofen alone or in combination with local anesthesia during the castration of bull calves on plasma cortisol, immunological, and inflammatory responses. *J Anim Sci*, 80 (4): 1044-1052, 2002. DOI: 10.2527/2002.8041044x
- Ting STL, Earley B, Hughes JML, Crowe MA:** Effect of ketoprofen, lidocaine local anesthesia, and combined xylazine and lidocaine caudal epidural anesthesia during castration of beef cattle on stress responses, immunity, growth, and behavior. *J Anim Sci*, 81, 1281-1293, 2003. DOI: 10.2527/2003.8151281x
- McMeekan C, Stafford KJ, Mellor DJ, Bruce RA, Ward RN, Gregory N:** Effects of a local anaesthetic and a non-steroidal anti-inflammatory



analgesic on the behavioural responses of calves to dehorning. *N Z Vet J*, 47 (3): 92-96, 1999. DOI: 10.1080/00480169.1999.36120

**13. Duffield TF, Heinrich A, Millman ST, DeHaan A, James S, Lissemore K:** Reduction in pain response by combined use of local lidocaine anesthesia and systemic ketoprofen in dairy calves dehorned by heat cauterization. *Can Vet J*, 51 (3): 283-288, 2010.

**14. Heinrich A, Duffield TF, Lissemore KD, Millman ST:** The effect of meloxicam on behavior and pain sensitivity of dairy calves following cautery dehorning with a local anesthetic. *J Dairy Sci*, 93 (6): 2450-2457, 2010. DOI: 10.3168/jds.2009-2813

**15. Stilwell G, Lima MS, Carvalho RC, Broom D:** Effects of hot-iron disbudding using regional anesthesia with and without carprofen, on cortisol and behavior of calves. *Res Vet Sci*, 92, 338-341, 2012. DOI: 10.1016/j.rvsc.2011.02.005

**16. Mintline EM, Stewart M, Rogers AR, Cox NR, Verkerk GA, Stookey JM, Tucker CB:** Play behavior as an indicator of animal welfare: Disbudding in dairy calves. *Appl Anim Behav Sci*, 144 (1-2): 22-30, 2013. DOI: 10.1016/j.applanim.2012.12.008

**17. Stafford KJ, Mellor DJ:** Addressing the pain associated with disbudding and dehorning in cattle. *Appl Anim Behav Sci*, 135, 226-231, 2011. DOI: 10.1016/j.applanim.2011.10.018

**18. Cadenas E:** Biochemistry of oxygen toxicity. *Annu Rev Biochem*, 58, 79-110, 1989. DOI: 10.1146/annurev.bi.58.070189.000455

**19. Young IS, Woodside JV:** Antioxidants in health and disease. *J Clin Pathol*, 54 (3): 176-186, 2001. DOI: 10.1136/jcp.54.3.176

**20. Biswas S, Chida AS, Rahman I:** Redox modifications of protein-thiols: Emerging roles in cell signaling. *Biochem Pharmacol*, 71 (5): 551-564, 2006. DOI: 10.1016/j.bcp.2005.10.044

**21. Circu ML, Aw TY:** Reactive oxygen species, cellular redox systems, and apoptosis. *Free Radic Biol Med*, 48 (6): 749-762, 2010. DOI: 10.1016/j.freeradbiomed.2009.12.022

**22. Matteucci E, Giampietro O:** Thiol signalling network with an eye to diabetes. *Molecules*, 15 (12): 8890-8903, 2010. DOI: 10.3390/molecules15128890

**23. Tetik S, Ahmad S, Alturfan AA, Fresko I, Disbudak M, Sahin Y, Aksoy H, Yardimci KT:** Determination of oxidant stress in plasma of rheumatoid arthritis and primary osteoarthritis patients. *Indian J Biochem Biophys*, 47, 353-358, 2010.

**24. Kuo LM, Kuo CY, Lin CY, Hung MF, Shen JJ, Hwang TL:** Intracellular glutathione depletion by oridonin leads to apoptosis in hepatic stellate cells. *Molecules*, 19 (3): 3327-3344, 2014. DOI: 10.3390/molecules19033327

**25. Prabhu A, Sarcar B, Kahali S, Yuan Z, Johnson JJ, Adam KP, Chinnaiyan P:** Cysteine catabolism: A novel metabolic pathway contributing to glioblastoma growth. *Cancer Res*, 74 (3): 787-796, 2013.

DOI: 10.1158/0008-5472.CAN-13-1423

**26. Fidan AF, Pamuk K, Ozdemir A, Saritas ZK, Tarakci U:** Effects of dehorning by amputation on oxidant/antioxidant status in mature cattle. *Rev Med Vet*, 161 (11): 502-508, 2010.

**27. Erel O, Neselioglu S:** A novel and automated assay for thiol/disulphide homeostasis. *Clin Biochem*, 47 (18): 326-332, 2014. DOI: 10.1016/j.clinbiochem.2014.09.026

**28. Ventura BA, von Keyserlingk MAG, Weary DM:** Animal welfare concerns and values of stakeholders within the dairy industry. *J Agric Environ Ethics*, 28, 109-126, 2015. DOI: 10.1007/s10806-014-9523-x

**29. Adams AE, Lombard JE, Shivley CS, Urie NJ, Roman-Muniz IN, Fossler CP, Koprak CA:** Management practices that may impact dairy heifer welfare on US dairy operations. *J Dairy Sci*, 98 (2): 105, 2015.

**30. Winder CB, LeBlanc SJ, Haley DB, Lissemore KD, Godkin MA, Duffield TF:** Practices for the disbudding and dehorning of dairy calves by veterinarians and dairy producers in Ontario, Canada. *J Dairy Sci*, 99 (12): 10161-10173, 2016. DOI: 10.3168/jds.2016-11270

**31. Vasseur E, Borderas F, Cue RI, Lefebvre D, Pellerin D, Rushen J, Wade KM, De Passillé AM:** A survey of dairy calf management practices in Canada that affect animal welfare. *J Dairy Sci*, 93, 1307-1315, 2010. DOI: 10.3168/jds.2009-2429

**32. Chirase NK, Greene LW, Purdy CW, Loan RW, Auvermann BW, Parker DB, Walborg EF, Stevenson DE, Xu Y, Klaunig JE:** Effect of transport stress on respiratory disease, serum antioxidant status, and serum concentrations of lipid peroxidation biomarkers in beef cattle. *Am J Vet Res*, 65 (6): 860-864, 2004. DOI: 10.2460/ajvr.2004.65.860

**33. Calabrese M, Magliozzi R, Ciccarelli O, Geurts JGG, Reynolds R, Martin R:** Exploring the origins of grey matter damage in multiple sclerosis. *Nat Rev Neurosci*, 16 (3): 147-158, 2015. DOI: 10.1038/nrn3900

**34. Polat M, Ozcan O, Sahan L, Üstündag Budak Y, Alisik M, Yilmaz N, Erel Ö:** Changes in thiol-disulfide homeostasis of the body to surgical trauma in laparoscopic cholecystectomy patients. *J Laparoendosc Adv Surg Tech A*, 26 (12): 992-996, 2016. DOI: 10.1089/lap.2016.0381

**35. Erkus ME, Altiparmak IH, Demirbag R, Gunebakmaz O, Kaya Z, Taskin A, Neselioglu S, Erel O:** The investigation of the dynamic thiol-disulfide homeostasis in acute coronary syndromes. *Am J Cardiol*, 115 (Suppl 1), S117-S118, 2015. DOI: 10.1016/j.amjcard.2015.01.401

**36. Jones DP, Liang Y:** Measuring the poise of thiol/disulfide couples *in vivo*. *Free Radic Biol Med*, 47 (10): 1329-1338, 2009. DOI: 10.1016/j.freeradbiomed.2009.08.021

**37. Durrieu G, LLau ME, Rascol O, Senard JM, Rascol A, Montastruc JL:** Parkinson's disease and weight loss: A study with anthropometric and nutritional assessment. *Clin Auton Res*, 2, 153-157, 1992. DOI: 10.1007/BF01818955