

Amelioration Effects of Vitamin E, Melatonin, L-carnitine, and Atorvastatin, on Destructive Effects of Busulfan in the Testes of Male Rats: A Gene Expression Evaluation

Forutan SALEHINEZHAD¹ Hamidreza ESHRAGHI¹ Ali KADIVAR^{2,3}
Sadegh SHIRIAN^{4,5} Ahmad ASGHARI⁶ Ehsan AALI⁷ Najmeh DAVOODIAN³

¹ Department of Veterinary Basic Science, Science and Research Branch, Islamic Azad University, Tehran, IRAN

² Department of Clinical Science, Faculty of Veterinary Medicine, Shahrekord University, Shahrekord, IRAN

³ Research Institute of Animal Embryo Technology, Shahrekord University, Shahrekord, IRAN

⁴ Department of Pathology, Faculty of Veterinary Medicine, Shahrekord University, Shahrekord, IRAN

⁵ Biotechnology Research Institute, Shahrekord University, Shahrekord, IRAN

⁶ Department of Veterinary Clinical Science, Science and Research Branch, Islamic Azad University, Tehran, IRAN

⁷ Department of Pharmacology, School of Medicine, Qazvin University of Medical Sciences, Qazvin, IRAN

Article ID: KVFD-2019-21726 Received: 11.01.2019 Accepted: 06.05.2019 Published Online: 06.05.2019

How to Cite This Article

Salehinezhad F, Eshraghi H, Kadivar A, Shirian S, Asghari A, Aali E, Davoodian N: Amelioration effects of vitamin E, melatonin, L-carnitine, and atorvastatin, on destructive effects of busulfan in the testes of male rats: A gene expression evaluation. *Kafkas Univ Vet Fak Derg*, 25 (5): 709-716, 2019. DOI: 10.9775/kvfd.2019.21726

Abstract

According to toxicity of various types of cancer treatments on different kind of cells with high division activities such as germ cells, antioxidants may protect these cells in testes against the toxic effects of the chemotherapeutic drugs. For this purpose, 24 h after busulfan treatment, 30 adult male wistar-rats were divided to six groups. Intra-peritoneally administrations of normal saline in control group and DMSO (as a busulfan solvent) in DMSO group were performed daily for 6 weeks beside the treatment contain vitamin E (Vit-E group), L-carnitine and melatonin (LM group), atorvastatin and melatonin (AM group), atorvastatin, L-carnitine, and melatonin (ALM group). After decapitation and removal of the testes, molecular evaluations were performed by the relative abundance measurement of *DAZL*, *Bcl2*, and *Casp3* transcripts. The results of this study exhibited high level of expression of *DAZL* in Vit-E treated rats compared to control counterparts ($P < 0.01$). The expression level of *Bcl2* is significantly down-regulated in LM ($P < 0.008$), and ALM groups ($P < 0.001$), and the relative abundance of *Casp3* transcripts was significantly lower in AM ($P < 0.001$) and ALM ($P < 0.007$) than that of control group. As well as, there was significant high expression of this gene in Vit E-treated rats compared to the rats of control group. In conclusion, busulfan destructive effects were moderated with Vit-E administration through regulation of the expression of *DAZL*. The other antioxidants used in different combinations had not amelioration effects on spermatogenesis in busulfan-induced male rats, though the positive effects of some of these antioxidants on apoptosis reduction.

Keywords: Rat, Busulfan, Vitamin E, Melatonin, L-carnitine, Atorvastatin, *DAZL*, *Bcl2*, *Casp3*

Erkek Rat Testislerinde Busulfan Kaynaklı Hasara Karşı Vitamin E, Melatonin, L-karnitin ve Atorvastatin'in Koruyucu Etkisi: Gen Ekspresyonunun Değerlendirilmesi

Öz

Çeşitli kanser ilaçlarının toksisitelerine bağlı olarak eşey hücreleri gibi yüksek bölünme kapasitesine sahip çeşitli hücrelerde meydana gelen hasara karşı antioksidanlar koruyucu olabilir ve bu durum testis dokusunda kemoterapötik ilaçlarla oluşan hasarda etkili olabilir. Bu amaçla, 24 saat busulfan uygulaması sonucunda 30 ergin erkek Wistar rat altı gruba ayrıldı. İntraperitoneal olarak 6 hafta süresince DMSO grubuna fizyolojik tuzlu su ve DMSO (busulfan çözücüsü olarak) uygulandı. Diğer gruplarda busulfan uygulamasına ilave olarak E (Vit-E grubu), L-karnitin ve melatonin (LM grubu), atorvastatin ve melatonin (AM grubu), atorvastatin, L-karnitin ve melatonin (ALM grubu) uygulamaları gerçekleştirildi. Hayvanlarda dekapitasyonu takiben testis dokuları alındı ve *DAZL*, *Bcl2*, ve *Casp3* miktarları karşılaştırmalı olarak moleküler açıdan değerlendirildi. Çalışma sonucu, Vit-E uygulanan ratlarda *DAZL* ekspresyonunun kontrol grubuna oranla daha fazla olduğu gözlemlendi ($P < 0.01$). LM ($P < 0.008$) ve ALM ($P < 0.001$) gruplarında *Bcl2* ekspresyonu aşağı yönde regüle edilirken, AM ($P < 0.001$) ve ALM ($P < 0.007$) gruplarında *Casp3* ekspresyonu anlamlı derecede kontrol grubuna oranla daha azdı. Kontrol ile karşılaştırıldığında Vit-E uygulanan grupta *Casp3* ekspresyonunun anlamlı derecede yüksek olduğu belirlendi. Sonuç olarak, busulfan kaynaklı hasarın Vit-E uygulaması sonucu *DAZL* ekspresyonu yolu ile normalize edildiği tespit edildi. Busulfan ile erkek ratlarda spermatogenez üzerine oluşturulan hasara karşı çalışmada uygulanan diğer antioksidanların hasarı azaltmada etkili olmadığı ancak bazılarının apoptozis üzerine etkisinin olduğu belirlendi.

Anahtar sözcükler: Rat, Busulfan, Vitamin E, Melatonin, L-karnitin, Atorvastatin, *DAZL*, *Bcl2*, *Casp3*



İletişim (Correspondence)



+98 917 7323202



kadivar.ali@sku.ac.ir

INTRODUCTION

Alkylating agents as chemotherapeutic drugs are often used to treat cancer and increase the survival rates of patients. Alkylating agents, affect cell division by adhesion to one strands of the DNA^[1]. Therefore, cells or tissues with high division activities such as germ cells and testes are more susceptible to these agents' side effects. Spermatogenesis makes round haploid spermatids through the reductive divisions of meiosis^[2]. It is clear that certain chemotherapeutic drugs specially alkylating agents influence spermatogenesis at least temporarily and in some cases permanently. Single doses of busulfan as an alkylating agent can permanently sterilize rats at non-lethal doses and cause long-term morphological damage to sperm produced by surviving spermatogonia^[3]. Busulfan toxicity occurs by several different mechanisms, including reactive oxygen species (ROS) formation and protein damage (oxidation)^[4]. ROS are directly involved in precarious oxidative damage of cellular macromolecules such as proteins and nucleic acids in germ cells, which can lead to cell death^[5]. Indeed, the overproduction of ROS and the consequent oxidative stress has a critical role in inhibition of development of germ cells. Forasmuch as, the rats treated with Busulfan exhibited a defined increase in apoptosis, using cytoprotective and anti-apoptotic agents such as antioxidants can neutralize both ROS and apoptosis procedure caused by them^[6]. ROS are scavenged by some antioxidants such as L-carnitine, Vitamin E (α -tocopherol), and Atorvastatin via their interfering with the lipid peroxidation chain reaction. Another small molecule antioxidant, Melatonin, which can detoxify highly reactive hydroxyl free radicals (OH) is involved in the detoxification of free radicals^[7]. Co-administration of these components also change produced ROS profile, prevented lipid peroxidation and improved antioxidant status, synergistically^[8]. Therefore, strategies that effectively preserve fertility during the course of cancer treatment especially through the decrease in apoptosis should be developed.

In this study all mentioned antioxidants, singly or in different combinations, were used to amelioration busulfan side effects in treated male rats and then the expression amount of deleted in azoospermia-like (*DAZL*) as a gene related to spermatogenesis resumption and B-cell lymphoma 2 (*Bcl2*) and Caspase 3 (*Casp3*) as the genes related to apoptosis was measured.

DAZL is a gene cluster which gets deletions in at least 10% of males with oligozoospermia or azoospermia^[9]. This gene is a DAZ autosomal homologue. As a result, *DAZL* has always been seen as a promising candidate for male infertility. *DAZL* plays an important role in the spermatogenic processes.

Bcl2 gene is an anti-apoptotic member of *Bcl2* Family, regulators of the cellular life-or-death switch, that prevents

cytochrome C release, and hence caspase-9 activation and subsequently several other caspases, independently of mitochondrial damage^[10]. Besides, caspase-3 is a cysteine protease that is activated early in a sequence of events associated with apoptosis^[11].

The aim of the present study was reduction the amount of apoptosis and subsequently improves the spermatogenesis and preserves fertility of azoospermic animal model using busulfan in rat, with a focus on related molecular pathways.

MATERIAL and METHODS

Animals

Thirty adult male wistar-Rats (2-3 months old and 200-250 g) were purchased from Faculty of Veterinary Medicine of Shahrekord University and were housed for 2 weeks at the animal lab of Veterinary Clinic under the Standard laboratory conditions (12 h dark and 12 h light cycle, temperature of $23\pm 3^{\circ}\text{C}$, and $50\pm 5\%$ humidity) for adaptability of rats to new living environment. Animal cages were kept clean, and commercial food (pellet) and water were provided *ad libitum*.

Experimental Design

In total, all rats were treated with two doses of 25 and 10 mg/kg busulfan with 14 days interval and after 24 h were randomly assigned to the following groups: Sham group that treated with DMSO as busulfan solvent, treated with busulfan (control group), and treatment groups including: treated with 100 mg/kg vitamin E (Vit E group), treated with 100 mg/kg L-carnitine and 1 mg/kg melatonin (LM group), treated with 20 mg/kg atorvastatin and 1 mg/kg melatonin (AM group), treated with 20 mg/kg atorvastatin, 100 mg/kg L-carnitine, and 1 mg/kg Melatonin (ALM group). Each group assigned by 5 rats. In all groups all administrations were performed intra-peritoneally daily for 6 weeks. Busulfan dose was assigned based on previous studies that established the toxic effect of busulfan on rat testes^[12]. The atorvastatin^[13], L-carnitine^[13], melatonin^[14] and Vit E^[15] doses were also selected based on previous reports demonstrating their anti-oxidative effect. This study was approved by the Institutional Ethics Committee for Animal Experimentation and was conducted in accordance with the international guidelines^[16].

Organ Removal and Tissue Processing

Animals were killed by decapitation under ether anesthesia, and testes of the animals were removed. One testis was used for sperm collection and the other one was preserved in -80°C freezer until molecular evaluations.

Sperm Collection and Evaluation

Sperm was collected from tail of left epididymis for all the

experimental rats. The left epididymis was immediately minced with scalpels and placed in pre-warmed microtube, containing 1 mL of Human Tubal Fluid (HTF) medium without BSA and placed in a 37°C incubator for 10 min. For sperm counting, 50 µL of the solution was diluted 10 times with 0.9% saline and a drop was transferred into chamber of Neubauer hemocytometer. Sperm counted under a standard optical microscope in order to determine the number of spermatozoa.

After 10 min of incubation, sperm motility was assessed by putting one drop of the on a warmed microscope slide. A cover slip was placed and at least 5 microscopic fields was observed at 400-fold magnifications. The percentage of nonmotile sperm was recorded for each rat.

Sperm viability was assessed with eosin-nigrosin staining. For this, 10 µL of sperm suspension were mixed with equal volume of eosin-nigrosin on a warmed microscope slide and a thin smear was prepared. Samples were observed under light microscope at a magnification of 1000×. At least 100 sperm were evaluated to discriminate death sperm (red stained) of live (not stained).

Molecular Evaluation

Total RNA Extraction, DNAase Treatment and cDNA Synthesis: To extract total RNA, tissues of testes (100 mg) were mechanical fragmented with a scalpel. Half mL RNX-PLUS solution (Sinaclon Bioscience, Karaj, Iran) containing phenol and guanidine was added and placed at room temperature for 5 min to homogenized samples. After adding 120 µL chloroform, the mixture was centrifuged at 8.000 g and 4°C for 5 min. The upper aqueous phase of supernatant was separated, and after addition of 400 µL isopropanol (100%), it was centrifuged (8000 g, 4°C, 5 min) and the RNA pellet was washed with 80% ethanol. Upon centrifugation at the same condition, the pellet was suspended in DEPC-treated water. To remove genomic DNA contamination, the extracted RNA was treated by RNase-free DNase I and its buffer (Sinaclon Bioscience, Karaj, Iran) and incubated at 37°C for 30 min. The reaction was stopped by EDTA at 65°C for 10 min, and the amount and quality of RNA were determined by spectrophotometry (Micro-volume Spectrophotometer System, Nano Mabna Iranian, Tehran, Iran). Only RNA of sufficient purity, having

an absorbance ratio (A260/280) between 1.8 and 2.2 was considered for synthesis of cDNA.

Shortly after extraction, total RNA (1 µg) was reversely transcribed into cDNA (less than 2 h) in a 5 µL reaction volume using the Easy cDNA synthesis kit, offered by the manufacturer (Pars tous biotechnology, Mashhad, Iran). The thermal program for cDNA synthesis included the following 3 steps: 25°C for 10 min (activation of the reverse transcriptase), 47°C for 60 min (reverse transcription), and 85°C for 5 min (inactivation of the reverse transcriptase). The synthesized cDNA was then stored at -20°C.

Quantitative Real-Time PCR: Real-time PCR was performed in two replicates for each sample (Rotor Gene Q 6000, Qiagen, USA). Primer sequences, the GenBank accession numbers, the size of amplified products, and annealing temperature of each primer are shown in Table 1. Half µL DNase I treated cDNA was added to 5 µL SYBR Premix Ex Taq II Mix and 0.5 µM of each specific primer in a total volume of 10 µL. The PCR program was comprised of 45 cycles of 94°C for 5 s (denaturation), and 54-60°C for 30 s (annealing & extension; Table 1) and 72°C for 30 s.

Considering the selection of an appropriate housekeeping gene as a reference gene for normalization, there are several studies demonstrating that *Actb* is highly reliable reference genes among the other genes used for RT-qPCR normalization and analysis of relative gene expression in the mouse testes [17].

Melt curve analysis was conducted to confirm the specificity of each product. The no-template control and no-reverse transcriptase control were considered to check contamination of the PCR reagents. Data were analyzed using LinReg PCR software version 2012.0 (USA), to give the threshold cycle number (Ct). Mean efficiency values (E) for each gene were also determined from the amplification profiles of individual samples using the same software [7]. The following formula was applied to determine the relative gene expression in tissue testes of treated rat compared to the control group [18,19].

Statistical Analysis

The differences in relative abundance of gene expression

Table 1. Sequence and annealing temperature of primers used for Real Time-PCR

Gene	Gene Bank Accession No.	Primer Sequence (5'-3')	Product Size (bp)	Annealing Temp (°C)	Reference
<i>Casp3</i>	NM_012922.2	F-GGACCTGTGGACCTGAAAAA R-GCATGCCATATCATCGTCAG	159	54	[20]
<i>Bcl2</i>	NM_016993.1	F-GGTGAACCTGGGGGAGGATTG R-GCATGCTGGGGCCATATAGT	197	60	[20]
<i>DAZL</i>	NM_001109414.1	F-TCTTCATCAGCAACCACCAG R-GACAAATCCATAGCCCTTCG	195	60	[12]
<i>B-Actin (Actb)</i>	NM_031144.3	F- CACCCGCGAGTACAACCTTC R- GAAGCCGGCCTTGACAT	127	60	Designed by writers

between groups were analyzed using one sample t-test student after ArcSin transformation with SPSS software version 20.0.0 (IBM Corp.; USA). Data were expressed as mean±SEM. Differences were considered significant at P<0.05.

RESULTS

The results of sperm evaluation showed that the mean percent of live and non-motile sperm and the sperm concentration were not different significantly between treatment groups. The difference was only significant between control and sham for all three parameters (Table 2).

The results of gene expression were presented by relative comparison of all of treatments groups with control group.

The relative abundance (RA) of transcripts in tissue of treated rat testes has been shown in Fig. 1A-B and Fig. 2. As shown in Fig. 1A, the expression level of *Bcl2* is significantly down regulated in those rats received both L-carnitine and melatonin (LM) (P<0.008), and those that received all three antioxidants, Atorvastatin, L-carnitine and melatonin (ALM) (P<0.001) in comparison with control group. No significant difference was observed between each of other treatment groups and control group in *Bcl2* expression.

According to Fig. 1B, the relative abundance of *Casp3* transcripts was significantly lower in AM (P<0.001) and ALM (P<0.007) than control group. As well as, there was significant high expression of this gene in Vit E-treated rats compared to the rats of control group. There was no significant difference between each of other groups (DMSO, and LM) and control group.

Table 2. The mean±SE percent of live and non-motile sperm and the sperm concentration

Sperm Characteristics in Treatment Groups		N	Mean	95% Confidence Interval for Mean	
				Lower Bound	Upper Bound
Live sperm (%)	Sham	8	67±1.52 ^a	63.40	70.60
	ALM	8	52±1.92 ^{ab}	47.46	56.54
	vitamin E	8	50.25±2.6 ^{ab}	44.09	56.41
	Control	8	24.75±2.23 ^b	19.47	30.03
	Total	32	48.5±2.9	42.58	54.42
Sperm count	Sham	8	564.38±15.63 ^a	527.40	601.35
	ALM	8	461.63±18.78 ^{ab}	417.21	506.04
	vitamin E	8	424.38±11.7 ^{ab}	389.93	458.82
	Control	8	225±23.28 ^b	197.35	252.65
	Total	32	418.84±131.72	371.35	466.34
Non-motile sperm	Sham	8	18.88±1.31 ^a	15.76	21.99
	ALM	8	28±1.45 ^{ab}	24.57	31.43
	vitamin E	8	31.13±1.7 ^{ab}	27.09	35.16
	Control	8	34.5±3.15 ^b	27.05	41.95
	Total	32	28.13±1.42	25.22	31.03

ALM: atorvastatin, L-carnitine, and melatonin group, Sham: treated with DMSO as busulfan solvent




Fig 1. Relative abundance of *Bcl2* (A) and *Casp3* (B) transcripts in rat testes tissue derived from treated rats with busulfan solvent (DMSO), Vitamin E (Vit E), L-carnitine and Melatonin (LM), atorvastatin and melatonin (AM), atorvastatin, L-carnitine, and Melatonin (ALM) compared to control group (busulfan treated rats). All reactions were normalized for β -Actin mRNA expression. Values with superscripts "*" refers to significant (P<0.05) differences in relative transcript abundance of each group compared to control group




Fig 2. Relative abundance of DAZL transcripts in rat testes tissue derived from treated rats with busulfan solvent (DMSO), Vitamin E (Vit E), L-carnitine and Melatonin (LM), atorvastatin and melatonin (AM), atorvastatin, L-carnitine, and Melatonin (ALM) compared to control group (busulfan treated rats). All reactions were normalized for β -Actin mRNA expression. Values with superscripts "*" refers to significant ($P < 0.05$) differences in relative transcript abundance of each group compared to control group

Table 3. Mean \pm SEM of relative abundance of *Bcl2*, *Casp3*, *DAZL* mRNA in treatment groups in comparison with control group

Groups	<i>Bcl2</i>	<i>Casp3</i>	<i>DAZL</i>
DMSO	0.82 \pm 0.13	0.65 \pm 0.24	1.29 \pm 0.11
Vit E	1.67 \pm 0.20	7.68 \pm 0.74*	1.97 \pm 0.08*
AM	0.37 \pm 0.03	0.03 \pm 0.02*	1.77 \pm 0.01
LM	0.2 \pm 0.08*	0.86 \pm 0.17	1.50 \pm 0.10
ALM	0.13 \pm 0.05*	0.49 \pm 0.31*	1.29 \pm 0.01

All reactions were normalized for β -Actin mRNA expression. "*" refers to significant ($P < 0.05$) differences in relative transcript abundance of each group compared to control group

Deleted in azoospermia-like, exhibited high level of expression in Vit E treated rats compared to control counterparts ($P < 0.01$) and no differences were found between each of other treatment groups (DMSO, AM, LM, and ALM) and control group (Fig. 2).

The relative abundance of *Bcl2*, *Casp3*, and *DAZL* transcripts in rat testes tissue derived from treated rats in all treatment groups compared to control group (Busulfan treated rats) are briefly shown in Table 3.

DISCUSSION

One of the most important problems in the life of couples' is infertility and its complications and the causes of some infertilities are related to men. The most common cause of male infertility is inability to produce sufficient numbers of active and healthy sperms [21]. Many factors can affect sperm production and infertility risks. Among these factors are using chemotherapy drugs for cancer, antibiotics, toxic substances, stress, pesticides, radiation, vitamin deficiency, and air pollution. These factors can reduce sperm concentration with generation of free radicals and oxidation of germ cells in the testes [21].

Busulfan, as a chemotherapy drug, is a bifunctional alkylating agent and is used for the treatment of various malignant diseases, such as polycythemia vera and chronic myelogenous leukemia [3]. Busulfan is a highly cytotoxic and genotoxic agent [3] that can induce various adverse effects, both acute and chronic, such as DNA damage and subsequently activates apoptosis or senescence in a cell type-dependent manner probably due to oxidative stress [4] in several biological organs such as hematologic [22], nervous [23] and reproductive organs [24]. It has been confirmed that chemotherapy with busulfan can induce apoptosis in sperm [6] and increase ROS generation and resulting in death of spermatozoa [25]. As well as, occurrence of teratospermia, the presence of spermatozoa with abnormal morphology in semen is the other possible result of busulfan treatment in mice [3]. These studies showed that busulfan is involved in the arrest of spermatogenesis, though some of the changes are reversible and dose dependent. In this study two dose of 25 and 10 mg/kg busulfan with 14 days interval were used which can destruct the spermatogenic process in rat as was shown in other studies [3,26,27].

Antioxidants neutralize free radicals and subsequently the oxidative reactions caused by them. The dietary antioxidants may be beneficial in reducing the lipid peroxidation and DNA damage in sperm during the treatment with busulfan [28]. Different antioxidants have examined in challenging procedures to reduce the adverse effects of ROS on rat spermatozoa [16,29].

Here, we studied the protective effect of vitamin E, melatonin, L-carnitine, and atorvastatin treatment, alone and in different combinations, against busulfan-mediated sperm damage in mice. Our results demonstrated that perform all treatments 48 h after the beginning of chemotherapy with busulfan and daily administration of antioxidants for 8 weeks can slightly reduce busulfan-mediated destructions.

To confirm the role of busulfan in germ cells destruction and apoptosis, respectively, *DAZL* activation and anti-apoptotic molecule *Bcl2* as well as activation of *Casp3* were measured in testes tissue after busulfan treatment.

Consistent with busulfan-induced increased mitochondrial membrane depolarization, significantly increased expression of *Bcl2*, was observed in testes tissue treated with vitamin E compared to other antioxidants-treated rats, suggesting busulfan-induced apoptosis is moderated with Vitamin E administration through regulation of the expression of *Bcl2*, but did not lead to subsequent decrease caspase-3 activation.

Vitamin E (α -tocopherol) is an organic fat soluble compound located generally in cell membranes. This dominant antioxidant extinguishes superoxide anions and free hydroxyl radicals thereby reducing lipid peroxidation initiated by

ROS in plasma membranes^[30] and thus protects the cell membrane from ROS-induced damages. This antioxidant reduces testicular tissue damages caused by cytotoxic agents through increasing the expression of antioxidant related genes. In the male reproductive tract, the antioxidant property of this vitamin in inhibition of destructive effects of free radicals in testes^[31] and sperm^[32] was confirmed. As well as, some studies showed vitamin E is efficient in protecting testes from induced damage by oxidative stress and mitigation of this damage can be achieved by vitamin E treatment^[33].

It was expected high expression of *Bcl2* reduces *Casp3* expression after administration of Vitamin E. But, it should be considered that there are at least two principal pathways for activating *Casp3*. The more ancient, which is induced by diverse intracellular stresses, including cytokine deprivation and genotoxic damage, is regulated by *Bcl2* and its relatives. Progression through the pathway usually leads to the activation of *Casp9* and subsequently *Casp3*. A more-recently evolved pathway is triggered when 'death receptors' on the plasma membrane engaged by cognate ligands of the tumour-necrosis factor (TNF) family, recruit *Casp8* through binding the adaptor protein FAS-associated death domain (FADD) (also called MORT1). This pathway eventually increases *Casp3*^[10]. Therefore, high expression of *Casp3* in vitamin E-treated rats may be due to activation of the second mentioned pathway.

Among the groups, Vitamin E more moderated spermatogenesis resumption as shown in increasing *DAZL* gene expression. This study was the first study in directly determining the effect of vitamin E on *DAZL* expression as a worthy candidate for male infertility that plays an important role in the spermatogenesis.

However, a number of studies suggested that supplementation with oral antioxidants such as vitamin E and carnitine could improve the sperm quality in infertile patients which can certainly be through the stimulation of related gene expression such as *DAZL*.

However, we observed lower *Casp3* gene expression in other treatments, especially in those rats treated with both AM and ALM. When combination groups were compared to each other, it was defined that down-regulation of *Casp3* mRNA in AM and ALM groups might be more due to the presence of melatonin, and on the other hand, high expression of this gene in LM in comparison of AM group and then ALM group was created by L-carnitine. This is consistent with Fan et al.^[34] study that expressed mRNA and protein levels caspase-8 was increased by L-carnitine treatment in hepa1c1c7 mouse cancer cells. As well as, it was noted that despite the inhibition effect of L-carnitine on the activity of recombinant caspases 3 in Jurkat cells (a human T lymphocyte cells line), its long-chain fatty acid derivative palmitoylcarnitine increases the activity of all the caspases. It was suggested that reversed effect

of palmitoylcarnitine on the inhibition of caspase activity by carnitine, may be regulated in part by the balance of palmitoylcarnitine and carnitine under physiological conditions^[35].

Many studies demonstrated that melatonin has an anti-apoptotic effect in somatic and germ cells^[36-38]. Melatonin has been reported to be protective in male reproductive health, which readily crosses the blood-testes barrier and has a very low toxicity^[39]. Studies have investigated the use of melatonin to relieve the side effects of environmental toxins and chemotherapy drugs during spermatogenesis^[36,40]. However, few systematic studies have investigated whether melatonin employs a protective role in the psychological stress-induced impairment of spermatogenesis as well as the mechanisms by which melatonin mitigates the damage in testes.

Many studies have found that L-carnitine and its derivatives can optimize sperm motion parameters^[41,42]. On the other hand, other studies failed to detect significant increases in sperm concentration following L-carnitine treatment^[43-45]. The relatively small doses and short duration of treatment employed may be the main reason why no substantial increases were detected. By the large, it has been found that L-carnitine further improves sperm motility and chromatin quality via antioxidant properties, the enhanced glucose uptake by sperm^[46], and long chain fatty acids transportation across the inner membrane of the mitochondria for use in metabolism^[47]. On the other hand, its effects in reducing some of the side effects of busulfan on the testes is lower than other used compounds^[25]. This may be the reason of low expression of *DAZL* in the L-carnitine-contained groups in comparison to the other groups.

In this study the lowest expression of *Bcl2* was seen in the ALM group. The main reason of this event can be atorvastatin. Atorvastatin improves the lipid profile, lipid oxidation, and oxidative/antioxidative status. These positive effects may be attributed to the antioxidant properties of statins. It has been suggested that statins increase apoptosis and change levels of *Bcl2* family members (e.g., *Bax* increasing and *Bcl2* reduction). Several reports found that statins reduce levels of the anti-apoptotic protein *Bcl2*, and increase apoptosis and cell death^[48-50]. Though, there is evidence that statins increase *Bcl2* abundance which would be desirable and in some instances reduce apoptosis and cell death^[51,52].

Generally at high statin concentrations apoptosis is increased, and *Bcl2* expression levels and cell viability are reduced^[53]. The mechanisms for the statin-induced reduction of *Bcl2* protein levels have not been forthcoming. Statins reduce cholesterol, protein prenylation, and two isoprenoids FPP (farnesyl pyrophosphate) and GGPP (geranylgeranyl pyrophosphate) but how these reductions trigger a weakening of the anti-apoptotic protein *Bcl2* and increase abundance of pro-apoptotic proteins such as *Bax* and *Bim* is not understood. There is evidence that statins has

function outside of the mevalonate pathway. Statins for example bind to a heterodimeric glycoprotein, lymphocyte function-associated antigen-1 (LFA-1) which is a member of the $\beta 2$ integrin family^[54]. Directly related to the subject of statins, these compounds increase *Bcl2* gene expression and protein levels, which do not involve the mevalonate pathway.

It can be concluded that, according to the antiapoptotic effects of these antioxidants, vitamin E retrieved spermatogenesis potential of busulfan-induced infertile male rats better than the other antioxidants used in different combinations. It was approved by increasing *DAZL* gene expression, a worthy candidate for male infertility evaluation.

ACKNOWLEDGMENT

The authors would like to thank Islamic Azad University, Science and Research Branch, Tehran, and Shahrekord University for their supports. We also thank Behnam Bakhtiarimoghdam for keeping rats and injection of drugs in treatment groups.

REFERENCES

- Li B, He X, Zhuang M, Niu B, Wu C, Mu H, Tang F, Cui Y, Liu W, Zhao B, Peng S, Li G, Hua J: Melatonin ameliorates busulfan-induced spermatogonial stem cell oxidative apoptosis in mouse testes. *Antioxid Redox Signal*, 28 (5): 385-400, 2018. DOI: 10.1089/ars.2016.6792
- Dun MD, Aitken RJ, Nixon B: The role of molecular chaperones in spermatogenesis and the post-testicular maturation of mammalian spermatozoa. *Hum Reprod Update*, 18 (4): 420-435, 2012. DOI: 10.1093/humupd/dms009
- Panahi M, Keshavarz S, Rahmanifar F, Tamadon A, Mehrabani D, Karimaghani N, Sepehrimanesh M, Aqababa H: Busulfan induced azoospermia: Stereological evaluation of testes in rat. *Vet Res Forum*, 6 (4): 273-278, 2015.
- Iwamoto T, Hiraku Y, Oikawa S, Mizutani H, Kojima M, Kawanishi S: DNA intrastrand cross-link at the 5'-GA-3' sequence formed by busulfan and its role in the cytotoxic effect. *Cancer Sci*, 95 (5): 454-458, 2004. DOI: 10.1111/j.1349-7006.2004.tb03231.x
- Aitken RJ, Gibb Z, Baker MA, Drevet J, Gharagozloo P: Causes and consequences of oxidative stress in spermatozoa. *Reprod Fertil Dev*, 28 (2): 1-10, 2015. DOI: 10.1071/RD15325
- Choi YJ, Ok DW, Kwon DN, Chung JI, Kim HC, Yeo SM, Kim T, Seo HG, Kim JH: Murine male germ cell apoptosis induced by busulfan treatment correlates with loss of c-kit-expression in a Fas/FasL- and p53-independent manner. *FEBS Lett*, 575 (1-3): 41-51, 2004. DOI: 10.1016/j.febslet.2004.08.034
- Reiter RJ, Tan DX, Terron MP, Flores LJ, Czarnocki Z: Melatonin and its metabolites: New findings regarding their production and their radical scavenging actions. *Acta Biochim Pol*, 54 (1): 1-9, 2007.
- Birben E, Sahiner UM, Sackesen C, Erzurum S, Kalayci O: Oxidative stress and antioxidant defense. *World Allergy Organ J*, 5:270, 2012.
- Reynolds N, Cooke HJ: Role of the DAZ genes in male fertility. *Reprod Biomed Online*, 10 (1): 72-80, 2005. DOI: 10.1016/S1472-6483(10)60806-1
- Cory S, Adams JM: The Bcl2 family: Regulators of the cellular life-or-death switch. *Nat Rev Cancer*, 2 (9): 647-656, 2002. DOI: 10.1038/nrc883
- Cui X, Zhang L, Magli AR, Catera R, Yan XJ, Griffin DO, Rothstein TL, Barrientos J, Koltitz JE, Allen SL, Rai KR, Chiorazzi N, Chu CC: Cytoplasmic myosin-exposed apoptotic cells appear with caspase-3 activation and enhance CLL cell viability. *Leukemia*, 30, 74-85, 2015. DOI: 10.1038/leu.2015.204
- Abd-Elrazek A, Ahmed-Farid OAH: Protective effect of L-carnitine and L-arginine against busulfan-induced oligospermia in adult rat. *Andrologia*, 50 (1): e12806, 2018. DOI: 10.1111/and.12806
- Javaherzadeh M, Shekarchizadeh A, Kafaei M, Mirafshrieh A, Mosaffa N, Sabet B: Effects of intraperitoneal administration of simvastatin in prevention of postoperative intra-abdominal adhesion formation in animal model of rat. *BEAT*, 4 (3): 156, 2016.
- Ferdosi Khosroshahi A, Bakhtiari M, Soleimani Rad J, Koroji M, Roshangar L, Janzadeh A, Kerdari M, Jameie B: Study of the effect of exogenous melatonin on sperm fertility in busulfan induced oligospermic of pinealectomized rat. *Razi J Med Sci*, 20 (110): 77-86, 2013.
- Soyalıç H, Gevrek F, Koç S, Avcu M, Metin M, Aladağ İ: Intraperitoneal curcumin and vitamin E combination for the treatment of cisplatin-induced ototoxicity in rats. *Int J Pediatr Otorhinolaryngol*, 89, 173-178, 2016. DOI: 10.1016/j.ijporl.2016.08.012
- Aksu EH, Akman O, Özkaraç M, Ömür A, Uçar Ö: Effect of *Maclura Pomifera* extract on cisplatin-induced damages in reproductive system of male rats. *Kafkas Univ Vet Fak Derg*, 21, 397-403, 2015. DOI: 10.9775/kvfd.2014.12662
- Gong ZK, Wang SJ, Huang YQ, Zhao RQ, Zhu QF, Lin WZ: Identification and validation of suitable reference genes for RT-qPCR analysis in mouse testis development. *Mol Genet Genomics*, 289 (6): 1157-1169, 2014. DOI: 10.1007/s00438-014-0877-6
- Dorak MT: Real-time PCR. CRC Press, 2007.
- Pfaffl MW: A new mathematical model for relative quantification in real-time RT-PCR. *Nucleic Acids Res*, 29 (9): e45, 2001. DOI: 10.1093/nar/29.9.e45
- Ghasemzadeh-Hasankolaei M, Eslaminejad MB, Batavani R, Ghasemzadeh-Hasankolaei M: Male and female rat bone marrow-derived mesenchymal stem cells are different in terms of the expression of germ cell specific genes. *Anat Sci Int*, 90 (3): 187-196, 2015. DOI: 10.1007/s12565-014-0250-1
- Amin A, Hamza AA: Effects of Roselle and Ginger on cisplatin-induced reproductive toxicity in rats. *Asian J Androl*, 8 (5): 607-612, 2006. DOI: 10.1111/j.1745-7262.2006.00179.x
- Albrecht M, Tackmann W, Pribilla W: Aplastic syndrome in myleran overdose. *Med Klin*, 66 (4): 126-130, 1971.
- Molenaar R, de Rooij DG, Rommerts FFG, Reuvers PJ, van der Molen HJ: Specific destruction of Leydig cells in mature rats after in vivo administration of ethane dimethyl sulfonate. *Biol Reprod*, 33 (5): 1213-1222, 1985. DOI: 10.1095/biolreprod33.5.1213
- Bollag W: The effect of myleran on the germ cells of rats. *Experientia*, 9 (7): 268, 1953.
- Dehghani F, Hassanpour A, Poost-Pasand A, Noorafshan A, Karbalay-Doust S: Protective effects of L-carnitine and homogenized testis tissue on the testis and sperm parameters of busulfan-induced infertile male rats. *Iran J Reprod Med*, 11 (9): 693-704, 2013.
- Mirzapour T, Movahedin M, Tengku Ibrahim TA, Koroji M, Haron AW, Nowroozi MR, Rafieian SH: Effects of basic fibroblast growth factor and leukaemia inhibitory factor on proliferation and short-term culture of human spermatogonial stem cells. *Andrologia*, 44 (Suppl. 1): 41-55, 2012. DOI: 10.1111/j.1439-0272.2010.01135.x
- Cakici C, Buyrukcu B, Duruksu G, Haliloglu AH, Aksoy A, Isik A, Uludag O, Ustun H, Subasi C, Karaoz E: Recovery of fertility in azoospermia rats after injection of adipose-tissue-derived mesenchymal stem cells: the sperm generation. *Biomed Res Int*, 2013: 529589, 2013. DOI: 10.1155/2013/529589
- Khaki A, Fathiazad F, Nouri M, Afshin Khaki A, Ozanci CC, Ghafari-Novin M, Hamadeh M: The effects of Ginger on spermatogenesis and sperm parameters of rat. *Iran J Reprod Med*, 7 (2): 7-12, 2009.
- Yildiz S, Öztürkler Y, Ari UÇ, Lehimcioğlu NC, Atakışi E, Kulaksiz R: The effects of L-ergothioneine, N-acetylcystein and cystein on freezing of ram semen. *Kafkas Univ Vet Fak Derg*, 21 (1): 81-86, 2015. DOI: 10.9775/kvfd.2014.11792
- Lü JM, Lin PH, Yao Q, Chen C: Chemical and molecular mechanisms

of antioxidants: Experimental approaches and model systems. *J Cell Mol Med*, 14 (4): 840-860, 2010. DOI: 10.1111/j.1582-4934.2009.00897.x

31. Mendiola J, Torres-Cantero AM, Vioque J, Moreno-Grau JM, Ten J, Roca M, Moreno-Grau S, Bernabeu R: A low intake of antioxidant nutrients is associated with poor semen quality in patients attending fertility clinics. *Fertil Steril*, 93 (4): 1128-1133, 2010. DOI: 10.1016/j.fertnstert.2008.10.075

32. Lu J, Wang Z, Cao J, Chen Y, Dong Y: A novel and compact review on the role of oxidative stress in female reproduction. *Reprod Biol Endocrinol*, 16: 80, 2018. DOI: 10.1186/s12958-018-0391-5

33. Aydin AF, Coban J, Dogan-Ekici I, Dogru-Abbasoglu S, Uysal M, Kocak-Toker N: Carnosine and vitamin E-a promising pair in the combat against testicular oxidative stress in aged rats. *Andrologia*, 47 (10): 1131-1138, 2015. DOI: 10.1111/and.12392

34. Fan JP, Kim HS, Han GD: Induction of apoptosis by L-carnitine through regulation of two main pathways in Hepa1c1c 7 cells. *Amino Acids*, 36: 365, 2009. DOI: 10.1007/s00726-008-0093-y

35. Mutomba MC, Yuan H, Konyavko M, Adachi S, Yokoyama CB, Esser V, McGarry JD, Babior BM, Gottlieb RA: Regulation of the activity of caspases by L-carnitine and palmitoylcarnitine. *FEBS Lett*, 478 (1-2): 19-25, 2000. DOI: 10.1016/S0014-5793(00)01817-2

36. Ji YL, Wang H, Meng C, Zhao XF, Zhang C, Zhang Y, Zhao M, Chen YH, Meng XH, Xu DX: Melatonin alleviates cadmium-induced cellular stress and germ cell apoptosis in testes. *J Pineal Res*, 52 (1): 71-79, 2012. DOI: 10.1111/j.1600-079X.2011.00921.x

37. Tunon MJ, San Miguel B, Crespo I, Jorquera F, Santamaria E, Alvarez M, Prieto J, Gonzalez-Gallego J: Melatonin attenuates apoptotic liver damage in fulminant hepatic failure induced by the rabbit hemorrhagic disease virus. *J Pineal Res*, 50 (1): 38-45, 2011. DOI: 10.1111/j.1600-079X.2010.00807.x

38. Jang SS, Kim WD, Park WY: Melatonin exerts differential actions on X-ray radiation-induced apoptosis in normal mice splenocytes and Jurkat leukemia cells. *J Pineal Res*, 47 (2): 147-155, 2009. DOI: 10.1111/j.1600-079X.2009.00694.x

39. Rocha CS, Rato L, Martins AD, Alves MG, Oliveira PF: Melatonin and male reproductive health: Relevance of darkness and antioxidant properties. *Curr Mol Med*, 15 (4): 299-311, 2015. DOI: 10.2174/1566524015666150505155530

40. Gobbo MG, Costa CFP, Silva DGH, de Almeida EA, Góes RM: Effect of melatonin intake on oxidative stress biomarkers in male reproductive organs of rats under experimental diabetes. *Oxid Med Cell Longev*, 2015: 614579, 2015. DOI: 10.1155/2015/614579

41. Vicari E, La Vignera S, Calogero AE: Antioxidant treatment with carnitines is effective in infertile patients with prostatovesiculopididymitis and elevated seminal leukocyte concentrations after treatment with nonsteroidal anti-inflammatory compounds. *Fertil Steril*, 78 (6): 1203-1208, 2002. DOI: 10.1016/S0015-0282(02)04350-9

42. Lenzi A, Lombardo F, Sgro P, Salacone P, Caponecchia L, Dondero

F, Gandini L: Use of carnitine therapy in selected cases of male factor infertility: A double-blind crossover trial. *Fertil Steril*, 79 (2): 292-300, 2003. DOI: 10.1016/S0015-0282(02)04679-4

43. Agarwal A, Said TM: Carnitines and male infertility. *Reprod Biomed Online*, 8 (4): 376-384, 2004. DOI: 10.1016/S1472-6483(10)60920-0

44. Moncada ML, Vicari E, Cimino C, Calogero AE, Mongioi A, D'Agata R: Effect of acetylcarnitine treatment in oligoasthenospermic patients. *Acta Eur Fertil*, 23 (5): 221-224, 1992.

45. Mongioi L, Calogero AE, Vicari E, Condorelli RA, Russo GI, Privitera S, Morgia G, La Vignera S: The role of carnitine in male infertility. *Andrology*, 4 (5): 800-807, 2016. DOI: 10.1111/andr.12191

46. Aliabadi E, Soleimani Mehranjani M, Borzoei Z, Talaei-Khozani T, Mirkhani H, Tabesh H: Effects of L-carnitine and L-acetyl-carnitine on testicular sperm motility and chromatin quality. *Iran J Reprod Med*, 10 (2): 77-82, 2012.

47. Matalliotakis I, Koumantaki Y, Evageliou A, Matalliotakis G, Goumenou A, Koumantakis E: L-carnitine levels in the seminal plasma of fertile and infertile men: Correlation with sperm quality. *Int J Fertil Womens Med*, 45 (3): 236-240, 2000.

48. Igarashi M, Yamaguchi H, Hirata A, Tsuchiya H, Ohnuma H, Tominaga M, Daimon M, Kato T: Mechanisms of inhibitory effects of cerivastatin on rat vascular smooth muscle cell growth. *J Cardiovasc Pharmacol*, 40 (2): 277-287, 2002. DOI: 10.1097/00005344-200208000-00013

49. Muck AO, Seeger H, Wallwiener D: Class-specific pro-apoptotic effect of statins on human vascular endothelial cells. *Z Kardiol*, 93 (5): 398-402, 2004. DOI: 10.1007/s00392-004-0081-5

50. Spanpanato C, De Maria S, Sarnataro M, Giordano E, Zanfardino M, Baiano S, Carteni M, Morelli F: Simvastatin inhibits cancer cell growth by inducing apoptosis correlated to activation of Bax and down-regulation of BCL-2 gene expression. *Int J Oncol*, 40 (4): 935-941, 2012. DOI: 10.3892/ijo.2011.1273

51. Kwak HB, Thalacker-Mercer A, Anderson EJ, Lin CT, Kane DA, Lee NS, Cortright RN, Bamman MM, Neuffer PD: Simvastatin impairs ADP-stimulated respiration and increases mitochondrial oxidative stress in primary human skeletal myotubes. *Free Radic Biol Med*, 52 (1): 198-207, 2012. DOI: 10.1016/j.freeradbiomed.2011.10.449

52. Zhao XH, Xu ZR, Zhang Q, Yang YM: Simvastatin protects human osteosarcoma cells from oxidative stress-induced apoptosis through mitochondrial-mediated signaling. *Mol Med Rep*, 5 (2): 483-488, 2012. DOI: 10.3892/mmr.2011.641

53. Wood WG, Igbavboa U, Muller WE, Eckert GP: Statins, Bcl-2, and apoptosis: Cell death or cell protection? *Mol Neurobiol*, 48 (2): 308-314, 2013. DOI: 10.1007/s12035-013-8496-5

54. Weitz-Schmidt G, Welzenbach K, Brinkmann V, Kamata T, Kallen J, Bruns C, Cottens S, Takada Y, Hommel U: Statins selectively inhibit leukocyte function antigen-1 by binding to a novel regulatory integrin site. *Nat Med*, 7 (6): 687-692, 2001. DOI: 10.1038/89058