

# First Molecular Evidence for *Mycoplasma haemocanis* and *Candidatus Mycoplasma haematoparvum* in Asymptomatic Shelter Dogs in Kyrgyzstan

Kursat ALTAY <sup>1,a</sup> Mehmet Fatih AYDIN <sup>2,b</sup> Ayperi AYTMIRZAKIZI <sup>3,c</sup>  
Zarima JUMAKANOVA <sup>3,d</sup> Ayday CUNUSOVA <sup>3,e</sup> Nazir DUMANLI <sup>3,4,f</sup>

<sup>1</sup> Department of Parasitology, Faculty of Veterinary Medicine, University of Sivas Cumhuriyet, TR-58140 Sivas - TURKEY

<sup>2</sup> Department of Public Health, Faculty of Health Sciences, University of Karamanoglu Mehmetbey, TR-70100 Karaman - TURKEY

<sup>3</sup> Faculty of Veterinary Medicine, Kyrgyz-Turkish Manas University, 720044, Bishkek - KYRGYZSTAN

<sup>4</sup> Department of Parasitology, Faculty of Veterinary Medicine, University of Firat, TR-23119 Elazig - TURKEY

<sup>a</sup> ORCID: 0000-0002-5288-1239; <sup>b</sup> ORCID: 0000-0002-8325-4887; <sup>c</sup> ORCID: 0000-0003-3363-0941; <sup>d</sup> ORCID: 0000-0001-5781-8004

<sup>e</sup> ORCID: 0000-0002-5331-7648; <sup>f</sup> ORCID: 0000-0002-6916-8210

Article ID: KVFD-2019-22196 Received: 07.03.2019 Accepted: 03.07.2019 Published Online: 03.07.2019

## How to Cite This Article

**Altay K, Aydin MF, Aytmirzakizi A, Jumakanova Z, Cunusova A, Dumanli N:** First molecular evidence for *Mycoplasma haemocanis* and *Candidatus Mycoplasma haematoparvum* in Asymptomatic shelter dogs in Kyrgyzstan. *Kafkas Univ Vet Fak Derg*, 2019 (Article in Press). DOI: 10.9775/kvfd.2019.22196

## Abstract

*Mycoplasma haemocanis* (Mhc) and *Candidatus Mycoplasma haematoparvum* (CMhp) have been investigated using species specific PCR and sequencing in 170 dogs from Kyrgyzstan. Maximum likelihood estimation (MLE) of the infection rates with 95% confidence intervals (CI) was calculated. The molecular prevalence of hemoplasma infection was 5.29% (CI 2.57-9.34). It was found that, five (2.94%, CI 1.06-6.22) samples were found to be infected with Mhc, one (0.59%, CI 0.03-2.57) sample with CMhp and three (1.76%, CI 0.44-4.52) samples with both species. These results demonstrate that dogs can be exposed to each haemoplasma species and provide first molecular evidence for these species in Kyrgyzstan.

**Keywords:** Canine, Haemoplasmas, PCR, Kyrgyzstan

## Kırgızistan'da Aseptomatik Barınak Köpeklerinde *Mycoplasma haemocanis* ve *Candidatus Mycoplasma haematoparvum* için İlk Moleküler Kanıt

## Öz

Kırgızistan'dan 170 köpekte tür spesifik PZR ve sekanslama ile *Mycoplasma haemocanis* (Mhc) ve *Candidatus Mycoplasma haematoparvum* (CMhp) araştırılmıştır. Enfeksiyon oranlarının en büyük olasılık tahmini (MLE) %95 güven aralığında (CI) hesaplandı. Hemoplasma enfeksiyonunun moleküler prevalansı %5.29 idi (CI 2.57-9.34). Beş (%2.94, CI 1.06-6.22) numunenin Mhc ile, bir (%0.59, CI 0.03-2.57) numunenin CMhp ile ve üç (%1.76, CI 0.44-4.52) numunenin de her iki tür ile enfekte olduğu bulundu. Bu sonuçlar, Kırgızistan'daki köpeklerin hemoplasma türlerinin her biri ile maruz kalabileceğini göstermekte ve bu türler için Kırgızistan'daki ilk moleküler kanıtı sağlamaktadır.

**Anahtar sözcükler:** Köpek, Hemoplasma, PZR, Kırgızistan

## INTRODUCTION

Vectors and vector-borne diseases have a considerable impact for domestic and wild animals in tropical and subtropical climatic regions worldwide. Haemotropic mycoplasmas or hemoplasmas are bacteria infect in a wide range of vertebrate

erythrocytes and recently renamed from *Haemobartonella* and *Eperythrozoon* <sup>[1]</sup>. Well known canine haemoplasma species, *Mycoplasma haemocanis* (Mhc) and *Candidatus Mycoplasma haematoparvum* (CMhp), cause subclinical or chronic disease in immunocompetent dogs and acute disease with hemolytic anemia in susceptible animals related



### İletişim (Correspondence)



+90 338 2262000/4310 Fax: +90 338 2262134



veterinermfa@gmail.com

to splenectomy, immunosuppression and concurrent infections. Lethargy, weight loss, fever and anorexia are the other symptoms for acute disease [2,3].

In several studies worldwide, *Mycoplasma* infections have been found in stray, wild and pet dogs in Turkey [4], Nigeria [5], United States [3], Brazil [6], Thailand [7], Iran [8,9], Italy [10], Spain, Portugal, Switzerland and France [2].

Recently, *Hepatozoon canis* infections with high prevalence in dogs from Kyrgyzstan were disclosed [11]. We aimed to investigate frequency of infection with Mhc and CMhp in dogs from Kyrgyzstan using polymerase chain reaction (PCR) and sequence analysis.

## MATERIAL and METHODS

### Ethic Statement

The Ethic statement was obtained from the Animal Experimentation Ethics Committee of Kyrgyz-Turkish Manas University (Document No: 29.06.2017/2017-06/01).

### Study Area and Samples

Bishkek, largest city and capital of Kyrgyzstan, is located at 42.87 latitude and 74.59 longitude, 800 meters above sea level and has a surface area of 169.9 km<sup>2</sup> for city center. Bishkek can show both temperate and continental climate

characteristics. Province has an average annual rainfall of 427 mm (Fig. 1). The study was conducted on 170 apparently asymptomatic dogs from May 2016-October 2017. Five mL of blood sample were taken from the *vena cephalica antebrachii* into tubes containing K3EDTA-anti-coagulant from shelter dogs with cooperation Kyrgyz-Turkish Manas University Veterinary Teaching Hospital.

### Nucleic Acid Extractions and PCR Assay


For genomic DNA isolation, 200 µL blood was processed with a commercial kit [PureLink Genomic DNA mini kit (Invitrogen, Carlsbad, USA)]. Target DNA's were kept at -20°C until analysis. To determine each species, a single PCR analysis were made in a final reaction volume of 25 µL containing PCR buffer [750 mM Tris-HCl (pH 8.8), 200 mM (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub>, 0.1% Tween 20], 5 mM MgCl<sub>2</sub>, 125 µM deoxy-nucleotide triphosphates, 1.25 U Taq DNA polymerase (Promega, Madison, WI, USA), primers (20 pmol/µL) and template DNA. Sequence, specificity, target gene and product sizes for primers were demonstrated in Table 1. PCR was performed with an initial denaturation step of 94°C for 5 min was followed by 32 cycles of 94°C for 1 min, 60°C for 1 min, and 72°C for 1 min. A final extension step at 72°C for 5 min was also applied [9]. Positive control DNA for Mhc (GenBank accession no: MG594502) and CMhp (GenBank accession no: MG594500) species and negative controls (nuclease-free water) were also used in the PCR



**Fig 1.** Map of Kyrgyzstan and location of Bishkek

**Table 1.** Primers used in this study

Primer	Sequence (5'-3')	Specificity	Target Gene	Product Length (bp)	Reference
Forward	GAAACTAAGGCCATAAATGACGC	Mhc	16S rRNA	309	[9]
Reverse	ACCTGTACCTCGATAACCTCTAC				
Forward	ACGAAAGTCTGATGGAGCAATAC	CMhp	16S rRNA	328	[9]
Reverse	TATCTACGCATTCCACCGCTAC				



**Fig 2.** Agarose-gel electrophoresis of *Mycoplasma haemocanis* (a) and *Candidatus Mycoplasma haematoparvum* (b) specific polymerase chain reaction. M: 1 kb DNA ladder-marker, lane a1-a8: *Mycoplasma haemocanis* positive dog blood samples; lane a9, a10: negative dog blood samples, lane a11: negative control distilled water; lane a12: positive control DNA from dog, lane b1-b4: *Candidatus Mycoplasma haematoparvum* positive dog blood samples; lane b5, b6: negative dog blood samples, lane b7: negative control distilled water; lane b8: positive control DNA from dog

**Table 2.** Molecular prevalence of canine hemoplasma species detected by PCR in Kyrgyzstan (n=170)

Overall Prevalence (n = 9)	<i>Mycoplasma haemocanis</i>	<i>Candidatus Mycoplasma haematoparvum</i>	Mhc+CMhp
5	5	-	-
3	-	-	3
1	-	1	-
Total	9 (5.29%) (CI; 2.57-9.34)	5 (2.94%) (CI; 1.06-6.22)	1 (0.59%) (CI; 0.03-2.57)

reaction. Five microliters of PCR product was separated using electrophoresis (100 V, 60 min) in a 1.5 % agarose gel stained with ethidium bromide and visualized using Gel Doc (Bio-Rad, Hercules, CA, USA) (Fig. 2).

### Sequencing and Molecular Classification

One positive sample for each species were selected to validate PCR results. After purification of PCR products by QIAquick PCR purification kit (Qiagen, Hilden, Germany) sequencing were performed by a commercial company (Macrogen, South Korea). Sequences were edited by

Chromas versiyon 2.6.5 (<http://technelysium.com.au/wp/>) and compared with the other sequences available in Genbank (<https://blast.ncbi.nlm.nih.gov/Blast.cgi>). The sequences of the partial 16S ribosomal RNA gene of Mhc and CMhp have been deposited in GenBank databases under accession no: MK015018 and MK026012 respectively.

## RESULTS

Distribution and frequency of hemoplasma species in 170 dogs were determined. The overall prevalence for hemoplasma infection was 5.29% (CI 2.57-9.34). Nine animals were found to be infected by one or more species (Table 2). Mixed infections were determined in 3 of 170 samples with a rate of 1.76% (CI 0.44-4.52) and single infections were in 6 with a rate of 3.53% (CI 1.41-7.03). While five (2.94%, CI 1.06-6.22) samples were found to be infected with Mhc, CMhp was detected in one (0.59%; CI 0.03-2.57) sample.

Obtained sequences comparisons exhibited that while Mhc sequence identified in this study (MK015018) showed 99-100% similarity with the previously reported sequences for the 16S ribosomal RNA gene of Mhc (KY117656, KP715858, EF416567, AB848714) and *Mycoplasma haemofelis* (KM275238, KR905462), CMhp sequence (MK026012) shared 99-100% identity with sequences for the 16S ribosomal RNA gene of CMhp (MG594500, KF366443, HQ918288) and 97-98% identity with *Candidatus Mycoplasma haemominutum* (JQ689947, AY150974, JQ044683).

## DISCUSSION

This study exhibits, for the first time, molecular evidence and prevalence of hemoplasma in dogs in Kyrgyzstan. The overall molecular prevalence of canine hemoplasma species was 5.29% (CI 2.57-9.34). Mhc, CMhp and Mhc + CMhp prevalences were 2.94% (CI 1.06-6.22), 0.59% (CI 0.03-2.57) and 1.76% (CI 0.44-4.52) respectively with this study. Low prevalence of hemoplasmas determined with this study was similar to data from Italy 4.5%<sup>[10]</sup> and Nigeria with 7.7%<sup>[5]</sup>. While higher prevalences were reported in Portugal 40%<sup>[2]</sup>, Iran 23%<sup>[9]</sup> and Turkey 15.3%<sup>[4]</sup>; low prevalence was from USA 1.3%<sup>[3]</sup>. Low prevalence in the studied area may be correlated to climate conditions and/or lack of vector diversity.

In the present study Mhc has higher molecular prevalence than CMhp and this result is in accordance with previous publications<sup>[5,9]</sup>. Three dogs (1.76%) were found to be infected for both Mhc and CMhp. Similar to our results, mix infections were determined in several studies<sup>[2,4,9]</sup>.

Diagnosis of hemotropic *Mycoplasma* species is based on microscopic examination of thin blood smears<sup>[8]</sup> or PCR analysis targeting 16S rRNA gene fragment<sup>[1,12]</sup>. Microscopic examination of smears may be useful and cheap in acute cases but it is not possible to discriminate species

and also determine carrier animals with this method. Molecular methods have always been found superior to microscopic examination for detection and differentiation of hemotropic *Mycoplasma* spp. and other tick-borne agents<sup>[13,14]</sup>. In this study a species-specific PCR assay with high sensitivity and specificity were applied to determine carrier animals for hemotropic *Mycoplasma* spp. in dogs. It was determined that overall prevalence of canine hemoplasma species was 5.29% in Kyrgyzstan and Mhc and CMhp circulate there. We recommend species-specific PCR for Mhc and CMhp in routine screening of blood donors.

Transmission of canine hemoplasma species is associated with haematophagous arthropods like fleas and ticks<sup>[2]</sup>. Also, blood transfusion from carrier dogs to splenectomized dogs induces transmission the organism<sup>[12]</sup>. Furthermore, CMhp is accepted as a zoonotic microorganism<sup>[15]</sup>. Since it was determined with this study dog population in Kyrgyzstan is carrier for both Mhc and CMhp, veterinarians and medical doctors should take these species into consideration in suspected cases.

In conclusion canine hemoplasma species were determined in Kyrgyzstan dog population for the first time and the molecular prevalence for hemotropic *Mycoplasma* spp. in Kyrgyzstan dog population is 5.29%. It was also determined that dog population in Kyrgyzstan exposure to either or both hemoplasma species of Mhc and CMhp and Mhc has higher molecular prevalence than CMhp. It is suggested that further studies aimed to determine molecular prevalence of canine hemoplasmas and potential arthropod vectors should be conducted in other provinces of Kyrgyzstan. We also suggest to veterinarians to be conscious for canine hemoplasma among anemic dogs and routine screening of blood donors may be useful to prevent spread of disease.

## ACKNOWLEDGMENTS

The authors thank Dr. Munir Aktas and Dr. Sezayi Ozubek (Department of Parasitology, College of Veterinary Medicine, Firat University, Turkey) for providing positive control DNA's of Mhc and CMhp and also, we are grateful to animal shelter staffs and Kyrgyz-Turkish Manas University Veterinary Teaching Hospital staffs for their kind help during sample collection. The authors declare that they have no conflicts of interest.

## REFERENCES

1. **Messick JB:** New perspectives about *Hemotropic mycoplasma* (formerly, *Haemobartonella* and *Eperythrozoon* species) infections in dogs and cats. *Vet Clin North Am Small Anim Pract*, 33 (6): 1453-1465, 2003. DOI:

10.1016/j.cvsm.2003.08.002

2. **Novacco M, Meli ML, Gentilini F, Marsilio F, Ceci C, Pennisi MG, Lombardo G, Lloret A, Santos L, Carrapic OT, Willi B, Wolf G, Lutz H, Hofmann-Lehmann R:** Prevalence and geographical distribution of canine hemotropic mycoplasma infections in Mediterranean countries and analysis of risk factors for infection. *Vet Microbiol*, 142 (3-4): 276-284, 2010. DOI: 10.1016/j.vetmic.2009.09.069

3. **Compton SM, Maggi RG, Breitschwerdt EB:** *Candidatus Mycoplasma haematoparvum* and *Mycoplasma haemocanis* infections in dogs from the United States. *Comp Immunol Microbiol Infect Dis*, 35 (6): 557-562, 2012. DOI: 10.1016/j.cimid.2012.06.004

4. **Aktas M, Ozubek S:** A molecular survey of hemoplasmas in domestic dogs from Turkey. *Vet Microbiol*, 221, 94-97, 2018. DOI: 10.1016/j.vetmic.2018.06.004

5. **Aquino LC, Kamani J, Haruna AM, Paludo GR, Hicks CA, Helps CR, Tasker S:** Analysis of risk factors and prevalence of haemoplasma infection in dogs. *Vet Parasitol*, 221, 111-117, 2016. DOI: 10.1016/j.vetpar.2016.03.014

6. **de Sousa KCM, Herrera HM, Secato CT, Oliveira ADV, Santos FM, Rocha FL, Barreto WTG, Macedo GC, de Andrade Pinto PCE, Machado RZ, Costa MT, André MR:** Occurrence and molecular characterization of hemoplasmas in domestic dogs and wild mammals in a Brazilian wetland. *Acta Trop*, 171, 172-181, 2017. DOI: 10.1016/j.actatropica.2017.03.030

7. **Kaewmongkol G, Lukkana N, Yangtara S, Kaewmongkol S, Thengchaisri N, Sirinarumitr T, Jittapalpong S, Fenwick SG:** Association of *Ehrlichia canis*, Hemotropic *Mycoplasma* spp. and *Anaplasma platys* and severe anemia in dogs in Thailand. *Vet Microbiol*, 201, 195-200, 2017. DOI: 10.1016/j.vetmic.2017.01.022

8. **Sharifiyazdi H, Abbaszadeh Hasiri M, Amini AH:** Intravascular hemolysis associated with *Candidatus Mycoplasma haematoparvum* in a non-splenectomized dog in the south region of Iran. *Vet Res Forum*, 5 (3): 243-246, 2014.

9. **Torkan S, Aldavood SJ, Sekhavatmandi A, Moshkelani S:** Detection of hemotropic *Mycoplasma* (*Haemobartonella*) using multiplex PCR and its relationship with epidemiological factors in dogs. *Comp Clin Pathol*, 23 (3): 669-672, 2014. DOI: 10.1007/s00580-012-1668-2

10. **Ravagnan S, Carli E, Piseddu E, Da Rold G, Porcellato E, Zanardello C, Carminato A, Vascellari M, Capelli G:** Prevalence and molecular characterization of canine and feline hemotropic mycoplasmas (hemoplasmas) in northern Italy. *Parasit Vectors*, 10 (1): 132, 2017. DOI: 10.1186/s13071-017-2069-9

11. **Altay K, Aydin MF, Aytmirzakizi A, Jumakanova Z, Cunusova A, Dumanli N:** Molecular survey of hepatozoonosis in natural infected dogs: First detection and molecular characterisation of *Hepatozoon canis* in Kyrgyzstan. *Kafkas Univ Vet Fak Derg*, 25 (1): 77-81, 2019. DOI: 10.9775/kvfd.2018.20352

12. **Messick JB:** Hemotropic mycoplasmas (hemoplasmas): A review and new insights into pathogenic potential. *Vet Clin Pathol*, 33 (1): 2-13, 2004. DOI: 10.1111/j.1939-165X.2004.tb00342.x

13. **Tasker S, Binns SH, Day MJ, Gruffydd-Jones TJ, Harbour DA, Helps CR, Jensen WA, Olver CS, Lappin MR:** Use of a PCR assay to assess the prevalence and risk factors for *Mycoplasma haemofelis* and '*Candidatus Mycoplasma haemominutum*' in cats in the United Kingdom. *Vet Rec*, 152, 193-198, 2003. DOI: 10.1136/vr.152.7.193

14. **Aydin MF, Aktas M, Dumanli N:** Molecular identification of *Theileria* and *Babesia* in sheep and goats in the Black Sea Region in Turkey. *Parasitol Res*, 112 (8): 2817-2824, 2013. DOI: 10.1007/s00436-013-3452-x

15. **Maggi RG, Mascarelli PE, Havenga LN, Naidoo V, Breitschwerdt EB:** Co infection with *Anaplasma platys*, *Bartonella henselae* and *Candidatus Mycoplasma haematoparvum* in a veterinarian. *Parasit Vectors*, 6:103, 2013. DOI: 10.1186/1756-3305-6-103