

# The Effect of Oregano Oil (*Origanum vulgare*) on the Fattening Performance and Blood Oxidant-Antioxidant Balance in Post-weaned Tuj Lambs

Özlem DURNA AYDIN<sup>1,a</sup> Oğuz MERHAN<sup>2,b</sup> Gültekin YILDIZ<sup>3,c</sup> Songül ULUFER<sup>4,d</sup>

<sup>1</sup> Kafkas University, Veterinary Faculty, Department of Animal Nutrition, and Nutritional Diseases, TR-36040 Kars - TURKEY

<sup>2</sup> Kafkas University, Veterinary Faculty, Department of Biochemistry, TR-36040 Kars - TURKEY

<sup>3</sup> Ankara University, Veterinary Faculty, Department of Animal Nutrition, and Nutritional Diseases, TR-06830 Ankara - TURKEY

<sup>4</sup> Kafkas University, Vocational School of Kağızman, Pharmacy Services, TR-36700 Kars - TURKEY

<sup>a</sup> ORCID: 0000-0003-4532-6795; <sup>b</sup> ORCID: 0000-0002-3399-0667; <sup>c</sup> ORCID: 0000-0002-1003-9254; <sup>d</sup> ORCID: 0000-0001-9533-0165

Article ID: KVFD-2019-22272 Received: 07.03.2019 Accepted: 16.07.2019 Published Online: 21.07.2019

## How to Cite This Article

Durna Aydın Ö, Merhan O, Yıldız G, Ulufer S: The effect of oregano oil (*Origanum vulgare*) on the fattening performance and blood oxidant-antioxidant balance in post-weaned Tuj lambs. *Kafkas Univ Vet Fak Derg*, 2019 (Article in Press). DOI: 10.9775/kvfd.2019.22272

## Abstract

The purpose of this study is to investigate the effect of oregano oil (*Origanum vulgare*) at on fattening performance and blood oxidant-antioxidant balance such as glutathione (GSH), superoxide dismutase (SOD) and glutathione peroxidase (GPx) catalase (CAT) enzyme activities, malondialdehyde, ceruloplasmin, nitric oxide, albumin, total protein and globulin in post-weaned Tuj breed lambs. The lambs were average 155 days old, and were divided into 3 groups regardless of gender, and in total 18 lambs were used. In the study the control group (C) was fed with basal ration; while the experimental groups were orally fed with oregano oil supplementation as a commercial product in daily doses of 1 mL/lamb/day (T1) and 2 mL/lamb/day (T2), respectively, in addition to the basal ration, via sterile injections. The experiment was maintained for a total of 52 days of which 10 days were adaptation period. In the study, it has been seen in post-weaned Tuj breed lambs that using different doses of oregano oil had no effect on live weight gain, daily live weight gain, daily feed consumption, and feed conversion ratio ( $P>0.05$ ). In the study, it was determined that the addition of oregano oil to the ration made no significant difference on fattening performance and MDA, GSH, CAT, ceruloplasmin, nitric oxide, albumin, total protein and globulin values of blood oxidant-antioxidant parameters. SOD and GPx values were significantly affected by the addition of oregano oil to the lamb rations at the point of examining the effect on blood oxidant-antioxidant balance ( $P<0.05$ ). As a result, it has been determined that oregano oil did not affect the performance parameters, it significantly affected the oxidant-antioxidant balance in lambs.

**Keywords:** Oxidant-antioxidant balance, Tuj, Lamb, Performance, Oregano oil

## Sütten Kesilmiş Tuj Irkı Kuzularda Kekik Yağı (*Origanum vulgare*) Kullanımının Besi Performansı ve Kan Oksidan-Antioksidan Denge Üzerine Etkisi

### Öz

Bu çalışmanın amacı sütten kesilmiş Tuj ırkı kuzularda kekik yağının performans ve glutatyon (GSH), süperoksit dismutaz (SOD) oranları, glutatyon peroksidaz (GPx) katalaz (CAT) enzim aktiviteleri, seruloplazmin, nitrik oksit, albumin, total protein ve globulin kan oksidan-antioksidan denge üzerine etkisinin araştırılmasıdır. Kuzular ortalama 155 günlük yaşta olup cinsiyet gözetilmeksizin 3 gruba ayrılmış ve toplamda 18 kuzu kullanılmıştır. Çalışmada kontrol grubu (K) bazal rasyonla beslenirken deneme gruplarına bazal rasyona ek olarak sırasıyla 1 mL/kuzu/gün (T1) ve 2 mL/kuzu/gün (T2) dozlarında ticari ürün olan kekik yağı ilavesi steril şırıngalar ile hayvanlara günlük olarak oral şekilde verilmiştir. Deneme, 10 gün adaptasyon dönemi olmak üzere toplam 52 gün sürdürülmüştür. Çalışmada kesilmiş Tuj ırkı kuzularda kekik yağının farklı dozlarda kullanımının canlı ağırlık artışı, günlük canlı ağırlık artışı, günlük yem tüketimi ve yemden yararlanma oranını etkilemediği görülmüştür ( $P>0.05$ ). Çalışmada rasyona kekik yağı katılmasının, besi performansı ve kan oksidan antioksidan parametrelerinden MDA, NO, GSH, CAT, seruloplazmin, albümin, total protein ve globulin değerlerinin üzerinde önemli bir farklılık oluşturmadığı saptanmıştır. Kan oksidan antioksidan denge üzerine etkinin irdelenmesi noktasında SOD ve GPx değerleri kuzu rasyonlarında kekik yağı ilavesinden anlamlı düzeyde etkilenmiştir ( $P<0.05$ ). Sonuç olarak kekik yağının kuzularda performans parametrelerini etkilemediği, kan oksidan-antioksidan denge üzerine olumlu etkilerinin olduğu tespit edilmiştir.

**Anahtar sözcükler:** Oksidan-antioksidan denge, Tuj, Kuzu, Performans, Kekik yağı



İletişim (Correspondence)



+90 474 242 68 36



odurna36@gmail.com.tr

## INTRODUCTION

Because the antibiotics and the antimicrobial agents, used in animal feeding for many years as growth stimulants, have given rise to the development of resistance in pathogenic bacteria in humans, and due to the residues they can generate in animal products, the demands for such products is decreasing day by day <sup>[1,2]</sup>. In recent years, with the prominence of consumer demand in terms of product characteristics, the desire to obtain more natural and more suitable products for human health has made the field of animal feeding with new searches a current issue, and many of the researchers have focused their attention on natural and aromatic plants, and essential oils obtained from them <sup>[3,4]</sup>. Oregano is a common term for the plant family *Lamiaceae*, which has more than 60 species, and known by their general aroma and taste <sup>[5]</sup>. Among these species *Thymus*, *Origanum*, *Satureja*, *Thymbra* and *Coridothymus* are of great importance both prevalently and economically <sup>[6]</sup>. Turkey is the country with the world's most important exporter thyme. A large part of the thyme exported from Turkey is *Origanum* species <sup>[7]</sup>. Both oregano itself and its extracts (oregano essential oil) contain substances that induce such as antiseptic, antioxidant, antimicrobial and aroma-regulating effects. It has been reported that oregano contains *phenols*, *thymo*, *carvacrol*, *monoterpene hydrocarbons*, *p-cymene* and *γ-terpinene* <sup>[8]</sup>. Oregano is rich in *carvacrol*, and at a lesser extent in *phenolic monoterpenoids* (particularly thymol) <sup>[9]</sup>.

The short-lived reactive atoms and molecules with un-conjugated electrons in their outer orbitals are known as free radicals and the enzymatic and non-enzymatic structures trying to prevent radicals and their reactions are defined as antioxidants <sup>[10]</sup>. In normal functioning cells, there is a balance between the free radicals produced as a by-product of cell function and the antioxidants which inactivate them, and is described as an oxidant-antioxidant balance. Oxidative stress occurs when there is a free radical production in the cells that overcomes the antioxidant defense system; leads to protein and DNA oxidation, and lipid peroxidation in cells; and consequently causes cell destruction and tissue damage <sup>[10]</sup>. Therefore, studies on the potential antioxidant effects of plant extracts have accelerated especially in recent years. Farag et al. <sup>[11]</sup> examined the relationship between the chemical composition and antioxidant properties of essential oils, and stated that due to the presence of phenolic OH groups acting as a hydrogen donor to the peroxide radicals released during the first step of the lipid oxidation, thymol owns high antioxidant activity by reducing the formation of hydroxy peroxide.

In recent years, several studies focused on the aromatic plants and their extracts. Oregano oil has antioxidant properties. Therefore, oregano oil has been used in many studies because of its antioxidant properties. Turkey is very important in the world in the export of oregano. Oregano

oil used in our study were obtained from the Mediterranean region of Turkey. There are many studies in the literature on the use of oregano in lambs. However, no study was seen in the literature about the use of oregano oil in Tuj breed lambs. Therefore, the aim of this study is to investigate the effect of oregano oil on fattening performance and blood oxidant-antioxidant balance such as glutathione (GSH), superoxide dismutase (SOD) and glutathione peroxidase (GPx) catalase (CAT) enzyme activities, malondialdehyde (MDA), ceruloplasmin, nitric oxide (NO), albumin, total protein and globulin in post-weaned Tuj breed lambs.

## MATERIAL and METHODS

### *Animals, Experimental Design and Feed*

This study was carried out with the permission of the Kafkas University Animal Experiments Local Ethics Committee (Decision No: KAU-HAYDEK/2018-086) report. In the study, Tuj breed lambs, being raised in Kars, Ardahan and Igdir provinces, which is a local sheep breed in Turkey were used as an animal material. The average age of the lambs was 155 days. Lambs were divided into 3 groups, regardless of gender, with a mean live weight of 34.75±0.22 kg, and a total of 18 lambs were used. The Experiment was conducted in Kafkas University Faculty of Veterinary Medicine, Prof. Dr. Ali Riza AKSOY Training, Research and Implementation Farm. The experiment was maintained for a total of 52 days of which 10 days were adaptation period. During the study, the animals were housed in individual boxes (180 cm × 150 cm × 120 cm; height, length, width, respectively) equipped with feeders dispensing pasture grass and concentrate feed separately. During the study, the animals were fed with pasture grass and lamb grower feed.

In the study; dry matter, energy and other nutrient requirements of animals were calculated according to NRC <sup>[12]</sup> standards. The daily amount of concentrate feed provided to the lambs was 600 g per animal. Pasture grass and water was supplied *ad libitum* during the trial. Animals were fed mainly with roughage. The content of the concentrated feed is given in *Table 1*. In adaptation period that lasted 10 days lambs got accustomed to the experimental diet. All lambs were treated against internal and external parasites. The study was conducted in Kars. Kars is a place where important pastures are at for Turkey. Tuj lambs used in the study are a region-specific race. Tuj lambs are fed with pasture-based feed in the region. For this reason, in our study, while feeding tuj lambs with the pasture *ad libitum*, 600 g of concentrated feed were given daily. In this wise, the effect of using oregano oil based on pasture feeding method on performance and blood oxidant and antioxidant balance was investigated.

In the study, control group (C) was fed with basal ration. In addition to basal ration, the experimental groups were fed with a supplementation of oregano oil (*Origanum vulgare*, BOTALIFE™), a commercial product at doses of 1 mL/lamb/

Ingredients	Amount, %
Wheat	3.75
Barley	17.5
Corn	12.5
Soya bean meal	2.5
DDGS	2.5
Sunflower seed meal (28-30 CP)	2.5
Sunflower seed meal (36 CP)	2.5
Safflower meal	6.25
Cotton seed meal (28-30 CP)	4.6
Wheat bran	20
Corn bran	6.3
Leaf cotton (25 CP)	6.25
Molasses	9
Marble powder	3.25
Salt	0.5
Vitamin mineral premix	0.1

<sup>1</sup> CP: Crude Protei  
<sup>2</sup> DDGS: Dried Distillers Grains with Solubles  
<sup>3</sup> The vitamin & mineral premix provided the following (per kg): 4,000,000 IU Vit. A, 800,000 IU Vit. D<sub>3</sub>, 5,000 IU Vit. E, 400 mg Vit. B<sub>2</sub>, 2 mg Vit. B<sub>12</sub>, 5,000 mg Vit. PP, 1,000 mg D-pantothenic acid, 20,000 mg choline, 50 mg Co, 5,400 mg Fe, 185 mg I, 6,900 mg Mn, 800 mg Cu, 6,400 mg Zn, 14 mg Se

day (T1) and 2 mL/lamb/day (T2), respectively. Oregano oil was given orally by sterile syringes to the experimental groups daily. Oregano oil used in the study, *Origanum vulgare* belong to the species of plants are grown in Turkey. Oregano oil used in the study contains 4.15% thymol and 75.3% carvacrol. The essential oil acid profile of the commercial oregano oil is presented in *Table 2*.

### Feed Analysis

Nutrient analysis of the feed was determined according to the method reported in AOAC [13]; while NDF (Neutral Detergent Fiber) and ADF (Acid Detergent Fiber) analyses for pasture grass were determined according to the method reported by Goering and Van Soest [14].

### Determination of Performance Parameters

Animals were weighed before morning feeding in the beginning and on the 0<sup>th</sup>, 10<sup>th</sup>, 31<sup>st</sup> and 52<sup>nd</sup> days of the trial. At the end of the trial, daily feed consumption and feed conversion ratios of each group were calculated. Feed conversion ratio was calculated as the proportion of daily feed consumption to daily weight gain (kg/kg).

### Biochemical Analyses

On the 10<sup>th</sup> and 52<sup>nd</sup> days during the experimental period, after separating a fair amount of the blood samples, taken from the *V. jugularis* of the animals with anticoagulant (EDTA) tubes, as whole blood, plasma of the remaining

Ingredient	%	Ingredient	%
$\alpha$ -pinene	0.29	spathulenol	0.37
$\alpha$ -thujene	0.30	b-caryophyllene oxide	1.01
$\alpha$ -terpinene	0.66	carvone	0.25
b-myrcene	0.79	$\alpha$ -terpineol	0.32
$\alpha$ -fellanderen	0.30	borneol	0.99
g-terpinen	2.39	b-bisabolen	2.08
p-simen	2.89	timol	4.15
Linalyl acetate	0.35	carvacrol	75.30
linalool	6.36		

blood was obtained. Samples taken were centrifuged at 3000 rpm for 15 min, and stored at -20°C until the analyses were carried out.

Superoxide dismutase, GPx and CAT antioxidant enzyme activities in plasma were determined by ELISA device (Epoch, Biotek, USA) using commercial kits (Cayman Chemical Co., USA). Whole blood reduced glutathione (GSH) analysis was determined colorimetrically (Epoch, Biotek, USA) according to the method issued by Beutler et al. [15]; while MDA in plasma by Yoshiko et al. [16], NO by Miranda et al. [17], ceruloplasmin by Colombo and Ricerich [18], and albumin and total protein levels by commercial test kit (Biolabo, France). The globulin was determined by subtraction of the albumin from the total protein according to Doumas et al. [19].

### Statistical Analysis

For the significance of the differences between the statistical calculations belonging to the groups and the mean values of the groups, one-way analysis of variance (ANOVA) method was used, and for the significance control of the difference between the groups, the appropriate next stage test (Tukey) was applied. For this purpose, SPSS packaged software was used [20].

## RESULTS

The amounts of nutrients and metabolizable energy values of concentrated feed and pasture grass are shown in *Table 3*. The performance parameters of the study are given in *Table 4*. In the study, statistically significant difference between the groups regarding average live weights, live weight gains, daily live weight gains, daily feed consumption and feed conversion ratio was not found ( $P > 0.05$ ). It has been observed that for the oregano oil use in post-weaned Tuj lambs, the difference between the groups, in terms of the performance parameters, was not statistically significant ( $P > 0.05$ ). It has been observed that the addition of oregano oil to lamb rations does not affect the performance parameters, but improves it.

Blood parameters of the study are given in *Table 5*. The differences between the groups on MDA, NO, GSH, CAT,

Feeds	DM	CP	EE	CF	CA	Calcium	Phosphorus	ADF	NDF	ME
Concentrate	87.36	16.25	2.36	11.03	7.99	0.64	0.42	-	-	2531
Pasture grass	92.1	9.13	-	-	7.6	0.65	0.16	38.70	62.60	1767

*DM: Dry matter, CP: Crude protein, EE: Ether extract, CF: Crude fiber, CA: Crude ash; ADF: Acid detergent fiber, NDF: Neutral detergent fiber, ME\*: Metabolic Energy (kcal/kg)*

Fattening Performance Parameters	Control		T1 1 mL/day/lamb		T2 1 mL/day/lamb		Significance
	X	Sx	X	Sx	X	Sx	P
<b>Live Weight, kg Days</b>							
0	34.58	0.23	34.83	0.42	34.83	0.52	0.886
10	35.81	0.25	36.13	0.34	36.06	0.47	0.814
31	38.18	0.2	38.7	0.23	38.93	0.34	0.160
52	41.05	0.07	41.4	0.33	41.65	0.44	0.441
<b>Live Weight Gain, kg, Days</b>							
0-10	1.23	0.12	1.3	0.11	1.23	0.12	0.889
10-31	2.36	0.08	2.56	0.19	2.86	0.24	0.195
31-52	2.86	0.23	2.7	0.18	2.71	0.18	0.818
0-52	6.46	0.25	6.56	0.37	6.81	0.48	0.805
<b>Daily Live Weight Gain, g Days</b>							
0-10	123.33	12.01	130	11.25	123.33	12.01	0.889
10-31	112.7	3.82	122.22	9.25	136.51	11.64	0.195
31-52	136.5	11.24	128.57	8.60	129.36	8.98	0.818
0-52	124.39	4.98	126.28	7.24	131.09	9.29	0.805
<b>Daily Feed Consumption (dry matter), g Days</b>							
0-52	1511.77	43.56	1448.30	101.34	1441.01	97.95	0.814
<b>Feed Conversion Ratio Days</b>							
0-52	12.32	0.84	11.91	1.56	11.46	1.37	0.898

*K: Control, T1: 1mL/day/lamb oregano oil, T2: 2 mL/day/lamb oregano oil. Statistically not significant (P>0.05). All values are given as mean ± standard error of mean (SEM), (n=6)*

ceruloplasmin, albumin, total protein and globulin values of oregano oil use in weaned Tuj lambs were not statistically significant ( $P>0.05$ ). According to the results, in terms of SOD and GPx, the use of oregano oil in post-weaned Tuj breed lambs was found to be statistically significant ( $P<0.05$ ).

## DISCUSSION

The use of aromatic plants and extracts in ruminant breeding is increasing each passing day. It has been reported that essential oils stimulate the digestive system of animals and increase the efficiency of digestive secretions [21]. Due to these features, essential oils have a positive effect on performance parameters. In the study conducted, it was observed that the use of oregano oil in different doses in post-weaned Tuj lambs did not affect live weight gain, daily live weight gain, daily feed consumption, and feed conversion rate. There are many current studies using

thyme and other plant extracts that support the results of our study [22-26]. In another study, it has been reported that the use of *cinnamic aldehyde* or *carvacrol* (200 mg/kg diet) does not affect dry matter consumption, live weight gain, and feed conversion ratio parameters [27]. In contrast to the results obtained, it was reported in a different study that the use of oregano oil in lamb feeding did not affect the feed conversion rate, and that the daily feed consumption and daily live weight gain were higher [28]. There are also studies reporting that the use of plant extracts in lambs has positive effects on dry matter consumption, live weight gain and feed conversion parameters [29-31].

The observed differences between the results are considered to be affiliated with the factors such as: plant species from which the oregano oil is obtained, oregano oil dose, the *carvacrol* and *thymol* ratio in oregano oil, the composition of the ration, the roughage-weighted feeding method, race

**Table 5.** The effect of oregano oil on MDA, NO, GSH, CAT, SOD, GPx, ceruloplasmin, albumin, total protein and globulin

Oxidant-Antioxidant Parameters	Days	Control		T1 1 mL/day/lamb		T2 1 mL/day/lamb		Significance P
		X	Sx	X	Sx	X	Sx	
MDA (µmol/L)	10	2.70	0.04	2.75	0.03	2.77	0.04	0.479
	52	2.69	0.04	2.73	0.02	2.76	0.03	0.407
NO (µmol/L)	10	12.24	0.94	12.32	1.30	12.75	1.14	0.948
	52	12.48	1.10	12.93	1.01	12.97	0.97	0.934
GSH (mg/dL)	10	38.21	1.93	38.16	2.14	38.24	1.95	1.000
	52	37.84	1.67	37.51	1.89	39.80	2.89	0.736
SOD (U/mL)	10	1.06 <sup>c</sup>	0.07	1.72 <sup>b</sup>	0.07	2.40 <sup>a</sup>	0.08	<b>0.000*</b>
	52	1.02 <sup>c</sup>	0.07	1.70 <sup>b</sup>	0.07	2.27 <sup>a</sup>	0.02	<b>0.000*</b>
CAT (nmol/min/mL)	10	28.07	1.39	32.23	1.55	32.42	2.05	0.156
	52	27.80	0.89	30.36	1.18	31.81	1.35	0.077
GPx (nmol/min/mL)	10	297.31 <sup>b</sup>	8.47	311.08 <sup>ab</sup>	4.22	32.54 <sup>a</sup>	6.00	<b>0.046*</b>
	52	294.81 <sup>b</sup>	6.04	320.97 <sup>ab</sup>	7.10	321.72 <sup>a</sup>	7.41	<b>0.022*</b>
Ceruloplasmin (mg/dL)	10	21.75	1.46	22.12	1.54	19.48	1.40	0.413
	52	19.57	0.9	18	1.02	18.14	1.08	0.488
Albumin (g/dL)	10	2.9	0.06	2.91	0.11	2.94	0.08	0.944
	52	2.87	0.07	2.95	0.08	2.87	0.06	0.709
Total protein (g/dL)	10	7.15	0.05	7.06	0.08	7.19	0.05	0.404
	52	7.18	0.10	7.16	0.13	7.20	0.06	0.952
Globulin (g/dL)	10	4.25	0.1	4.15	0.16	4.24	0.07	0.819
	52	4.31	0.16	4.21	0.14	4.33	0.07	0.789

**K:** Control, **T1:** 1 mL/day/lamb oregano oil, **T2:** 2 mL/day/lamb oregano oil. Statistically not significant ( $P > 0.05$ ). All values are given as mean  $\pm$  standard error of mean (SEM), (n=6); <sup>a,b,c</sup> The differences between the mean values with a different letter in the same row were statistically significant ( $P < 0.05$ ) \*  $P < 0.001$

of the animal, oregano oil's administration route to the animal.

Oxidation events occur constantly in the metabolism of a living being, and reactive oxygen substances, taken from outside, accelerate these oxidation events. Antioxidants are substances that prevent autooxidation/peroxidation progress by reacting with the radicals very quickly [32]. Endogenous antioxidants are non-enzymatic antioxidants composed of glutathione, albumin and ceruloplasmin, and enzymatic antioxidants that form the SOD, CAT, GPx and glutathione reductase enzymatic defense line [33,34]. The SOD enzyme is the first line of the antioxidant defense system. It plays a critical role in eliminating superoxide radicals [35]. Glutathione peroxidase is found in the cytoplasm of cells, and protects cells against oxidative damage caused by H<sub>2</sub>O<sub>2</sub> [36]. In the present study, it was observed that the use of oregano oil in lambs did not affect MDA, NO, GSH, CAT, ceruloplasmin, albumin, total protein, and globulin values, but significantly affected SOD and GPx values. Compared to the control group, SOD and GPx values increased in direct proportion to the increasing doses of oregano oil. The research results are consistent with current studies using thyme and other plant extracts. In the study in which 11 different plant extracts, including thyme, were used for lamb feeding, it was reported that

the blood antioxidant activity was not affected [37]. Gumus et al.[24] reported in their study that the use of oregano oil in the lambs not effected liver SOD values but GSH and CAT values, and also affected GSH, SOD, and CAT enzyme activities in *M. longissimus dorsi* tissue muscle significantly. Furthermore, in some studies using lamb extract from lambs, glutathione peroxidase activity ( $P < 0.05$ ) has been positively affected and antioxidant parameters have been improved [26,31,38].

In conclusion, it was determined that the addition of oregano oil into the ration did not have a significant effect on fattening performance and MDA, NO, GSH, CAT, ceruloplasmin, albumin, total protein, and globulin values of blood oxidant-antioxidant parameters. SOD and GPx values were significantly affected by the addition of oregano oil in lamb rations at the point of examining the effect on blood oxidant-antioxidant balance. Oregano oil had no effect on performance parameters in lambs, but significantly affects SOD and GPx values in blood. Therefore, it is determined that blood oxidant-antioxidant balance parameter results; have the potential to protect the cells against oxidative damage caused by free radicals, are able to decrease the peroxidation by strengthening the antioxidant structure in blood, and can be effective in protecting the oxidative stress which decreases the efficiency and resistance of the animals.

In the light of the data obtained from this study, it was concluded that the use of oregano oil, which is among the natural herbal products as an alternative to antibiotics, in Tuj lambs will make an important contribution to the breeders and the researchers, who will work later in this field, since it clarifies the changes occurred in the performance and blood parameters.

## REFERENCES

- Jouany JP, Morgavi DP:** Use of 'natural' products as alternatives to antibiotic feed additives in ruminant production. *Animal*, 1, 1443-1466, 2007. DOI: 10.1017/S1751731107000742
- Patra AK, Saxena J:** The effect and mode of action of saponins on the microbial populations and fermentation in the rumen and ruminant production. *Nurt Res Rev*, 22 (2): 204-219, 2009. DOI: 10.1017/S0954422409990163
- Hart KJ, Yanez-Ruiz DR, Duval SM, McEwan NR, Newbold CJ:** Plant extract to manipulate rumen fermentation. *Anim Feed Sci Technol*, 147 (1-3): 8-35, 2008. DOI: 10.1016/j.anifeeds.2007.09.007
- Szumacher-Strabel M, Cieslak A:** Potential of phytofactors to mitigate rumen ammonia and methane production. *J Anim Feed Sci*, 19, 319-337, 2010. DOI: 10.22358/jafs/66296/2010
- Olivier GW:** The world market of oregano. In, Padulosi S (Ed): Oregano. 14. *Proceedings of the IPGRI International Workshop*. Italy, Rome, 142-146, 1997.
- Çakmakçı R, Erdoğan Ü, Kotan R, Erdoğan Y, Mete E:** Çoruh vadisinde origanum acutidens populasyonlarının uçucu yağ kompozisyonu ve oranlarındaki varyasyon. *Türkiye VIII. Tarla Bitkileri Kongresi*, 19-22 Ekim, Hatay, 311-314, 2009.
- Karık Ü, Öztürk M:** Türkiye dış ticaretinde tıbbi ve aromatik bitkiler. *Bahçe*, 38 (2): 21-31, 2009.
- Grosso C, Figueiredo AC, Burillo J, Mainar AM, Urieta JS, Barroso JG, Coelho JA, Palavra AMF:** Composition and antioxidant activity of *Thymus vulgaris* volatiles: Comparison between supercritical fluid extraction and hydrodistillation. *J Sep Sci*, 33, 2211-2218, 2010. DOI: 10.1002/jssc.201000192
- D'antuono LF, Galletti GC, Bocchini P:** Variability of essential oil content and composition of *Origanum vulgare* L. Populations from a north mediterranean area (Liguria Region, Northern Italy). *Ann Bot*, 86, 471-478, 2000. DOI: 10.1006/anbo.2000.1205
- Dündar Y, Aslan R:** Hekimlikte Oksidatif Stres ve Antioksidanlar. AKÜ Yayınları, Yayın No, 29, Uyum Ajans, Ankara, 2000.
- Farag RS, Daw ZY, Abo-Raya SH:** Influence of some spice essential oils on *Aspergillus parasiticus* growth and production of aflatoxins in a synthetic medium. *J Food Sci*, 54, 74-76, 1989. DOI: 10.1111/j.1365-2621.1989.tb08571.x
- NRC:** Nutrient Requirements of Small Ruminants. The National Academy Press, Washington D.C., USA, 2006.
- AOAC:** Official Methods of Analysis of AOAC International. 18<sup>th</sup> edn., Gaithersburg, USA, 2005.
- Goering H, Van Soest PJ:** Forage fiber analyses (apparatus, reagents, procedures, and some applications), US Agricultural Research Service, Washington, USA, 1970.
- Beutler E, Duron O, Kelly BM:** Improved method for the determination of blood glutathione. *J Lab Clin Med*, 61, 882-888, 1963.
- Yoshioka T, Kawada K, Shimada T, Mori M:** Lipid peroxidation in maternal and cord blood and protective mechanism against activated-oxygen toxicity in the blood. *Am J Obstet Gynecol*, 135, 372-376, 1979. DOI: 10.1016/0002-9378(79)90708-7
- Miranda KM, Espey MG, Wink DA:** A rapid, simple spectrophotometric method for simultaneous detection of nitrate and nitrite. *Nitric Oxide*, 5, 62-71, 2001. DOI: 10.1006/niox.2000.0319
- Colombo JP, Richterich R:** Zur bestimmung des caeruloplasmin im plasma [on the determination of ceruloplasmin in plasma]. *Schweiz Med Wochenschr*, 23, 715-720, 1964.
- Doumas BT, Watson WA, Biggs HG:** Albumin standards and the measurement of serum albumin with bromocresol green. *Clin Chim Acta*, 31, 87-96, 1971. DOI: 10.1016/0009-8981(71)90365-2
- SPSS:** Statistical Packages for the Social Sciences, 20<sup>th</sup> edn., IBM Inc, Chicago, USA, 2011.
- Simsek ÜG, Güler T, Çiftçi M, Ertas ON, Dalkılıç B:** Esans yağ karışımının (kekik, karanfil ve anason) broylerde canlı ağırlık, karkas ve etlerin duyuşal özellikleri üzerine etkisi. *YYÜ Vet Fak Derg*, 16 (2): 1-5, 2005.
- Ünal A, Kocabağlı N:** Effect of different dosages of oregano oil on performance and some blood parameters in lambs. *Ankara Univ Vet Fak Derg*, 61, 199-204, 2014.
- Baytok E, Kara K, Aksu T, Güçlü BK, Özkaya S, Denek N, Kamalak A, Kaya DA, Önel SE, Akçay A:** The effect of Mediterranean thyme (*Thymbra spicata* L var *spicata*) essential oil on fattening performance and ruminal parameters in lamb. *J Anim Feed Sci*, 26, 319-325, 2017.
- Gümüş R, Erol HS, İmik H, Halıcı M:** The effects of the supplementation of lamb rations with oregano essential oil on the performance, some blood parameters and antioxidant metabolism in meat and liver tissues. *Kafkas Univ Vet Fak Derg*, 23 (3): 395-401, 2017. DOI: 10.9775/kvfd.2016.16791
- Ozdoğan M, Ustundag AO, Yaralı E:** Effect of mixed feeds containing different levels of olive cake on fattening performance, carcass, meat quality and fatty acids of lambs. *Trop Anim Health Prod*, 49 (8): 1631-1636, 2017. DOI: 10.1007/s11250-017-1369-6
- Liu H, Li K, Zhao J, Deng W:** Effects of polyphenolic extract from *Eucommia ulmoides* Oliver leaf on growth performance, digestibility, rumen fermentation and antioxidant status of fattening lambs. *Anim Sci J*, 89 (6): 888-894, 2018. DOI: 10.1111/asj.12998
- Yang WZ, Benchaar C, Ametaj BN, Chaves AV, He ML, McAllister TA:** Effects of garlic and juniper berry essential oils on ruminal fermentation and on the site and extent of digestion in lactating cows. *J Dairy Sci*, 90, 5671-5681, 2007. DOI: 10.3168/jds.2007-0369
- Khamisabadi H, Kafilzadeh F, Charaie B:** Effect of thyme (*Thymus vulgaris*) or peppermint (*Mentha piperita*) on performance, digestibility and blood metabolites of fattening Sanjabi lambs. *Biharean Biol*, 10 (2): 118-122, 2016.
- Salem AZ, Kholif AE, Olivares M, Elghandour MM, Mellado M, Arece J:** Influence of *S. babylonica* extract on feed intake, growth performance and diet in vitro gas production profile in young lambs. *Trop Anim Health Prod*, 46 (1): 213-219, 2014. DOI: 10.1007/s11250-013-0478-0
- Bandeira PAV, Filho JMP, de Azevedo Silva AM, Cezar MF, Bakke OA, Silva UL, Borburema JB, Bezerra LR:** Performance and carcass characteristics of lambs fed diets with increasing levels of *Mimosa tenuiflora* (Willd.) hay replacing Buffel grass hay. *Trop Anim Health Prod*, 49 (5): 1001-1007, 2017. DOI: 10.1007/s11250-017-1291-y
- Zhao J, Li K, Su R, Liu W, Ren Y, Zhang C, Du M, Zhang J:** Effect of dietary Tartary buckwheat extract supplementation on growth performance, meat quality and antioxidant activity in ewe lambs. *Meat Sci*, 134, 79-85, 2017. DOI: 10.1016/j.meatsci.2017.07.016
- Dündar Y, Aslan R:** Hücre moleküler statüsünün anlaşılması ve fizyolojik önem açısından radikaller, antioksidanlar. *İnsizyon Cerrahi Tıp Bil Derg*, 2 (2): 134-142, 1999.
- Aydemir B, Karadağ Sarı E:** Antioksidanlar ve büyüme faktörleri ile ilişkisi. *Kocatepe Vet J*, 2 (2): 56-60, 2009.
- Sen S, Chakraborty R:** The role of antioxidants in human health. In, *Oxidative Stress: Diagnostics, Prevention and Therapy*. Chapter-1, 1-37, ACS Symposium Series, Vol. 1083, 2011. DOI: 10.1021/bk-2011-1083.ch001
- Halıcı M, İmik H, Koc, Gumus R:** Effects of  $\alpha$ -lipoic acid, vitamins E and C upon the heat stress in Japanese quails. *J Anim Physiol Anim Nutr*, 96, 408-415, 2012. DOI: 10.1111/j.1439-0396.2011.01156.x
- Sen S, Chakraborty R, Sridhar C, Reddy YSR, De B:** Free radicals, antioxidants, diseases and phytomedicines: Current status and future prospect. *Int J Pharm Sci Rev Res*, 3 (1): 91-100, 2010.
- Leal LN, Jordán MJ, Bello JM, Otal J, Den Hartog LA, Hendriks WH, Martín-Tereso J:** Dietary supplementation of 11 different plant extracts on the antioxidant capacity of blood and selected tissues in lightweight lambs. *J Sci Food Agric*, 99, 4296-4303, 2019. DOI: 10.1002/jsfa.9662
- Zhang Y, Luo H, Liu K, Jia H, Chen Y, Wang Z:** Antioxidant effects of liquorice (*Glycyrrhiza uralensis*) extract during aging of *longissimus thoracis* muscle in Tan sheep. *Meat Sci*, 105, 38-45, 2015. DOI: 10.1016/j.meatsci.2015.03.002