

# Screening of Chinese Medicinal Herbs for Anthelmintic Efficacy Against *Gyrodactylus kobayashii* (Monogenea) in Goldfish (*Carassius auratus*)

Kaiqi LIAN <sup>1,a</sup> Mingliang ZHANG <sup>1,b</sup> Lingling ZHOU <sup>1,c</sup> Yuwei SONG <sup>1,d</sup> Xianjun GUAN <sup>1,e</sup>

<sup>1</sup> Academician Workstation of Animal Disease Control and Nutrition Immunity in Henan Province, Henan Joint International Research Laboratory of Veterinary Biologics Research and Application, School of Biotechnology and Food Science, Anyang Institute of Technology, Anyang 455000, CHINA

<sup>a</sup> ORCID: 0000-0003-0681-207X; <sup>b</sup> ORCID: 0000-0002-9512-8781; <sup>c</sup> ORCID: 0000-0002-9512-8781; <sup>d</sup> ORCID: 0000-0002-7012-6405

<sup>e</sup> ORCID: 0000-0002-3096-0593

Article ID: KVFD-2020-23196 Received: 16.08.2019 Accepted: 31.12.2019 Published Online: 06.01.2020

## How to Cite This Article

Lian K, Zhang M, Zhou L, Song Y, Guan X: Screening of Chinese medicinal herbs for anthelmintic efficacy against *Gyrodactylus kobayashii* (Monogenea) in goldfish (*Carassius auratus*). *Kafkas Univ Vet Fak Derg*, 2020 (Article in Press). DOI: 10.9775/kvfd.2020.23196

## Abstract

Monogenean infection can cause high mortality and significant financial losses in commercially farmed fish. Existing chemical drugs for the treatment of such infections often have serious drawbacks. In order to find alternative agents of chemical drugs, fourteen medicinal plants were tested for their in vivo anthelmintic activity against *Gyrodactylus kobayashii* in goldfish (*Carassius auratus*). Ethanol extracts of *Evodia rutaecarpa*, *Cnidium monnieri* and *Sophora flavescens* had 100% anthelmintic efficacy at low concentrations (100, 100 and 300 mg/L, respectively), after 48 h of exposure. The ethyl acetate extract of *C. monnieri* was the most effective, with an EC<sub>50</sub> value of 11.0 mg/L, after 48 h of exposure, showing 100% anthelmintic efficacy against *G. kobayashii* at 50.0 mg/L. Higher anthelmintic activity was also observed for remaining extracts of *C. monnieri* and *E. rutaecarpa* except for the water extracts of the two plants. The ethyl acetate extract of *C. monnieri* had the highest therapeutic index (TI, LC<sub>50</sub>/EC<sub>50</sub>) value of 31.8, and the lowest EC<sub>50</sub>, which indicates that this extract was the safest to goldfish among all extracts. For the remaining extracts of *C. monnieri* and *E. rutaecarpa*, the 48-h LC<sub>50</sub> values were about 10-fold higher than the corresponding EC<sub>50</sub> values. This result indicates these extracts have low toxicity in goldfish. The ethyl acetate extract of *C. monnieri* was the most effective and the safest among the tested extracts. Therefore, the two plants are expected to be alternative agents to control monogenean infection.

**Keywords:** Disease control, *Gyrodactylus kobayashii*, Chinese Medicinal Herb, Goldfish, Anthelmintic efficacy

## Japon Balıklarında (*Carassius auratus*) *Gyrodactylus kobayashii*'ye (Monogenea) Karşı Çin Şifalı Otlarının Anthelmintik Etkinliklerinin İncelenmesi

## Öz

Monogenean enfeksiyonları, ticari olarak yetiştirilen balıklarda yüksek mortalite ve önemli finansal kayıplara neden olabilir. Bu tür enfeksiyonların tedavisi için mevcut kimyasal ilaçların genellikle ciddi dezavantajları vardır. Kimyasal ilaçların alternatif ajanlarını belirleyebilmek için on dört şifalı bitki, akvaryum balığındaki (*Carassius auratus*) *Gyrodactylus kobayashii*'ye karşı in vivo antelmintik aktiviteleri açısından test edildi. *Evodia rutaecarpa*, *Cnidium monnieri* ve *Sophora flavescens*'in etanol özleri, 48 saat maruziyetten sonra düşük konsantrasyonlarda (sırasıyla 100, 100 ve 300 mg/L) %100 antelmintik etkinliğe sahipti. *C. monnieri*'nin etil asetat özütü, 48 saat maruziyetten sonra EC<sub>50</sub> değeri 11.0 mg/L ile etkili oluyordu ve 50.0 mg/L'de *G. kobayashii*'ye karşı %100 antelmintik etkinlik gösterdi. *C. monnieri* ve *E. rutaecarpa*'nın su özütleri dışındaki özütleri için de daha yüksek antelmintik aktivite gözlemlendi. *C. monnieri*'nin etil asetat ekstresi, en yüksek terapötik indeks (TI, LC<sub>50</sub>/EC<sub>50</sub>) değeri 31.8 ve en düşük EC<sub>50</sub>'ye sahipti, ve bu bulgu bu ekstraktın tüm ekstraktlar arasında akvaryum balığı için en güvenli olduğunu gösterdi. *C. monnieri* ve *E. rutaecarpa*'nın geri kalan ekstraktları için 48 saatlik LC<sub>50</sub> değerleri, ilgili EC<sub>50</sub> değerlerinden yaklaşık 10 kat daha yüksekti. Bu sonuç, bu ekstraktların Japon balıklarında düşük toksisiteye sahip olduğunu gösterdi. *C. monnieri*'nin etil asetat özütü, test edilen özütler arasında en etkili ve en güvenli olanıydı. Bu nedenle, iki bitki Monogenean enfeksiyonlarının kontrolünde alternatif ajanlar olabilir.

**Anahtar sözcükler:** Hastalık kontrolü, *Gyrodactylus kobayashii*, Çin şifalı bitki, Japon balığı, Antelmintik etkinlik

## INTRODUCTION

Ornamental fish trade is one of the most rapidly developing

areas of the aquaculture industry, and the value of international trade exports of ornamental fish has grown by an average of about 14% per year since 1985 <sup>[1,2]</sup>. The



İletişim (Correspondence)



+86 372 2909876



mingliang6969@sina.com

rapid development of the ornamental fish industry has been overwhelmed by the occurrence of large-scale parasitic diseases. Goldfish (*Carassius auratus*) is one of most popular ornamental fish worldwide, due to its easy maintenance and its attractive coloration. The most common ectoparasites infecting goldfish are gyrodactylids<sup>[3]</sup>. Viviparous gyrodactylids have a direct life-cycle. They can directly spread among hosts by contact, and undergo continuous transmission throughout their life-cycle<sup>[4,5]</sup>. Serious infection can cause evident clinical symptoms<sup>[3]</sup>. Monogeneans can damage the fish epidermis, which results in secondary infections by other pathogenic microorganisms<sup>[6,7]</sup>.

It is difficult to control monogeneans. Many chemical drugs have been used against these parasites, but bring some drawbacks (e.g., low efficacy, toxicity to host and environmental and human health problems)<sup>[8,9]</sup>. Additionally, the long-term use of chemical drugs can lead to drug resistance in parasites<sup>[10]</sup>. Recently, increasing attention has been paid to the use of traditional plant-based medicines to control diseases in aquaculture<sup>[11,12]</sup>. Zhou et al.<sup>[13]</sup> showed that herbal medicines were able to control *Gyrodactylus kobayashii* (*G. kobayashii*) infection in goldfish. Huang et al.<sup>[14]</sup> screened plant extracts with anthelmintic activity against *Dactylogyrus intermedius* (Monogenea) in goldfish.

In order to find safe and efficient alternatives to treat monogenean infection in aquaculture, we evaluated the anthelmintic efficacy of ethanol extracts of 14 plants against *G. kobayashii* in goldfish, using *in vivo* anthelmintic efficacy assays.

## MATERIAL and METHODS

### **Establishment of a Goldfish-*G. kobayashii* Model and Animal Ethics**

Goldfish weighing  $3.7 \pm 0.7$  g were selected from a fish farm in Anyang city, Henan province, China. All experiments complied with institutional animal care guidelines and were approved by the Animal Care Committee of Anyang Institute of Technology and Academician Workstation of Animal Disease Control and Nutrition Immunity in Henan Province (License no. SCXK(AWADCNI)2018-0001). These goldfish were fed and treated to remove all ectoparasites, as described previously<sup>[13]</sup>. A goldfish experimental infection model using *G. kobayashii* was performed according to a method described previously<sup>[15]</sup>. Uninfected fish were anesthetized with 0.02% MS-222. In order to allow parasite transfer between hosts, the caudal fins of uninfected fish were placed in contact with the caudal fins of heavily-infected fish reared in our laboratory. Experimentally-infected fish were then placed in a 1 L container. The success of the infection was determined by daily examination of the fins using a stereomicroscope. Ten days after infection, two parasites were collected from the infected fish for morphological and molecular identification according

to Li et al.<sup>[16]</sup> in order to verify that the infective agent was *G. kobayashii*. Uninfected goldfish were introduced periodically into an 80 L aquarium containing infected goldfish, in order to increase the number of infected fish for the experiments.

### **Preparation of Plant Materials**

The fourteen Chinese medicinal herbs listed in *Table 1* were purchased from a drug store and prepared according to a method described previously<sup>[13]</sup>. The herbal medicines were washed thoroughly, air-dried and oven-dried at 45°C for 48 h. In order to ensure the complete removal of water, dry plant materials were powdered, strained using a filter net of 30-40 mesh (450-600  $\mu$ m), and freeze-dried at -54°C. The powder of all the medicinal plants was extracted with ethanol. The powder of *E. rutaecarpa*, *C. monnieri* and *S. flavescens* was extracted for 48 h with water, methanol, petroleum ether, or ethyl acetate; the extraction process was repeated three times. To obtain solidified crude extracts, the extract solutions were filtered, and then concentrated at reduced pressure in a vacuum rotary evaporator, until all solvents had evaporated. Solid extracts then were dissolved in dimethyl sulfoxide (DMSO) at 0.5 mg/L as stock solutions for anthelmintic efficacy assays.

### **Screening Experiment**

The dry powder (50.0 g) of herbal medicines was extracted three times with 500.0 mL ethanol for 48 h. Ethanol filtrates were evaporated at reduced pressure in a vacuum rotary evaporator in order to obtain solidified crude extracts<sup>[14]</sup>. The solidified crude extract of each plant was dissolved in DMSO as a stock solution at 0.5 g/mL; these stock solutions were then screened for anthelmintic efficacy.

In addition, three kinds of herbal medicines (*E. rutaecarpa*, *C. monnieri* and *S. flavescens*) with higher antiparasitic activities (*Table 1*) were fractionated with different solvents (water, methanol, petroleum ether, and ethyl acetate), and then tested in anthelmintic efficacy assays. Each solid extract was dissolved in DMSO to obtain the stock solution at the concentration of 0.5 g/mL, which was used in anthelmintic efficacy assays.

### **In vivo Anthelmintic Efficacy Assay**

The anthelmintic assay against *G. kobayashii* in goldfish was performed according to a previous method<sup>[13]</sup>. Two goldfish infected with *G. kobayashii* were placed in 20×12×10 cm tanks containing 0.5 L of the test solutions at  $23 \pm 1^\circ\text{C}$  for 48 h. Two control groups without extract, or with the highest percentage of DMSO, were set at the same experimental conditions. All treatments and control groups were performed using five replicates. Before manipulation, goldfish were anesthetized with 0.02% MS-222, and the parasites in the caudal fin were counted under a stereomicroscope.

**Table 1.** The tested medicinal plants and the anthelmintic efficacy (AE) against *Gyrodactylus kobayashii* in goldfish of the ethanol extracts

Species	Plant Part Used	Mean Intensity (±SD)	The Best AE (%)	The Concentrations of the Best AE (mg/L)	The Concentrations of Fish Died (mg/L)
<i>Evodia rutaecarpa</i> (Juss.) Benth.	Fruit	102.6±45.2	100	100	500
<i>Cnidium monnieri</i> (L.) Cuss.	Fruit	87.6±35.4	100	100	200
<i>Sophora flavescens</i> Ait.	Root	95.9±50.4	100	300	500
<i>Areca catechu</i> L.	Seed	101.8±39.6	100	150	150
<i>Citrus reticulata</i> Blanco	Bark	96.9±51.5	100	400	800
<i>Mentha haplocalyx</i> Briq.	Leaf	117.8±40.1	100	400	1000
<i>Punica granatum</i> L.	Bark	105.8±42.4	100	600	800
<i>Agrimonia pilosa</i> Ledeb.	Aerialparts	109.1±37.1	100	600	600
<i>Omphalia lapidescens</i> Schroet.	Sclerotium	112.3±43.4	100	800	800
<i>Quisqualis indica</i> L.	Fruit	82.7±30.4	94.7	800	1000
<i>Dryopteris crassirhizoma</i> Nakai	Root, stem and leaf	101.7±25.7	89.0	200	200
<i>Pharbitis nil</i> (L.) Choisy	Seed	95.9±28.5	54.4	200	200
<i>Stemona sessilifolia</i> (Miq.) Miq.	Root	91.1±24.0	43.3	700	700
<i>Cynanchum paniculatum</i> (Bge) Kitaga	Rhizome	98.1±39.1	9.9	300	300

The anthelmintic efficacy of each extract was calculated according to the following formula:  $E = (L - L_t) / L \times 100\%$  for  $L > L_t$ ,  $E = 0$  for  $L \leq L_t$ .  $E$  is anthelmintic efficacy,  $L$  is the number of *G. kobayashii* on the caudal fin before treatment, and  $L_t$  is the number of surviving parasites after the treatment. Mean anthelmintic efficacy was calculated from five replicates per treatment.

### Acute Toxicity Tests

The extract with the strongest anthelmintic efficacy in *in vivo* anthelmintic efficacy assay, was tested for safety to goldfish. Acute toxicity tests were conducted in 26.5×16.5×12.5 cm tanks, containing ten healthy goldfish, and 2 L of aerated tap water at 23±1°C. Fish in tanks without plant extracts constituted the control group. Three replicates were set for the group of control fish and for the groups of fish exposed to plant's extracts. Fish mortality in each tank was recorded after 48 h. Fish were not fed during the experiment, and dead fish were continuously removed in order to avoid deterioration of water quality.

### Statistical Analyses

The homogeneity of replication samples was evaluated through the Mann-Whitney U test. At the 95% confidence level, median lethal concentration (LC<sub>50</sub>, LC<sub>90</sub>) and median effective concentration (EC<sub>50</sub>, EC<sub>90</sub>) were determined using the probit analysis. The therapeutic index (TI) was calculated as LC<sub>50</sub>/EC<sub>50</sub>. Each statistical analysis was performed using SPSS 19.0. A value of  $P < 0.05$  was considered significant and  $P < 0.01$  was considered highly significant.

## RESULTS


### Anthelmintic Efficacy of Extracts Against *G. kobayashii* *in vivo*

As shown in Table 1, of the 14 medicinal plants selected,

only the ethanol extracts of *E. rutaecarpa*, *C. monnieri* and *S. flavescens* had 100% anthelmintic efficacy at 100.0, 100.0 and 300.0 mg/L, respectively, after 48 h of exposure. Additionally, the extracts of *Citrus reticulata*, *Mentha haplocalyx*, *Punica granatum*, *Agrimonia pilosa* and *Omphalia lapidescens* showed good anthelmintic efficacy, but only at high concentrations (Table 1). There was either very weak anthelmintic activity, or highly toxicity to goldfish, in the remaining ethanol extracts of the herbal medicines tested. In the control groups, the number of *G. kobayashii* increased on the caudal fin (from 98.2±25.7 to 160.5±42.7 in the group with no extract, and from 113.1±21.5 to 150.5±31.3 in the DMSO group).



For the different extracts of *E. rutaecarpa*, *C. monnieri* and *S. flavescens*, their anthelmintic efficacy is showed in Fig. 1, 2, 3, and the corresponding EC<sub>50</sub> and EC<sub>90</sub> values are showed in Table 2. The ethyl acetate extract of *C. monnieri* exhibited 100% anthelmintic efficacy against *G. kobayashii* at 50.0 mg/L, and was the most effective against *G. kobayashii*, with an EC<sub>50</sub> value of 11.1 mg/L, and an EC<sub>90</sub> value of 28.3 mg/L, after 48 h of exposure. In addition, the anthelmintic activity against *G. kobayashii* was also determined for the methanol and petroleum ether extracts of *C. monnieri*, with EC<sub>50</sub> values of 23.6 and 44.6 mg/L, and EC<sub>90</sub> values of 51.0 and 84.0 mg/L, respectively. In all the extracts of *C. monnieri*, the water extract had the weakest anthelmintic efficacy of 42.7% at 1 000.0 mg/L.

The ethyl acetate extract of *E. rutaecarpa* had good anthelmintic efficacy, with EC<sub>50</sub> and EC<sub>90</sub> values of 24.0 and 50.3 mg/L, respectively, after a 48-h exposure. The petroleum ether and methanol extracts of *E. rutaecarpa* showed 100% anthelmintic efficacy at 150.0 and 120.0 mg/L, with EC<sub>50</sub> and EC<sub>90</sub> values of 71.9 and 108.3 mg/L (petroleum ether), and 40.9 and 91.2 mg/L (methanol), respectively. The water extract of *E. rutaecarpa* had the weakest anthelmintic efficacy of 25.6% at 800.0 mg/L. All



**Fig 1.** Anthelmintic activity of the different extracts of *Evodia rutaecarpa* against *Gyrodactylus kobayashii* after the exposure for 48 h. MEE, methanol extract; EAE, ethyl acetate extract; PEE, petroleum ether extract; WAE, water extract

**Fig 2.** Anthelmintic activity of the different extracts of *Cnidium monnieri* against *Gyrodactylus kobayashii* after the exposure for 48 h. MEE, methanol extract; EAE, ethyl acetate extract; PEE, petroleum ether extract; WAE, water extract



**Fig 3.** Anthelmintic activity of the different extracts of *Sophora flavescens* against *Gyrodactylus kobayashii* after the exposure for 48 h. MEE, methanol extract; EAE, ethyl acetate extract; PEE, petroleum ether extract; WAE, water extract

the extracts of *S. flavescens* showed effective anthelmintic efficacy against *G. kobayashii* after 48 h of exposure, but at high concentrations, with  $EC_{50}$  values of 234.6 mg/L (methanol), 224.4 mg/L (petroleum ether) and 238.6 mg/L (ethyl acetate).

#### Acute Toxicity of the Tested Extracts in Goldfish

The results of the acute toxicity assay of the extracts with higher anthelmintic efficacy at the low concentrations (methanol, petroleum ether, and ethyl acetate extracts of

*E. rutaecarpa* and *C. monnieri*) are summarized in Table 3. After 48 h of exposure, the  $LC_{50}$  values of the extracts of *E. rutaecarpa* were 759.3 (methanol), 771.9 (petroleum ether) and 235.9 (ethyl acetate) mg/L, which is 18.6, 10.7 and 9.9 times higher than the corresponding  $EC_{50}$ , respectively. The methanol, petroleum ether, and ethyl acetate extracts of *C. monnieri* had  $LC_{50}$  values of 214.1, 405.3 and 350.0 mg/L, which is 9.1-, 9.1- and 31.8 times higher than the corresponding  $EC_{50}$ , respectively. The water extracts of *E. rutaecarpa* and *C. monnieri* exhibited very weak toxicity to goldfish, and killed only two and

**Table 2.** Anthelmintic efficacy ( $EC_{50}$  and  $EC_{90}$ ) of different extracts from *Evodia rutaecarpa*, *Cnidium monnieri* and *Sophora flavescens* against *Gyrodactylus kobayashii* after 48 h of exposure

Plants	Extraction Solvent	$EC_{50}$ (mg/L)	95% CI	$EC_{90}$ (mg/L)	95% CI	P value
<i>Evodia rutaecarpa</i>	Methanol	40.9	0-56.0	91.2	76.3-131.1	0.06
	Petroleum ether	71.9	66.7-76.2	108.3	102.8-115.7	0.23
	Ethyl acetate	24.0	20.8-26.9	50.3	46.0-56.0	0.38
<i>Cnidium monnieri</i>	Methanol	23.6	15.8-28.4	51.0	47.6-55.8	0.46
	Petroleum ether	44.6	37.8-49.8	84.0	78.5-91.4	0.25
	Ethyl acetate	11.1	7.7-13.5	28.3	25.82-31.69	0.94
<i>Sophora flavescens</i>	Methanol	234.6	161.5-303.0	319.8	266.0-543.6	0.07
	Petroleum ether	224.4	0-292.8	400.0	328.1-714.7	0.11
	Ethyl acetate	238.6	0-315.3	416.4	335.9-914.6	0.06

$EC_{50}$ , effective concentration with 50% anthelmintic efficacy;  $EC_{90}$ , effective concentration with 90% anthelmintic efficacy; 95% CI, 95% confidence interval

**Table 3.** Acute toxicity for goldfish exposed to different extracts from *Evodia rutaecarpa* and *Cnidium monnieri* after 48 h of exposure

Plants	Extraction Solvent	$LC_{50}$ (mg/L)	95% CI	$LC_{90}$ (mg/L)	95% CI	TI ( $LC_{50}/EC_{50}$ )	P value
<i>Evodia rutaecarpa</i>	Methanol	759.3	713.5-806.6	854.7	807.3-975.2	18.6	0.45
	Petroleum ether	771.9	719.6-826.5	892.8	835.5-1039.1	10.7	0.32
	Ethyl acetate	235.9	172.6-298.8	385.9	317.2-579.7	9.9	0.70
	Water	>1000.0	-	-	-	-	-
<i>Cnidium monnieri</i>	Methanol	214.1	186.8-240.2	274.6	246.7-344.5	9.1	0.70
	Petroleum ether	405.3	332.0-462.8	552.9	488.5-735.8	9.1	0.30
	Ethyl acetate	350.0	301.5-398.5	453.6	403.7-579.8	31.8	0.59
	Water	>1000.0	-	-	-	-	-

$LC_{50}$ , 50% lethal concentration;  $LC_{90}$ , 90% lethal concentration; 95% CI, 95% confidence interval; TI, therapeutic index; -, not calculated

one goldfish, respectively, at the concentration of 1000.0 mg/L.

## DISCUSSION

Diseases caused by the common ectoparasites, *Gyrodactylus*, can result in high mortality and huge financial losses in commercially farmed fish [6,17]. Additionally, prolonged and frequent use of common chemicals such as formaldehyde, rotenone, and praziquantel results in increasing drug resistance and adverse effects on the environment [8,10]. Therefore, there is an urgent need for effective strategies to control *Gyrodactylus* infections. In our study, the ethyl acetate extract of *C. monnieri* was the most efficient, having the lowest  $EC_{50}$  and  $EC_{90}$  values (11.0 and 28.3 mg/L, respectively) after 48 h of exposure, and it had 100% anthelmintic efficacy against *G. kobayashii* at 50.0 mg/L. The dried fruit of *C. monnieri* (L.) Cuss., called "Shechuangzi" in Chinese, is a commonly used traditional Chinese medicine with an impressive range of health benefits [18]. Previous research has found that the ethanol extract had better anthelmintic efficacy against *Dactylogyrus intermedius* (*D. intermedius*) in goldfish, and had the minimal effective concentration compared to the chloroform, petroleum ether, water, and acetone extracts of *C. monnieri*. This

extract had 100% anthelmintic efficacy at 70.0 mg/L after 48 h of exposure. The active compounds of the ethanol extract against *D. intermedius* were identified as osthol and isopimpinellin [19,20]. In the present study, the ethyl acetate extract of *C. monnieri* was the most effective against *G. kobayashii* in goldfish at the minimal concentration, showing 100% anthelmintic efficacy at 50.0 mg/L, which was far lower than the concentration of ethanol extract with 100% anthelmintic efficacy (100.0 mg/L). Differences in anthelmintic efficacy of the same extracts may be due to the differences in the evaluation methods. In previous research, anthelmintic efficacy was calculated by comparison with the control group. In our paper, the number of *G. kobayashii* was recorded before and after exposure, and anthelmintic efficacy was calculated based on the changes in intensity of *Gyrodactylus* in goldfish. Besides, the difference in the main parasitic sites may be another cause. The main parasitic site of *Dactylogyrus* sp. is the gill, which may provide partial protection for this parasite by reducing the exposure to extracts. Nevertheless, *Gyrodactylus* sp. is mainly parasitic on the fins or the skin, which directly exposes this parasite to the extracts.

"Wuzhuyu", derived from the dried fruit of *E. rutaecarpa* (Juss.) Benth., is well described in the Chinese medical matter

and the Chinese pharmacopoeia, and it is recommended for the treatment of dizziness, headache, emesis, diarrhea and other symptoms<sup>[21,22]</sup>. Pharmacological studies indicate that the bioactive constituents of *E. rutaecarpa* have many properties, including anti-inflammatory, antihypertensive, antinociceptive, antimicrobial and anthelmintic activities<sup>[23,24]</sup>. For example, Perrett and Whitfield<sup>[25]</sup> reported that atanine (3-Dimethylallyl-4-methoxy-2-quinolone), an alkaloid isolated from the hexane extract of *E. rutaecarpa* caused 100% immobility of the cercarial and miracidial larvae of *Schistosoma mansoni* at 100.0 mg/L, after 7 min of exposure. Liu et al.<sup>[26]</sup> have shown that ethanol extracts of *E. rutaecarpa* are effective at killing intestinal nematode parasites of pigs and leeches *in vitro*. Also, the n-hexane extract of *E. rutaecarpa* has been demonstrated to have strong antifeedant activity against the grain storage insects *Sitophilus zeamais* and *Tribolium castaneum*<sup>[26]</sup>. Moreover, two alkaloids isolated from the methanol extract of *E. rutaecarpa*, evodiamine and rutaecarpine, have shown insecticidal activity against fruit flies (*Drosophila melanogaster*), with LC<sub>50</sub> values of 3.9 and 3.6 mg per adult, respectively<sup>[27]</sup>. In the present study, the ethyl acetate extract of *E. rutaecarpa* showed good anthelmintic efficacy, with EC<sub>50</sub> and EC<sub>90</sub> values of 24.0 and 50.3 mg/L, after 48 h of exposure. Although the active compounds against *G. kobayashii* that are contained in the ethyl acetate extract have not been identified, some of the compounds described above are thought to jointly or independently control *G. kobayashii* infection.

The therapeutic index (TI), a parameter for the quantitative relationship between efficacy and safety, is used to evaluate the potential application of the extracts in aquaculture<sup>[28]</sup>. TI is calculated as the ratio of LC<sub>50</sub> to EC<sub>50</sub>, and higher TI values reflect safer effects. In our study, we found that the ethyl acetate extract of *C. monnieri* had the highest TI value of 31.8, and the lowest EC<sub>50</sub>, which indicated this extract was safest to goldfish among all extracts and had the potential to be used in aquaculture. Additionally, the methanol extract had a higher TI value of 18.6, but the EC<sub>50</sub> was also higher compared with the ethyl acetate extract for *E. rutaecarpa*. The 48-h LC<sub>50</sub> values of the remaining extracts of *C. monnieri* and *E. rutaecarpa* were about 10-fold higher than the corresponding EC<sub>50</sub> values, which reflects low toxicity to goldfish. The bioactive ingredients that play a major role in anthelmintic activity are necessary for further bioassay-guided isolation and identification. In addition, field trials need to be performed before the use of extracts from *C. monnieri* and *E. rutaecarpa* in aquaculture.

Among the 14 Chinese medicinal herbs screened, the extracts of *C. monnieri* and *E. rutaecarpa* had the highest anthelmintic efficacy against *G. kobayashii* in goldfish. The ethyl acetate extract of *C. monnieri* was the most effective and the safest of all the extracts tested. *C. monnieri* and *E. rutaecarpa* may become novel therapeutic agents against *G. kobayashii* infection.

## ACKNOWLEDGEMENTS

This work was supported by the following funding: the National Natural Sciences Foundation of China (31802170), Colleges Key Scientific Research Projects of Henan Province (19B230001), and PhD Research Start-up Foundation of Anyang Institute of Technology (BSJ2016008, BSJ 016013).

## COMPETING INTERESTS

The authors declare they have no competing interests.

## REFERENCES

- 1. Tlustý M:** The benefits and risks of aquacultural production for the aquarium trade. *Aquaculture*, 205 (3-4): 203-219, 2002. DOI: 10.1016/S0044-8486(01)00683-4
- 2. FAO:** FishStat Plus 2010. Aquaculture Production Statistics: 1997–2008. Food and Agricultural Organisation of the United Nations, Fisheries and Aquaculture Department, Rome, Italy, 2010.
- 3. Cone DK, Odense PH:** Pathology of five species of *Gyrodactylus nordmanni*, 1832 (Monogenea). *Can J Zool*, 62 (6): 1084-1088, 1984. DOI: 10.1139/z84-156
- 4. Jorgensen TR, Raida MK, Kania PW, Buchmann K:** Response of rainbow trout (*Oncorhynchus mykiss*) in skin and fin tissue during infection with a variant of *Gyrodactylus salaris* (Monogenea: Gyrodactylidae). *Folia Parasitol*, 56 (4): 251-258, 2009. DOI: 10.14411/fp.2009.029
- 5. Zhou S, Zou H, Wu SG, Wang GT, Marcogliese DJ, Li WX:** Effects of goldfish (*Carassius auratus*) population size and body condition on the transmission of *Gyrodactylus kobayashii* (Monogenea). *Parasitology*, 144 (9): 1221-1228, 2017. DOI: 10.1017/S0031182017000543
- 6. Bakke TA, Cable J, Harris PD:** The biology of gyrodactylid monogeneans: The "Russian-doll killers". *Adv Parasitol*, 64, 161-376, 2007. DOI: 10.1016/S0065-308x(06)64003-7
- 7. Zhang S, Zhi T, Xu X, Zheng Y, Bilong Bilong CF, Pariselle A, Yang T:** Monogenean fauna of alien tilapias (Cichlidae) in south China. *Parasite*, 26:4, 2019. DOI: 10.1051/parasite/2019003
- 8. Kang YJ, Wakabayashi C, Kim KH:** Anthelmintic potential of quinacrine and oxyclozanide against gill parasite *Microcotyle sebastis* in black rockfish *Sebastes schlegelii*. *Dis Aquat Organ*, 119, 259-263, 2016. DOI: 10.3354/dao02997
- 9. Forwood JM, Harris JO, Deveney MR:** Efficacy of bath and orally administered praziquantel and fenbendazole against *Lepidotrema bidyana* Murray, a monogenean parasite of silver perch, *Bidyana bidyanus* (Mitchell). *J Fish Dis*, 36 (11): 939-947, 2013. DOI: 10.1111/jfd.12089
- 10. Buchmann K, Roepstorff A, Waller PJ:** Experimental selection of mebendazole-resistant gill monogeneans from the European eel, *Anguilla anguilla* L. *J Fish Dis*, 15 (5): 393-408, 1992. DOI: 10.1111/j.1365-2761.1992.tb01238.x
- 11. Ling F, Jiang C, Liu G, Li M, Wang G:** Anthelmintic efficacy of cinnamaldehyde and cinnamic acid from cortex cinnamon essential oil against *Dactylogyrus intermedius*. *Parasitology*, 142 (14): 1744-1750, 2015. DOI: 10.1017/S0031182015001031
- 12. Militz TA, Southgate PC, Carton AG, Hutson KS:** Efficacy of garlic (*Allium sativum*) extract applied as a therapeutic immersion treatment for *Neobenedenia* sp. management in aquaculture. *J Fish Dis*, 37 (5): 451-461, 2014. DOI: 10.1111/jfd.12129
- 13. Zhou S, Li WX, Wang YQ, Zou H, Wu SG, Wang GT:** Anthelmintic efficacies of three common disinfectants and extracts of four traditional Chinese medicinal plants against *Gyrodactylus kobayashii* (Monogenea) in goldfish (*Carassius auratus*). *Aquaculture*, 466, 72-77, 2017. DOI: 10.1016/j.aquaculture.2016.09.048
- 14. Huang AG, Yi YL, Ling F, Lu L, Zhang QZ, Wang GX:** Screening of plant extracts for anthelmintic activity against *Dactylogyrus intermedius* (Monogenea) in goldfish (*Carassius auratus*). *Parasitol Res*, 112 (12): 4065-

4072, 2013. DOI: 10.1007/s00436-013-3597-7

**15. Harris PD:** Changes in the site specificity of *Gyrodactylus turnbulli* Harris, 1986 (Monogenea) during infections of individual guppies (*Poecilia reticulata* Peters, 1859). *Can J Zool*, 66 (12): 2854-2857, 1988. DOI: 10.1139/z88-414

**16. Li R, Li W, Wu X, Wang G:** Identification of gyrodactylus species in goldfish (*Carassius auratus*) through morphological study and the analysis of the rDNA ITS sequence. *Acta Hydrobiol Sin*, 38 (5): 903-909, 2014. DOI: 10.7541/2014.135

**17. Heggberget TG, Johnsen BO:** Infestations by *Gyrodactylus* sp of Atlantic salmon, *Salmo salar* L, in Norwegian rivers. *J Fish Biol*, 21 (1): 15-26, 1982. DOI: 10.1111/j.1095-8649.1982.tb02819.x

**18. Li YM, Jia M, Li HQ, Zhang ND, Wen X, Rahman K, Zhang QY, Qin LP:** *Cnidium monnieri*: A review of traditional uses, phytochemical and ethnopharmacological properties. *Am J Chinese Med*, 43 (5): 835-877, 2015. DOI: 10.1142/s0192415x15500500

**19. Wang GX, Zhou Z, Cheng C, Yao JY, Yang ZW:** Osthol and isopimpinellin from *Fructus cnidii* for the control of *Dactylogyrus intermedius* in *Carassius auratus*. *Vet Parasitol*, 158 (1-2): 144-151, 2008. DOI: 10.1016/j.vetpar.2008.07.034

**20. Wang G, Chao C, Chen A, Shen Y, Xing Z:** Dactylogyrus killing efficacies of 22 plants extracts and its 6 compounds. *Acta Botanica Boreali-Occidentalia Sinica*, 26, 2567-2573, 2006.

**21. Pan X, Wang M, Wu Y, Lu X, Shang Y, Xu Y, Zhai Y, Li J, Li Z, Gong M:** Identification of active ingredients in Wuzhuyu decoction improving migraine in mice by spectral efficiency association. *Mol Med Rep*, 12 (1):

1524-1534, 2015. DOI: 10.3892/mmr.2015.3506

**22. Pan XQ, Wu YC, Gong MX, Xu YS, Wang ZM, Zhang QW, Shang YW, Lu XR, Song YF:** Study on pharmacological ingredients of wuzhuyu tang treating migraine by correlating absorption ingredients in everted intestinal sac and pharmacodynamics. *Zhongguo Zhong Yao Za Zhi*, 39 (1): 126-133, 2014.

**23. Liao JF, Chiou WF, Shen YC, Wang GJ, Chen CF:** Anti-inflammatory and anti-infectious effects of *Evodia rutaecarpa* (Wuzhuyu) and its major bioactive components. *Chinese Med*, 6:6 2011. DOI: 10.1186/1749-8546-6-6

**24. Zhou X, Zhao Y, Lei PH, Cai ZW, Liu H:** Chromatographic fingerprint study on *Evodia rutaecarpa* (Juss.) Benth by HPLC/DAD/ESI-MS<sup>n</sup> technique. *J Sep Sci*, 33 (15): 2258-2265, 2010. DOI: 10.1002/jssc.201000035

**25. Perrett S, Whitfield PJ:** Atanine (3-dimethylallyl-4-methoxy-2-quinolone), an alkaloid with anthelmintic activity from the chinese medicinal plant, *Evodia rutaecarpa*. *Planta Med*, 61 (3): 276-278, 1995. DOI: 10.1055/s-2006-958073

**26. Liu ZL, Goh SH, Ho SH:** Screening of Chinese medicinal herbs for bioactivity against *Sitophilus zeamais* Motschulsky and *Tribolium castaneum* (Herbst). *J Stored Prod Res*, 43 (3): 290-296, 2007. DOI: 10.1016/j.jspr.2006.06.010

**27. Miyazawa M, Fujioka J, Ishikawa Y:** Insecticidal compounds from *Evodia rutaecarpa* against *Drosophila melanogaster*. *J Sci Food Agric*, 82, 1574-1578, 2002. DOI: 10.1002/jsfa.1215

**28. Muller PY, Milton MN:** The determination and interpretation of the therapeutic index in drug development. *Nat Rev Drug Discov*, 11, 751-761, 2012. DOI: 10.1038/nrd3801