

Association of Single Nucleotide Polymorphism in Bone Morphogenetic Protein Receptor 1B (BMPR-1B) Gene with Growth Traits in Chicken

Ashraf AWAD¹ Mahmoud Salah EL-TARABANY¹ 

¹ Department of Animal Wealth Development, Faculty of Veterinary Medicine, Zagazig University, Sharkia - EGYPT

Article Code: KVFD-2015-13515 Received: 03.04.2015 Accepted: 13.07.2015 Published Online: 16.07.2015

Abstract

Growth traits are under the control of multiple genes. Understanding the genetic information of related genes is helpful for the selection and breeding course through marker assisted selection. The aim of the current study was to investigate the association of A287G SNP in BMPR-1B gene with growth traits in chicken. A single nucleotide polymorphism was identified in 240 individuals using the PCR-RFLP technique and confirmed by sequencing. The allelic and genotypic frequencies were compared, using the Chi-squared test. Associations between the genotype of each polymorphism and the traits were analyzed using the General Linear Model of statistical software SAS. Three genotypes (AA, AG and GG) were detected in Fayoumi and Rhold Island Red chicken. Sequencing revealed one mutation (287 A→G) in the genotype AA in comparison to the genotype GG. The A287G SNP of BMPR-1B gene was associated significantly with body weight at 2nd (P=0.022), 3rd (P=0.034), 4th (P=0.011), 5th (P=0.035), 6th (P=0.001), 7th (P=0.008) and 8th (P=0.016) week of age. In conclusion, BMPR-1B gene may be associated with body weight in chicken and may be considered in Marker Assisted Selection program to improve chicken growth performance.

Keywords: Chicken, BMPR-1B, SNP, Growth traits

Tavuklarda Kemik Morfogenetik Protein Reseptörü 1B (BMPR-1B) Genindeki Tek Nükleotid Polimorfizmi ile Büyüme Özellikleri Arasındaki İlişki

Özet

Büyüme özellikleri çok sayıda genin kontrolü altındadır. İlgili genlerdeki genetik bilgiyi anlamak, marker-destekli seleksiyon yoluyla seçim ve üreme sürecinde yardımcı olur. Bu çalışmanın amacı, tavuklarda BMPR-1B genindeki A287G SNP ile büyüme özellikleri arasındaki ilişkiyi araştırmaktır. PCR-RFLP tekniği kullanılarak, 240 bireyde bir tek nükleotid polimorfizmi tespit edildi ve sekanslama ile doğrulandı. Allellik ve genotipik frekanslar Ki-kare testi kullanılarak karşılaştırıldı. Her polimorfizm genotipi ve özellikler arasındaki ilişkiler SAS istatistiksel yazılımının Genel Lineer Modeli kullanılarak analiz edildi. Fayoumi ve Rhold Adası Kırmızı tavuklarında üç genotipi (AA, AG ve GG) saptandı. Sıralama, genotip GG'ye kıyasla genotip AA'da bir mutasyon (287 A→G) olduğunu gösterdi. BMPR-1B geninin A287G SNP'si, vücut ağırlığı ile 2. (P=0.022), 3. (P=0.034), 4. (P=0.011), 5. (P=0.035), 6. (P=0.001), 7. (P=0.008) ve 8. haftalarda (P=0.016) önemli ölçüde ilişkili idi. Sonuç olarak, BMPR-1B geni tavuklarda vücut ağırlığı ile ilişkili olabilir ve Markör Destekli Seçim programında tavukların büyüme performansını artırmak için dikkate alınabilir.

Anahtar sözcükler: Tavuk, BMPR-1B, SNP, Büyüme özellikleri

INTRODUCTION

Growth and egg production traits of chicken are controlled by a series of major genes and/or quantitative trait loci (QTL). Analyses of genetic markers in animals could lead to discernment of the genetic architecture of quantitative traits. There are two basic methods of QTLs

identification: approach of the candidate gene and whole-genome scanning ^[1,2]. The candidate gene approach is an effective method for finding QTLs responsible for genetic variation in the traits of interest in agricultural animal species and calibrating whether specific genes are associated to economic traits in farm animals ^[3]. Several trials have been concerned in the fields of association



İletişim (Correspondence)



+20 122 3668785, Fax: +20 552283683



mahmoudtarabany2887@yahoo.com; mmohamedibrahim@zu.edu.eg

analysis between candidate gene SNPs with animal growth and body composition traits [3-7].

Fayoumi is a native chicken breed originated in Egypt, reared for meat and egg production. It is adapted to subtropical environmental conditions and performed well under intensive management conditions [8]. Rhode Island Red (RIR) is a dual-purpose breed of American class. It is well adapted to the local environmental conditions [9]. Genetic polymorphisms are playing an important role as genetic markers in many sectors of animal breeding. As the molecular genetic techniques developed, it has become possible to obtain a new class of gene markers based upon the variability at DNA sequence level. Application of these molecular genetic markers potentially will greatly enhance the intensity of selection and will most efficiently uncover the productive potential of birds. Additionally, marker-associated selection (MAS) based on the studies concerned candidate genes and their effect on the phenotypic manifestations. An important intent of modern breeding in the poultry industry is to synthesis high-performance poultry lines and breeds in two main directions of productivity, meat and eggs [10].

The bone morphogenetic proteins (BMPs) are members of the transforming growth factor b (TGFb) superfamily. They are multifunctional proteins that regularize growth and differentiation in many cell types and play fundamental functions during embryogenesis and the fertility in mammals [11,12]. BMPs are potent inducers of cartilage and bone formation and has an important role in the bone healing process and in improving therapeutic efficacy [13,14]. In multipotential mesenchymal cells derived from equine adipose cells, BMP-2 increase under magnetic field conditions which affect the osteogenic properties of the cells and enhance vascularization process [15]. A non-conservative substitution (Q249R) in the *BMPR-1B* sequence was related with the proliferation characteristics of some ewe breeds [16]. In the chicken ovary, granulosa cells are major target for BMPs and it was proposed that mRNA levels for *BMPR-1B* in granulosa cells are higher than in theca cells [17].

The objectives of the present study were to detect single nucleotide polymorphisms (SNPs) of *BMPR-1B* gene in Fayoumi and RIR chicken by PCR-RFLP and sequencing. In addition, investigating the association between these SNP and growth traits in chicken.

MATERIAL and METHODS

This study was carried out in accordance with the Zagazig University Animal Ethics Committee guidelines (ANWD-206), at the Biotechnology unit belonging to Department of Animal Wealth Development, Faculty of Veterinary Medicine, Zagazig University.

Experimental Flock and Management

A total of 300 day old chicks, Fayoumi and RIR breeds, were used in this study. Chicks were physically examined and wing banded. Brooding and growing requirements were provided using conventional floor system. All chicks were subjected to the same managerial, hygienic and climatic conditions. A standard diet was provided *ad libitum* at rearing period (8 weeks), including 22.5% crude protein and 2975.8 K.cal ME/kg. body weight to the nearest gram was recorded at hatch, 1st, 2nd, 3rd, 4th, 5th, 6th, 7th and 8th week of age.

DNA Extraction

A total of 240 blood samples were collected from plumage veins of Fayoumi and RIR chickens and conserved in tubes containing EDTA as an anticoagulant. DNA was isolated using Gene JET whole blood genomic DNA purification mini kit (Fermentas, Thermo Fisher Scientific, USA), following the manufacturer protocol. The quality of DNA was checked by running on 0.8% agarose gel and was quantified by reading absorbance at A260/A280 nm in a UV spectrophotometer.

PCR Amplification

A 581 bp fragment of *BMPR-1B* gene covering exon 6, intron 6 and exon 7 was amplified by polymerase chain reaction using Primers suggested by Zhang et al. [18] and DreamTaq Green PCR Master Mix (Thermo Scientific, fermentas, USA). PCR was carried out in a total volume of 25 µl reaction mixture containing 12.5 µl of Master Mix, 2 µl DNA, 1 µl each primer (10 µM), 8.5 µl deionized water and used T-professional thermal cycler (Biometra, Germany) according to the following program: initial denaturation at 95°C for 5 min; 40 cycles consisting of 30 s at 95°C, 30 s at 56°C and 30 s at 72°C for denaturation, annealing and extension steps, respectively; and a final extension at 72°C for 7 min. The PCR products were checked by electrophoresis on 1.5% agarose gel in 1×TAE buffer. The amplified product was visualized under UV transilluminator.

PCR-RFLP Analysis

A 20 µl of digestion mixture, consisting of 15 µl of the PCR product, 1×of recommended buffer, 2 µl of deionised water, and 6 units of restriction enzyme *Hind* III was placed in a 0.5 ml microcentrifuge tube. The digestion mixture was mixed thoroughly in a vortex mixer and incubated at 37°C for 15 min. The digested product was run on a 2% agarose gel in TAE buffer along with 100 bp plus DNA ladder as molecular size marker. Gels were visualized under UV light and the genotype patterns were screened in gel documentation system.

Nucleotide Sequencing

The desired PCR product band was excised using a clean, sterile razor blade or scalpel and excised quickly to

minimize exposure of the DNA to UV light. The minimum agarose slice was transferred to a 1.5 ml micro centrifuge or screw cap tube and then purified by using a commercially available gel extraction kit (Fermentas, Thermo Fisher Scientific, USA). Samples were labeled and sent for sequencing. Sequencing was done by European Custom Sequencing Centre (GATC Biotech AG, Germany) using both forward and reverse primers of PCR amplification. The obtained sequences were edited manually using ChromasLiteVer. 2.01, (http://www.tech_nelysium.com.au/chromas.html) and aligned with Clustal Omega software to identify nucleotide polymorphism.

Statistical Analysis

All statistical procedures were performed using SAS statistical system package V9.1 [19]. Allelic and genotypic

frequencies of the single nucleotide polymorphism (SNP) were calculated and Chi-Square test was performed to examine Hardy-Weinberg equilibrium. Marker-trait association analysis was conducted using the one-way analysis of variance (ANOVA) through the general linear models (GLM) procedure. The comparison of means was carried out with Duncan's multiple range tests.

RESULTS

PCR Amplification

Genomic DNA of the two chicken breeds was amplified using specific primers for *BMPR-1B* gene. PCR products were detected by running a 1.5% agarose gel electrophoresis (Fig. 1). The amplified products (581 bp)




Fig 1. PCR Amplification of chicken *BMPR-1B* gene exon 6 to exon 7. M: 100 bp plus ladder

Şekil 1. Tavuk *BMPR-1B* geni ekson 6'dan 7. M ekson'a kadar PCR Amplifikasyonu: 100 bp artı merdiveni




Fig 2. Genotyping of the A287G SNP in chicken *BMPR-1B* gene by (A) PCR-RFLP and (B) sequencing. Arrow indicates A/G substitution

Şekil 2. Tavuk *BMPR-1B* genindeki A287G SNP'nin (A) PCR-RFLP ve (B) sıralaması (sekanslaması) ile genotiplendirilmesi. Ok, A/G ikamesini belirtir




Table 1. Frequency of genotypes and alleles of *BMPR-1B* gene in Fayoumi and RIR chicken breeds**Tablo 1.** Fayoumi ve RIR tavuk ırklarındaki *BMPR-1B* geninin genotip ve allel sıklığı

Breed	Genotype Frequency (n)			Allele Frequency		χ^2
	AA	AG	GG	A	G	
Fayoumi	0.09 (11)	0.55 (66)	0.36 (43)	0.37	0.63	4.07 ¹
RIR	0.12 (14)	0.60 (72)	0.28 (34)	0.42	0.58	6.59 ²

¹ Significant at level ($P < 0.05$), ² Significant at level ($P < 0.01$)

Table 2. Least squares means of body weight (g) according to genotypes at the SNP A287G of *BMPR-1B* gene**Tablo 2.** *BMPR-1B* geninin SNP A287G'indeki genotiplere göre vücut ağırlığının (g) en küçük kareleri

Traits	Genotypes				
	AA	AG	GG	RSD	P-value
BW Day old (g)	28.7	28.6	27.2	2.7	0.652
BW 1 st wk (g)	71.3	73.9	68.3	8.1	0.559
BW 2 nd wk (g)	150.1 ^{ab}	156.8 ^a	126.6 ^b	12.9	0.022
BW 3 rd wk (g)	235.2 ^{ab}	242.9 ^a	200.5 ^b	22.7	0.034
BW 4 th wk (g)	320.4 ^a	328.7 ^a	266.6 ^b	24.6	0.011
BW 5 th wk (g)	382.5 ^{ab}	398.6 ^a	327.2 ^b	43.1	0.035
BW 6 th wk (g)	477.4 ^a	499.2 ^a	412.7 ^b	45.7	0.001
BW 7 th wk (g)	571.2 ^b	677.4 ^a	518.3 ^b	62.6	0.008
BW 8 th wk (g)	673.3 ^b	773.8 ^a	606.7 ^b	67.4	0.016

RSD: residual standard deviation

^{a,b} Values within a row with different superscripts differ significantly

were consistent with the target fragments and had a good specificity, which could be directly analyzed by RFLP and sequencing.

RFLP Analysis

The PCR-RFLP method was developed successfully for genotyping the A287G SNP in intron 6 of the chicken *BMPR-1B* gene, where all individuals have been screened. Three genotypes of AA, AG, and GG were detected and confirmed by sequencing (Fig. 2 A,B). The fragment sizes of 581 bp for the GG genotype, 294/287 for the AA genotype, and a combination of 581, 294 and 287 bp for AG genotype.

Genotyping and Frequencies

Allele and genotype frequencies of *BMPR-1B* gene were calculated within each breed (Table 1). The frequencies of GG genotype in Fayoumi and RIR chickens were 0.36 and 0.28, respectively; which obviously greater than AA frequencies. Therefore, the allele G was predominant in the populations (0.63 and 0.58, respectively). Chi-test showed that two chicken breeds were not in Hardy-Weinberg equilibrium, in which genotype frequencies had been distorted by recent selection, mutation, or migration.

Association of the *BMPR-1B* Genotypes with Growth Traits

The least squares means of body weight according to different genotypes of *BMPR-1B* gene in chicken populations were presented in Table 2. The A287G SNP of *BMPR-1B* gene is associated significantly with body weight at 2nd ($P=0.022$), 3rd ($p=0.034$), 4th ($P=0.011$), 5th ($P=0.035$), 6th ($P=0.001$), 7th ($P=0.008$) and 8th ($P=0.016$) week of age. Heterozygous genotype AG had a higher body weight than GG genotype over the whole experimental period. The clear significant differences between heterozygous AG genotype and AA genotype were detected ultimately at 7th and 8th week of age.

DISCUSSION

The primary objectives of the current study were to detect single nucleotide polymorphisms (SNPs) of *BMPR-1B* gene in Fayoumi and RIR chicken by PCR-RFLP and sequencing. Growth and body composition are an inclusive reflection of development of various parts of the chicken body and its final expression is the result of interaction among genetic, nutritional and environmental factors [20]. Growth is under complex genetic monitoring, and uncovering the molecular mechanism of growth

results in more efficient selection for growth in broiler chickens [21]. Identifying the QTL will facilitate poultry breeding programs for the economic important traits. Molecular genetic information is required to be used to consolidate genetic improvement of animal species. The candidate gene approach is a very powerful method to examine associations of gene polymorphisms with economically important traits in farm animals [22]. Application of breeding programs that utilize marker-assisted selection requires advances in some areas like detection and estimation of associations of identified genes and their genetic markers with economic traits. Phenotypic evaluation is critical to establish marker-assisted associations or carry out the candidate gene validations required to conduct MAS [23]. Up to now, majority of association studies, especially in chicken, have been performed using phenotypic information.

The A/G transition at the base position of 287 in *BMPR-1B* gene was investigated in two chicken breeds. The allele frequency of G was higher than that of A in those two breeds and was in the range that reported by Zhang et al. [18] in Zang chickens. While the allele frequency of A was higher than that of G in three Chinese native chickens, a synthetic broiler line [18] and Mazandaran native chicken [24]. Our results showed that A287G SNP of chicken *BMPR-1B* gene is associated significantly with body weight from the 2nd till the 8th week of age. On the contrary, Zhang et al. [18] and Niknafs et al. [24] recorded non-significant association between *BMPR-1B* gene and growth traits. Zhang et al. [18] stated that A287G SNP of chicken *BMPR-1B* is associated with egg production from 47 to 56 weeks. Previous studies showed that *BMPR-1B* gene as a well known effective gene for reproductive traits [16,25].

Chicken *BMPR-1B* mRNA sequences were first identified by Sumitomo et al. [26] and Lim et al. [27]. Lim et al. [27] reported that BMP signaling, including *BMPR-1B*, is involved in chick diencephalic development, and the expression level of *BMPR-1B* reduced in the theca of chicken ovary from F1 to F3 follicles. Onagbesan et al. [17] proposed that *BMPR-1B* is possibly concerned in follicular differentiation and maintenance of the follicular hierarchy. Therefore, the expression level or the activity of *BMPR-1B* in the granulosa and/or theca of chicken ovary may be associated with oocyte maturation.

BMPR-1B as a member of the transforming growth factor β (TGF- β) receptor superfamily played the important actions in signal transduction. The current model of induction of signalling responses is at the cell surface, the ligand binds a complex of transmembrane receptor serine/threonine kinases (types I and II) and incites transphosphorylation of the Gly-Ser (GS) segments in the type I receptor by the type II receptor kinases. The consequently activated type I receptors phosphorylate selected Smads at C-terminal serines, and these receptor activated Smads (R-Smads) then form a complex with a

common Smad4. Energetic Smad complexes translocate into the nucleus, where they regularize transcription of target genes, through physical interaction, CBP or p300 coactivators and functional cooperation with DNA-binding transcription factors (X) [28].

In conclusion, the broiler chickens have been subjected to intensive breeding with so many objectives that should be simultaneously considered to reduce costs, improve health and product quality. So, several traits such as growth and body composition have been included in selection policies. In addition to difficulty of measurement of these traits, the correlations among them are complex. MAS can be a perfect option to improve selection programs. The results from the current study indicated that a SNP marker in the *BMPR-1B* gene was associated with growth traits in chickens, therefore, a potential marker for molecular MAS programs in chicken. However, the conclusion was only preliminary; it was worth increasing the number of chicken breeds, and expanding the number of samples to make in-depth study.

ACKNOWLEDGEMENTS

We sincerely thank Prof. Dr. Khairy M. El-Bayoumi, and Dr. Bassanat Magdi for their kindful help and assistance.

CONFLICT OF INTEREST

The authors declare that they have no conflict of interest.

REFERENCES

1. Ikeobi CON, Woolliams JA, Morrice DR, Law A, Windso D, Burt DW, Hocking PM: Quantitative trait loci affecting fatness in the chicken. *Anim Genet*, 33, 428-435, 2002. DOI: 10.1046/j.1365-2052.2002.00911.x
2. Kim TH, Choi BH, Lee HK, Park HS, Lee HY, Yoon DH, Lee JW, Jeon GJ, Cheong IC, Oh SJ, Han JY: Identification of quantitative traits loci (QTL) affecting growth traits in pigs. *Asian Australas J Anim Sci*, 18, 1524-1528, 2005. DOI: 10.5713/ajas.2005.1524
3. Li H, Deeb N, Zhou H, Mitchell AD, Ashwell CM, Lamont SJ: Chicken quantitative trait loci for growth and body composition associated with transforming growth factor- β genes. *Poult Sci*, 82, 347-356, 2003. DOI: 10.1093/ps/82.3.347
4. Jiang YL, Fan XZ, Lu ZX, Tang H, Xu JQ, Du LX: Detection of 881A→881G mutation in tyrosinase gene and associations with the Black ear Coat Color in rabbits. *Asian-Australas J Anim Sci*, 15, 1395-1397, 2002. DOI: 10.5713/ajas.2002.1395
5. Yan B, Deng X, Fei J, Hu X, Wu C, Li N: Single nucleotide polymorphism analysis in chicken growth hormone gene and its associations with growth and carcass traits. *Chinese Sci Bull*, 48, 1561-1564, 2003. DOI: 10.1007/BF03183961
6. Zhang BZ, Lei MG, Deng CY, Xiong YH, Zuo B, Li FE: Association between PCR-RFLP polymorphism of the fifth intron in lipoprotein lipase gene and productive traits in pig resource family. *Asian Australas J Anim Sci*, 18, 458-462, 2005. DOI: 10.5713/ajas.2005.458
7. Meng H, Zhao JG, Li ZH, Li H: Single nucleotide polymorphism on peroxisome proliferators-activated receptor genes associated with fatness traits in chicken. *Asian-Australas J Anim Sci*, 18, 1221-1225, 2005. DOI: 10.5713/ajas.2005.1221
8. Hossary MA, Galal ESE: Improvement and adaptation of the Fayoumi chickens. *Anim Genet Resour*, 14, 33-39, 1994. DOI: 10.1017/

S1014233900000328

9. Ashraf M, Mahmood S, Ahmed F: Comparative reproductive efficiency and egg characteristics of Lypur silver Black and Rhode Island Red breeds of poultry. *Int J Agric Biol*, 5, 449-451, 2003.

10. Kulibaba RA, Podstreshnyi AP: Prolactin and growth hormone gene polymorphisms in chicken lines of Ukrainian selection. *Cytol Genet*, 46, 390-395, 2012.

11. Fang X, Xu H, Zhang C, Chen H, Hu X, Gao X, Gu C, Yue W: Polymorphism in BMP4 gene and its association with growth traits in goats. *Mol Biol Rep*, 36, 1339-1344, 2009. DOI: 10.1007/s11033-008-9317-1

12. Monget P, Fabre S, Mulsant P, Lecerf F, Elsen JM, Mazerbourg S, Pisselet C, Monniaux D: Regulation of ovarian folliculogenesis by IGF and BMP system in domestic animals. *Domest Anim Endocrinol*, 23, 139-154, 2002. DOI: 10.1016/S0739-7240(02)00152-2

13. Nishii N, Arai M, Yanai N, Togari A, Nakabayashi T: Effect of bone morphogenetic protein-2 (BMP-2) or troglitazone, as an inducer of osteogenic cells or adipocytes, on differentiation of a bone marrow mesenchymal progenitor cell line established from temperature-sensitive (ts) simian virus (SV) 40 T-antigen gene transgenic mice. *Biol Pharm Bull*, 32, 10-17, 2009.

14. Seol YJ, Kim KH, Park YJ, Lee YM, Ku Y, Rhyu IC, Lee SJ, Han SB, Chung CP: Osteogenic effects of bone-morphogenetic-protein-2 plasmid gene transfer. *Biotechnol Appl Biochem*, 49, 85-96, 2008. DOI: 10.1042/BA20060247

15. Monika M, Krzysztof M, Daniel L, Anna S, Agnieszka Ś: Static magnetic field enhances synthesis and secretion of membrane-derived microvesicles (MVs) rich in VEGF and BMP-2 in equine adipose-derived stromal cells (EqASCs) - A new approach in veterinary regenerative medicine. *In Vitro Cell Dev Biol Anim*, 51, 230-240, 2015. DOI: 10.1007/s11626-014-9828-0

16. Souza CJH, MacDougall C, Campbell BK, McNeilly AS, Baird DT: The Booroola (FecB) phenotype is associated with a mutation in the bone morphogenetic receptor type 1B (BMPR1B) gene. *J Endocrinol*, 169, R1-R6, 2001.

17. Onagbesan OM, Bruggeman V, Van As P, Tona K, Williams J, Decuypere E: BMPs and BMPRs in chicken ovary and effects of BMP-4 and -7 on granulosa cell proliferation and progesterone production *in vitro*.

Am J Physiol Endocrinol Metabol, 285, 973-83, 2003.

18. Zhang NB, Tang H, Kang L, Ma YH, Cao DG, Lu Y, Hou M, Jiang YL: Associations of single nucleotide polymorphisms in BMPR-1B gene with egg production in a synthetic broiler line. *Asian Australas J Anim Sci*, 21, 628-632, 2008.

19. SAS Institute: SAS/STAT users guide. SAS Institute Inc., Cary, NC, USA. 2003.

20. Scanes CG, Harvey S, Marsh JA, King DB: Hormones and growth in poultry. *Poult Sci*, 63, 2062-2074, 1984. DOI: 10.3382/ps.0632062

21. Deeb N, Lamont SJ: Genetic architecture of growth and body composition in unique chicken population. *J Hered*, 93, 107-118, 2002. DOI: 10.1093/jhered/93.2.107

22. Rothschild MF, Soller M: Candidate gene analysis to detect genes controlling traits of economic importance in domestic livestock. *Prob Newslet Agric Genom*, 8, 13-20, 1997.

23. Guimarães EP, Ruane J, Scherf BD, Sonnino A, Dargie JD: Marker-Assisted Selection. Food and Agricultural Organization of the United Nations, Rome, Italy, 427-440, 2007.

24. Niknafs S, Javaremi AN, Sadeghi M: Single nucleotide polymorphisms in BMPR-1B and STAT5B genes and their association with growth and reproductive traits in chicken. *Songklanakarin J Sci Technol*, 36, 137-142, 2014.

25. Wilson T, Wu XY, Juengel JL, Ross IK, Lumsden JM, Lord EA, Dodds KG, Walling GA, McEwan JC, O'Connell AR, McNatty KP, Montgomery GW: Highly prolific Booroola sheep have a mutation in the intracellular kinase domain of bone morphogenetic protein 1B receptor (ALK-6) that is expressed in both oocytes and granulosa cells. *Biol Reprod*, 64, 1225-1235, 2001.

26. Sumitomo S, Saito T, Nohno T: A new receptor protein kinase from chick embryo related to type II receptor for TGF-beta. *DNA Seq*, 3, 297-302, 1993.

27. Lim Y, Cho G, Minarcik J, Golden J: Altered BMP signaling disrupts chick diencephalic development. *Mech Dev*, 122, 603-620, 2005. DOI: 10.1016/j.mod.2004.12.001

28. Derynck R, Zhang YE: Smad-dependent and Smad independent pathways in TGF-beta family signaling. *Nature*, 425, 577-584, 2003. DOI: 10.1038/nature02006