

The Effect of Single and Combined Use of Probiotic and Humate in Quail (*Coturnix coturnix Japonica*) Diet on Fattening Performance and Carcass Parameters

Tarkan ŞAHİN *  Dilek AKSU ELMALI ** İsmail KAYA * Mehmet SARI *** Özlem KAYA *

* Department of Animal Nutrition and Nutritional Diseases, Faculty of Veterinary Medicine, University of Kafkas, TR-36100 Kars - TURKEY

** Department of Animal Nutrition and Nutritional Diseases, Faculty of Veterinary Medicine, University of Mustafa Kemal, TR-31034 Hatay - TURKEY

*** Department of Animal Science, Faculty of Veterinary Medicine, University of Kafkas, TR-36100 Kars - TURKEY

Makale Kodu (Article Code): KVFD-2010-640

Summary

This experiment was carried out to determine the both single and combined effects of humates (Farmagulator XP™) and probiotics (Biosacc™) in quail diets on fattening performance and carcass yields. A total of 300 one-day old Japanese Quails (*Coturnix coturnix japonica*) were used in this experiment. Animals were divided into one control group and three treatment groups containing 75 birds each. Each group was further divided into three replicates containing 25 birds each. The experimental period lasted for 35 days. The control group was fed with unsupplemented basal diets. The rations of treatment groups were supplemented with 1 g/kg Farmagulator XP™ (Group H), 0.5 g/kg Biosacc™ (Group B) and 1 g/kg Farmagulator XP™ + 0.5 g/kg Biosacc™ combination (Group H+B), respectively. At the end of the study there were no statistically significant differences among the groups in terms of body weight gain, feed consumption, feed conversion ratio and carcass yield ($P>0.05$). But, body weight scores were found higher at the 4th and 5th weeks in the group H ($P<0.05$). It is concluded that the use of probiotic and humic acid alone and combination has no additional effect on quail performance.

Keywords: *Quail, Humic acide, Probiotic, Performance, Carcass*

Bıldırcın Rasyonlarına Propiyotik ve Humik Asitin Yalnız ve Kombine Katılmasının Besi Performansı ve Karkas Kalitesine Etkisi

Özet

Bu araştırma, bıldırcın rasyonlarına ilave edilen humik asit (Farmagulator XP™) ve probiyotik (Biosacc™)'ın, yalnız veya kombine kullanımının besi performansı ve karkas kalitesine etkilerini belirlemek amacıyla yapılmıştır. Araştırmada toplam 300 adet günlük Japon bıldırcın (*Coturnix coturnix japonica*) civcivi kullanılmıştır. Her grupta 75 civciv bulunan bir kontrol ve 3 deneme grubu oluşturulmuştur. Gruplar kendi aralarında 3'erli alt gruba ayrılmıştır. Deneme beş hafta sürdürülmüştür. Kontrol grubu basal rasyonla beslenmiştir. Deneme grubu rasyonlarına sırasıyla 1 g/kg Farmagulator XP™ (Grup H), 0.5 g/kg Biosacc™ (Grup B) ve 1 g/kg Farmagulator XP™ + 0.5 g/kg Biosacc™ (Grup H+B) kombinasyonu ilave edilmiştir. Araştırma sonunda, canlı ağırlık artışı, yem tüketimi, yemden yararlanma oranı ve karkas verimleri bakımından istatistik açıdan önemli bir fark görülmemiştir ($P>0.05$). Bununla beraber canlı ağırlık değişimleri 4 ve 5. haftalarda Grup H' de daha yüksek bulunmuştur ($P<0.05$). Sonuç olarak, probiyotik ile humik asitin yalnız ve birlikte kullanılmasının performansta ilave bir artışa yol açmadığı görülmüştür.

Anahtar sözcükler: *Bıldırcın, Humik asit, Probiyotik, Performans, Karkas*



İletişim (Correspondence)



+90 474 2426807/1137



tarkants7@hotmail.com

INTRODUCTION

Humates, originated from decomposed plants in the soil, have a very complex structure with molecular weight ranging from 5.000 to 200.000. Humates are composed of humic, ulmic and fulvic acids. Humic acids have ingredients of carbohydrates, aminoacids and fenolic compounds¹⁻³. They are also long chain heavy molecules which are able to transfer electrons. By this way they can play important roles in excreting toxic compounds from the body^{1,4}.

To enhance nutrient utilization, improve feed conversion efficiency, and maintain health status, inclusion of probiotics and humates in rations is preferable to antibiotics primerly, because they cause no harmful effects on consumers^{5,6}. Because of this reason, the effects of humic acids on fattening performance and some carcass parameters are investigated⁷⁻¹⁰.

Probiotic is a generic term, and these products can contain yeast cells, bacterial cultures, or both that stimulate microorganisms capable of modifying the gastrointestinal environment to favour health status and improve feed efficiency¹¹. Therefore, the major outcomes from using probiotics include improvement in growth and feed conversion ratio¹².

Although the exact mechanisms of probiotics are not clearly identified, their effects varies in terms of the microorganism species, dose, animal species and environment^{7,13}. Studies with broilers¹⁴, laying hens¹⁵, market turkeys¹⁶, and turkey breeder hens¹⁷⁻¹⁹ have shown that the incorporation of a yeast culture into the feeds resulted in enhanced performance.

This experiment was carried out to determine the effect of both single and combination of humates (Farmagulator XP™) and probiotics (Biosacc™) in Japanese Quails diets on body weight (BW), body weight gain (BWG), feed consumption (FC), feed conversion ratio (FCR) and carcass yield.

MATERIAL and METHODS

A total of 300 one-day old Japanese quail chicks (*Coturnix coturnix japonica*) of both gender were randomly divided into 4 experimental groups of 75 birds in each group with 3 replicates following measurement of body weighing. The chicks were housed in electrically heated battery cages and exposed to light for 24 h from hatching to 5 weeks of age. The experimental period lasted for 35 days. The control and experimental groups were fed with a basal diet of including 21.50% CP

and 3050 kcal/kg ME. Diets were formulated to meet NRC²⁰ nutrient requirements. The experimental design consisted of four dietary treatments: 1) Control group was fed with unsupplemented basal diets; 2) 1 g/kg Farmagulator XP in diet (Grup H); 3) 0.5 g/kg Biosacc in diet (Grup B); 4) 1 g/kg Farmagulator XP + 0.5 g/kg Biosacc combination in diet (Grup H+B). Quails consumed the diets and water ad libitum.

Farmagulator XP used in this study contained humic, ulmic and fulvic acids, and organic minerals. Biosacc's ingredient was *Saccharomyces cerevisiae* (2.5 x 10⁹ cfu/g). The compositions of diets are shown in [Table 1](#). Chemical compositions of feeds were analyzed by the methods of AOAC²¹. The metabolisable energy levels of rations were calculated according to TSE²².

The body weight and body weight gain of the quails were determined at the beginning (0) and 7, 14, 21, 28 and 35th days of the study. At the same time all the replicates feed residues were weekly weighed to define the feed consumption levels and feed conversion ratios. Following measurement of individual body weights, a total of 40 chicks, 5 male and 5 female birds from each group, were randomly chosen and slaughtered for determining the carcass yield at the end of the study. Hot carcass weights were determined after slaughter and cold carcass weight was determined after storage at

Table 1. Composition and calculated analysis of basal ration

Tablo 1. Temel Rasyonun bileşimi ve hesaplanan analiz değerleri

| Ingredient | (%) | Analyzed Contents of Nutrients |
|--------------------------|--------|--------------------------------|
| Maize | 60.50 | Dry matter, %90.17 |
| Soybean meal | 29.50 | Crude protein, %21.50 |
| Fish meal | 4.00 | Crude extract, %5.78 |
| Oil | 3.30 | Crude fiber, %3.35 |
| Limestone | 1.20 | Crude ash, %3.02 |
| Dicalcium phosphate | 0.50 | N- free extract, %56.52 |
| Salt | 0.30 | ME**, kcal/kg 3050 |
| DL- Methionine | 0.10 | |
| L- Lysine | 0.10 | |
| Vitamin -mineral premix* | 0.50 | |
| Total | 100.00 | |

* 1 kg : 20.000.000 IU Vit A, 3.000.000 IU Vit D₃, 25 g Vit E, 4 g Vit B₁, 8 g Vit B₂, 5 g Vit B₆, 20 mg Vit B₁₂, 20 g Nicotinamide, 12 g Ca-D-Pantothenate, 200 g Choline Chloride, 50 g Mn, 50 g Fe, 50 g Zn, 10 g Cu, 0.8 g I, 0.15 g Co, 0.15 g Se

** Calculated

Supplemented of the treatment groups:

1 g/kg Farmagulator XP™ (Group H),

0.5 g/kg Biosacc™ (Group B)

1 g/kg Farmagulator XP™ + 0.5 g/kg Biosacc™ combination (Group H+B)

+4°C for 18 h. Hot and cold carcass yield rates were determined as follow; dividing hot and cold carcass weights by body weight before the slaughter.

Statistical analyses of body weight and carcass parameters of the groups were determined by analysis of variance. Duncan test was used to determine differences between treatment groups. Kruskal Wallis test was employed to determine the effect of groups on BWG, FC and FCR²³. The statistical analyses were performed of SPSS 10.0 (Inc. Chicago. IL. USA).

RESULTS

The body weights are shown in [Table 2](#). Body weight gain, feed consumption and feed conversion ratio are

shown in [Table 3](#). The carcass yield values are shown in [Table 4](#).

Body weights of quail chicks in this study were found statistically higher than the that of group H at the fourth and fifth weeks of the study. Body weight gain, feed consumption, feed conversion ratio and carcass parameters values were not statistically significant between all groups.

DISCUSSION

Although there was no statistically significant difference between the control and experimental groups in the first 3 weeks of the study a significant improvements in body weight of the humic acid supplemented (Group

Table 2. Mean body weights of groups (g)

Tablo 2. Gruplarda ortalama canlı ağırlıklar (g)

| Age (Weeks) | Control X±Sx | Group H X±Sx | Group B X±Sx | Group H+B X±Sx | P |
|-------------|--------------|--------------|--------------|----------------|---|
| 0 | 8.11±0.07 | 8.41±0.10 | 8.31±0.11 | 8.19±0.10 | - |
| 1 | 17.12±0.43 | 17.31±0.49 | 16.74±0.48 | 17.00±0.46 | - |
| 2 | 37.13±1.24 | 36.82±1.05 | 35.95±1.00 | 35.65±0.94 | - |
| 3 | 67.30±1.67 | 69.80±1.67 | 64.56±1.61 | 66.65±1.65 | - |
| 4 | 108.13±2.38b | 118.27±2.15a | 104.23±1.85b | 108.99±2.04b | * |
| 5 | 136.67±2.83b | 148.34±2.69a | 139.51±2.33b | 137.11±2.28b | * |

a,b: Differences between values having different letters in the same line are statistically significant ($P<0.05$)

- : Differences among the groups were not statistically significant ($P>0.05$)

Table 3. Mean body weight gain (g), feed consumption (g), and feed conversion ratio* in groups

Tablo 3. Gruplarda ortalama canlı ağırlık artışları (g), yem tüketimleri (g/civciv) ve yemden yararlanma oranı*

| Age (Weeks) | Parameters | Control X±Sx | Group H X±Sx | Group B X±Sx | Group H+B X±Sx | P |
|-------------|-----------------------|--------------|--------------|--------------|----------------|---|
| 1 | Feed consumption | 18.57±0.18 | 19.76±0.23 | 18.48±0.28 | 19.60±0.46 | - |
| | Body weight gain | 9.01±0.41 | 8.90±0.74 | 8.41±0.72 | 8.81±0.46 | - |
| | Feed conversion ratio | 1.92±0.09 | 2.25±0.16 | 2.24±0.23 | 2.47±0.15 | - |
| 2 | Feed consumption | 39.92±2.40 | 45.61±0.73 | 42.73±0.74 | 43.43±1.98 | - |
| | Body weight gain | 20.00±1.14 | 19.80±0.49 | 18.97±0.99 | 18.63±0.50 | - |
| | Feed conversion ratio | 2.00±0.05 | 2.31±0.07 | 2.26±0.09 | 2.34±0.15 | - |
| 3 | Feed consumption | 82.83±1.34 | 86.17±1.58 | 81.03±1.97 | 79.89±1.83 | - |
| | Body weight gain | 30.24±2.48 | 32.71±0.20 | 28.91±0.53 | 30.93±1.40 | - |
| | Feed conversion ratio | 2.78±0.27 | 2.63±0.06 | 2.80±0.02 | 2.60±0.15 | - |
| 4 | Feed consumption | 128.21±4.79 | 142.86±1.14 | 130.28±3.82 | 127.46±3.06 | - |
| | Body weight gain | 40.79±0.92 | 48.43±0.48 | 39.48±4.48 | 42.40±0.34 | - |
| | Feed conversion ratio | 3.14±0.10 | 2.95±0.03 | 3.38±0.35 | 3.01±0.10 | - |
| 5 | Feed consumption | 111.67±3.70 | 128.87±4.26 | 119.82±1.49 | 121.38±2.10 | - |
| | Body weight gain | 28.59±1.00 | 30.12±1.65 | 35.66±6.38 | 28.22±1.32 | - |
| | Feed conversion ratio | 3.92±0.27 | 4.29±0.12 | 3.57±0.58 | 4.32±0.19 | - |
| 1-5 | Feed consumption | 385.84±9.78 | 432.75±3.57 | 405.14±9.34 | 417.08±7.99 | - |
| | Body weight gain | 128.64±3.52 | 139.96±1.56 | 131.44±2.58 | 128.98±2.50 | - |
| | Feed conversion ratio | 3.00±0.11 | 3.09±0.01 | 3.08±0.02 | 3.23±0.03 | - |

* (kg, feed consumption /kg, body weight gain)

- : Differences among the groups were not statistically significant ($P>0.05$)

Table 4. Mean live weight, carcass weight and yields of experimental groups**Tablo 4.** Grupların ortalama kesim ve karkas ağırlıkları (g) ile karkas randımanları (%)

| Parameters | Control X±Sx | Group H X±Sx | Group B X±Sx | Group H+B X±Sx | P |
|------------------------|-----------------|-----------------|-----------------|-------------------|---|
| Female | | | | | |
| Live weight (g) | 152.19±2.78 | 148.25±4.38 | 150.97±2.93 | 153.98±2.96 | - |
| Warm carcass (g) | 103.47±1.87 | 101.53±3.92 | 104.26±2.60 | 106.57±2.03 | - |
| Cold carcass (g) | 101.83±1.69 | 98.42±3.48 | 100.12±1.92 | 103.20±2.53 | - |
| Warm carcass yield (%) | 68.01±0.68 | 68.42±0.85 | 69.03±0.49 | 69.26±1.22 | - |
| Cold carcass yield (%) | 66.93±0.62 | 66.34±0.52 | 66.33±0.54 | 67.02±0.97 | - |
| Male | | | | | |
| Live weight(g) | 143.88±1.15 | 141.02±1.66 | 143.85±0.92 | 145.60±3.98 | - |
| Warm carcass (g) | 101.57±1.18 | 98.32±1.40 | 98.97±2.02 | 102.31±3.07 | - |
| Cold carcass (g) | 98.88±1.53 | 97.32±1.27 | 96.63±2.37 | 98.39±3.71 | - |
| Warm carcass yield (%) | 70.59±0.47 | 69.73±0.74 | 68.78±1.00 | 70.26±0.77 | - |
| Cold carcass yield (%) | 68.72±0.84 | 69.02±0.75 | 67.15±1.28 | 67.51±1.11 | - |

- : Differences among the groups were not statistically significant ($P>0.05$)

H) group at 4th and 5th weeks ($P<0.05$) was noted. Feeding humate during the grower period had the most beneficial effect on broiler performance in terms of growth. This may be explained by the fact that humic acid stabilizes the intestinal microflora and thus ensures an improved utilization of nutrients in animal feed²⁴. It is defined that the supplementation of humic acid enhances the growth performance of quails. This study is in accordance with the studies where humic acid had effect ($P<0.05$) on body weight gains on broilers^{8,10}. On the other hand, insignificant differences between the control and experimental groups (Group B ve Group H+B) in terms of body weight is paralell to the studies where *S. cerevisiae* supplementation to diet of the quail²⁵ and turkey^{19,26} diets did not affect body weight. The results of the studies on humate and probiotic addition to the broiler diets^{27,28} revealed similar results to ours.

In this study, feed consumption and feed conversion ratio were determined as 385.84, 432.75, 405.14, 417.08 g and 3.19, 3.15, 3.14 and 2.87, respectively. Feed consumption was found lower in control group than the treatment groups but this was not significant. Our results were in agreement with the studies in which probiotic and humate addition single or in combination did not affect the feed consumption^{8,10,25,27,29,30}. Also, the results from the studies conducted with humate⁹, humate and probiotic²⁸, L-carnitine and sodium humate³¹, antibiotic, prebiotic, humic acid mixture^{25,32} were similar to the results of this study.

No statistical significant difference among the treatment groups with respect to hot and cold carcass yields was similar to previous results obtained from studies where humate/humic acid⁸⁻¹⁰, humate and probiotic combined and/or separately²⁸ were added to broiler diets^{8,10,25,33,34}.

It is concluded that the use of probiotic and humic acid alone or in combination has no additional effect on quail performance.

REFERENCES

- Anonim:** The effects of humates on crops. <http://www.organicapproach.com/humates.htm>, 2002. Accessed: 30.10.2009.
- Faust RH:** Humic acids- humates. <http://www.agrihelp.com/humates.htm>, 1996. Accessed: 22.09.2009.
- Levinsky BV:** All about humate effects and benefits. <http://www.unic.com/>, 1997. Accessed: 22.09.2009.
- Spark KM, Wells JD, Johnson BB:** The interaction of a humic acid with heavy metals. *Aust J Soil Res*, 35, 89-101, 1997.
- Fedorova LA, Dudinov VM, Levinsky BV:** Experimental and commercial application of potassium/sodium humate (Gumat) at State Production Company Fowl Farm Severy. http://www.alt.irkutsk.ru/humate/result_e.html, 1998. Accessed: 30.01.2010.
- Onifade AA, Odunsi AA, Babatunde GM, Olorede BR, Muma E:** Comparison of the supplemental effects of *Saccharomyces cerevisiae* and antibiotics in low-protein and high-fiber diets fed to broiler chickens. *Arch Tierernahr*, 52, 29-39, 1999.
- Bailey CA, White KE, Donke SL:** Evaluation of menefee humate on performance of broilers. *Poult Sci*, 75 (Suppl): 84 (Abstr), 1996.
- Eren M, Deniz G, Gezen ŞŞ, Türkmen İİ:** Broiler yemlerine katılan humatların besi performansı, serum mineral konsantrasyonu ve kemik küllü üzerine etkileri. *Ankara Univ Vet Fak Derg*, 47, 275-317, 2000.
- Karaoğlu M, Macit M, Esenbuga N, Durdag H, Turgut L, Bilgin ÖC:** Effect of supplemental humate at different levels on the growth performance, slaughter and carcass traits of broilers. *Int J Poult Sci*, 3 (6): 406-410, 2004.
- Kocabağlı N, Alp M, Acar N, Kahraman R:** The effects of dietary humate supplementaion on broiler growth and carcass yield. *Poultry Sci*, 81, 227-230, 2002.
- Dierck NA:** Biotechnology aids to improve feed and feed digestion: Enzymes and fermentation. *Arch Anim Nutr Berl*, 39, 241-261, 1989.

12. **Yeo J, Kim K:** Effect of feeding diets containing an antibiotic, a probiotic, or yucca extract on growth and intestinal urease activity in broiler chicks. *Poult Sci*, 76, 381-385, 1997.
13. **Jin LZ, Ho YW, Abdullah N, Jalaludin S:** Probiotics in poultry; modes of action. *World's Poult Sci J*, 53, 351-368, 1997.
14. **Thayer RH, Jackson CD:** Improving phytate phosphorus utilization by poultry with live yeast culture. *Okla Agric Exp Stn Res Rep*, MP-103, 131-139, 1975.
15. **Tonkinson LV, Gleaves EW, Dunkelgod KE, Thayer RH, Sirny RJ, Morrison RD:** Fatty acid digestibility in laying hens fed yeast culture. *Poult Sci*, 44, 159-164, 1965.
16. **Savage TF, Nakaue HS, Holmes ZA:** Effects of feeding a live yeast culture on market turkey performance and cooked meat characteristics. *Nutr Rep Int*, 31, 695-703, 1985.
17. **Thayer RH, Burkitt RF, Morrison RD, Murray EE:** Efficiency of utilization of dietary phosphorus by caged turkey breeder hens fed rations supplemented with live yeast culture. *Okla Agric Exp Stn Res Rep*, MP-103, 173-181, 1978.
18. **Savage TF, Mirosh LW:** Breeder performance of Medium White turkey hens fed diet containing a 2.5% yeast culture. *Poult Sci*, 69, 118 (Abstr), 1990.
19. **Savage TF, Mirosh LW:** The effects of feeding Medium White turkey hens a breeder diet containing a 1.5% yeast culture. *Poult Sci Tech*, 69, 189 (Abstr), 1990.
20. **NRC:** Nutrient Requirements of Poultry. 9th Revised ed, Natl Acad Press, Washington, 1994.
21. **AOAC:** Official Methods of Analysis of the Association of Official Analytical Chemist. 16th ed., Arlington, VA, 1996.
22. **TSE:** Hayvan Yemleri, Metabolik (Çevrilebilir) Enerji Tayini (Kimyasal Metot). TSE No: 9610, Türk Standartları Enstitüsü, Ankara, 1991.
23. **Düzgüneş O, Kesici T, Gürbüz F:** İstatistik Metotları. Ankara Univ Zir Fak, Yayın No: 861, 1983.
24. **Shermer CL, Maciorowski KG, Bailey CA, Byers FM, Ricke SC:** Caecal metabolites and microbial populations in chickens consuming diets containing a mined humate compound. *J Sci Food Agric*, 77, 479-486, 1998.
25. **Sahin T, Kaya I, Unal Y, Aksu Elmali D:** Dietary supplementation of probiotic and prebiotic combination (combiotics) on performance, carcass quality and blood parameters in growing quails. *J Anim Vet Adv*, 7 (11): 1370-1373, 2008.
26. **Savage TF, Mirosh LW:** Reproductive performance of turkey breeder hens fed a yeast culture, *Proc 25th Annual Pacific NW Animal Nutrition Conf*, 6-8 November 1990.
27. **Kaya CA, Tuncer ŞD:** The Effects of humates on fattening performance, carcass quality and some blood parameters of broilers. *J Anim Vet Adv*, 8 (1): 94-98, 2009.
28. **Yalçın S, Şehu A, Onbaşılar EE, Şahin T:** Broiler rasyonlarına humat ve probiyotik ilavesinin performans üzerine etkileri. *Ankara Univ Vet Fak Derg*, 50, 239-244, 2003.
29. **Avcı M, Denek N, Kaplan O:** Effects of humik asit at different levels on growth performance, carcass yield and some biochemical parameters of quails. *J Anim Vet Adv*, 6 (1): 1-4, 2007.
30. **Küçükersan S, Küçükersan K, Göncüoğlu E, Şahin T:** The effects of dietary humate supplementation on laying hen egg production and egg quality. *Indian Vet J*, 81, 674-678, 2004.
31. **Özçelik H, Yalçın S:** Broiler rasyonlarında L-karnitin ve sodyum humat kullanımı. *Ankara Univ Vet Fak Derg*, 51, 63-69, 2004.
32. **Ceylan N, Çiftçi İ, İlhan Z:** Büyütme faktörü antibiyotiklere alternatif yem katkılarının etlik piliçlerde besi performansı ve bağırsak mikroflorası üzerine etkileri. *Türk J Vet Anim Sci*, 27, 727-733, 2003.
33. **Eren M, Deniz G, Biricik H, Gezen ŞŞ, Türkmen İİ, Yavuz HM:** Broiler yemlerine zink basitrasın, probiyotik ve mannan-oligosakkaritleri katkısının besi performansı üzerine etkileri. *Uludağ Univ Vet Fak Derg*, 3 (18): 73-85, 1999.
34. **Aksu T, Bozkurt AS:** Effect of dietary essential oils and / or humic acids on broiler performance, microbial population of intestinal content and antibody titres in the summer season. *Kafkas Univ Vet Fak Derg*, 15 (2): 185- 190, 2009.