

# The Effect of Orally Administrated $\beta$ -glucan and Dietary Restriction on Faecal Microflora in Rats <sup>[1]</sup>

Tülay ELAL MUŞ <sup>1</sup> Füsün AK SONAT <sup>2</sup> 

<sup>[1]</sup> This study was financially supported by the Uludag University Research Fund (Project No, HDP(V) 2013-45)

<sup>1</sup> Department of Food Hygiene and Technology, Faculty of Veterinary Medicine, University of Uludag, TR-16059 Bursa - TURKEY

<sup>2</sup> Department of Physiology, Faculty of Veterinary Medicine, University of Uludag, TR-16059 Bursa - TURKEY

Article Code: KVFD-2014-11481 Received: 29.04.2014 Accepted: 02.10.2014 Published Online: 13.10.2014

## Abstract

In this study we aimed to evaluate the effect of orally administered  $\beta$ -glucan and/or dietary restriction on *Lactobacillus* spp., coliforms, *Enterobacteriaceae* and enterococci counts in rat faeces. For this aim, rats were divided into three experimental groups: i) first group animals received normal diet for 6 months and administrated orally with  $\beta$ -glucan (20 mg/kg for bodyweigh) over the last 14 day of experiment, ii) second group was dietary restricted animals for 6 months and receiving  $\beta$ -glucan as those of first group animals, iii) last group was the control group rats receiving only *ad libitum* feed. Compared to control group, numeric increase was observed in the number of coliforms, *Enterobacteriaceae* and lactobacillus counts in first group but this was not statistically important. The increase in coliforms and *Enterobacteriaceae* counts was nearly 2 log while this was 1 log for lactobacillus counts. Interestingly, dietary restriction +  $\beta$ -glucan administration had no significant influence on the increase of defined bacterial groups. The results of the present study showed that orally administration of the  $\beta$ -glucan, widely used as prebiotic, has the potential to modify faeces microbiota in rat model.

**Keywords:**  $\beta$ -glucan, Probiotic, Prebiotic, *Lactobacillus*

## Oral Yolla Uygulanan $\beta$ -glukan ve Diyet Kısıtlamasının Ratların Fekal Mikroflorasına Etkisi

### Özet

Bu çalışmanın amacı oral yolla verilen  $\beta$ -glukan ve/veya diyet kısıtlamasının rat feçesindeki *Lactobacillus* spp., koliform, *Enterobacteriaceae* ve enterokok sayıları üzerine etkisini ortaya koymaktır. Bu amaçla ratlar üç deneysel gruba ayrıldı: i) birinci grup 6 ay süresince normal diyetle beslenen ve son 14 gün oral yolla  $\beta$ -glukan (20 mg/kg, canlı ağırlığa) verilen hayvanlar ii) ikinci grup 6 ay boyunca diyet kısıtlaması uygulanan ve ilk gruba aynı şekilde  $\beta$ -glukan verilen hayvanlar iii) yalnızca *ad libitum* beslenen kontrol grubu ratlardan oluşmaktadır. Kontrol grubu ile karşılaştırıldığında, ilk grupta koliform, *Enterobacteriaceae* ve laktobasil sayılarında artış gözlemlendi, fakat bu artış istatistiksel olarak önemli değildi. Koliform ve *Enterobacteriaceae* sayılarında artış yaklaşık 2 log iken, laktobasil sayılarında bu artış 1 log idi. İlginç olarak diyet kısıtlaması ile birlikte  $\beta$ -glukan uygulaması adı geçen bakterilerin artışında önemli bir etki oluşturmadı. Mevcut çalışmanın sonuçları, prebiyotik olarak geniş kullanım alanı bulunan  $\beta$ -glukan'ın oral yolla takviyesinin rat modelinde dışkıdaki mikroorganizmaları modifiye etme potansiyeline sahip olduğunu gösterdi.

**Anahtar sözcükler:**  $\beta$ -glukan, Probiyotik, Prebiyotik, Laktobasil

## INTRODUCTION

Prebiotic is defined as "a selectively fermented ingredient that allows specific changes, both in the composition and/or activity in the gastrointestinal microbiota that confers benefits upon host wellbeing and health" <sup>[1,2]</sup>. Today prebiotics have been widely used to enhance health in the human life <sup>[3]</sup>. Additionally they

benefits on growth and performance of animals <sup>[4]</sup>. Probiotics and prebiotics are mostly consumed for strenghtening of friendly microflora of gut <sup>[5]</sup>.  $\beta$ -glucan, which is a novel prebiotic, has many health benefits. Some of these effects are immunomodulation, increasing the defence system to invading pathogens, anti-tumor activity, lowering the blood cholesterol level and beneficially affect to gut health <sup>[4,6]</sup>. Because of this effects,  $\beta$ -glucan



**İletişim (Correspondence)**



+90 224 2941229



fusunak@uludag.edu.tr

is considered as biological response changers and regard from the pharmaceutical and functional food industries [7]. In addition to these positive properties  $\beta$ -glucan has prebiotic effect by improving the growth of lactobacilli in gut microflora [8] and is one of the most studied prebiotics [9]. Degradation of  $\beta$ -glucan by microbiota resulted with some fermentation products such as short chain fatty acids (SCFA) which are important for the development of the colon microflora. Presence of SCFA reduce the colonic pH and increases the bacterial cultivation [10]. Production of SCFA also prevents growth of potential pathogens [11]. Many researchers reported that  $\beta$ -glucan has positive effects on probiotic bacteria such as lactobacilli and bifidobacteria in gut microflora by animal experiments [12-14]. But there is not enough research about impact of  $\beta$ -glucan to intestinal microflora such as *Lactobacillus*, coliform, *Enterobacteriaceae*, enterococci.

The composition of the gastrointestinal microorganisms is intensively influenced by several factors that include the main gut flora at birth, host genetics, immunological factors, antimicrobial consumption and dietary effects. Change the intakes of the carbohydrates, proteins and fats can significantly modify the composition of the microflora [15].

Dietary restriction (DR) that underfeeding without malnutrition, extends rats life-span in comparison with *ad libitum* feeding [16,17]. DR is the only experimental orientation that has been presented to delay aging, decrease diseases, health risks and the occurrence and proceeding of tumors [16]. DR delays onset of various age-related neurodegenerative diseases such as Alzheimer's disease and insuline resistance. DR is also ameliorate oxidative stress-related impairment in tissues [17]. Although these positive effects there is not enough research about impact of dietary restriction to intestinal microflora.

The aim of this study was to evaluate the in vivo prebiotic potential of  $\beta$ -glucan and the efficiency of long term dietary restriction supported with  $\beta$ -glucan on rat faecal microbiota such as *Lactobacillus*, coliform, *Enterobacteriaceae*, enterococci.

## MATERIAL and METHODS

### Animals and Management

Male, 3-months old Sprague-Dawley rats were used in this study. Rats were purchased from Experimental Animals Breeding and Research Centre of Uludag University. Thirty Sprague-Dawley rats were housed under controlled humidity (50-60%) and temperature (20-23°C) with a 12 h light-dark cycle. Rats had free access to tap water, and fed with a laboratory chow diet. The composition of the diet was as follows: protein 18% (min), lipid 2.5% (min), fiber 4% (max), ash 5.5% (max), nitrogen free extract 57.0% (max), metabolic energy 2650 kcal/kg (min), water 13% (max) plus various amino acids, minerals and vitamins (data

obtained from the supplier). This study was performed after approval of the Local Ethical Committee of Animal Experiments of Uludag University with decision number 2012-14/05.

### Study Design

All rats were divided to 3 groups and each group consisted of 10 rats. All animals were cared during 6 months. First and third (control) groups were given tap water and standard laboratory chow diet with *ad libitum*. Also, first group was orally administered by  $\beta$ -glucan. Second group rats was applied by DR (monday, wednesday and friday mornings) (the food hoppers were placed the following morning as an every-other-day feeding schedule) and tap water *ad libitum* [16,17]. During last 14 days of 6 months housing, daily  $\beta$ -glucan intake in first and second groups were 20 mg/kg dose by intragastric way.

### Microbiological Analyses

Rat faecal samples were collected from cage litters at 1<sup>st</sup>, 7<sup>th</sup>, and 14<sup>th</sup> days under sterile conditions and stored at -20°C for microbiological analysis until the last required sample is collected. Approximately 5 g of each faecal sample were homogenised in 45 ml sterile peptone water (0.1%) solution using a stomacher for at least two minutes. Decimal dilutions were made in sterile pepton water (0.1%) and plated in duplicate on the selective agars. Coliforms were isolated on Violet Red Bile agar (CM0107, Oxoid, UK) with overnight (20-24 h) incubation at 35-37°C. For enterococci the culture was grown on Slanetz and Bartley medium (CM0377, Oxoid, UK) at 35-37°C for 48 h. *Enterobacteriaceae* count was performed by Violet Red Bile Glucose agar (CM0485, Oxoid, UK) and incubated 20-24 h at 35-37°C [18]. *Lactobacillus* spp. was enumerated on de Man Rogosa Sharpe medium (CM0361, Oxoid, UK) and plates were incubated under anaerobic conditions at 35-37°C for 72 h [19].

### Statistical Analyses

Statistical analyses were performed by SPSS 20 programme by Kruskal-Wallis test. When differences among the groups were significant, Mann-Whitney test was used. For interpreting results  $P < 0.05$  significance level was used.

## RESULTS

The results related to the counts of coliforms, *enterobacteriaceae*, *enterococci* and *Lactobacillus* spp. according to groups are summarized in Table 1. Increase in coliform counts was almost 2 logs in the first group, while this increase was slightly lower in other groups with a change of 1 log. But there was no statistically significant difference ( $P > 0.05$ ) among the groups. Only the first group animals exhibited changes in term of *Enterobacteriaceae* counts which were increased 2 log in 2 weeks period. In

**Table 1.** *Lactobacillus* spp., Coliform bacteria, Enterobacteriaceae and Enterococci counts (log<sub>10</sub> cfu/g) according to groups and sampling days**Table 1.** Gruplara ve örnekleme günlerine göre *Lactobacillus* spp., koliform bakteri, Enterobacteriaceae ve enterokok sayıları (log<sub>10</sub> kob/g)

Microorganism	Day 1			Day 7			Day 14		
	First (β-glucan)	Second (DR+ β-glucan)	Third (Control)	First (β-glucan)	Second (DR+ β-glucan)	Third (Control)	First (β-glucan)	Second (DR+ β-glucan)	Third (Control)
	Mean±SEM			Mean±SEM			Mean±SEM		
<i>Lactobacillus</i> spp.	8.67±0.07*	9.02±0.17*	8.92±0.15*	8.53±0.07	9.02±0.16	8.64±0.05	9.04±0.40	9.11±0.10	8.66±0.09
Coliform	4.62±0.13	5.13±0.59	4.27±0.28	5.11±0.27	5.17±0.31	5.01±0.38	6.37±0.37	5.36±0.42	5.01±0.30
Enterobacteriaceae	4.80±0.09	5.27±0.60	4.97±0.14	5.18±0.22	5.20±0.27	5.06±0.42	6.35±0.51	5.44±0.45	4.77±0.39
Enterococci	5.77±0.23	5.81±0.48	5.57±0.42	6.08±0.14	5.83±0.12	5.75±0.07	6.26±0.22	5.34±0.23	6.07±0.38

\* Values sharing the same symbol in the same row are not significantly (P>0.05) different among groups

other two (control and DR + β-glucan) groups, the counts of *Enterobacteriaceae* did not change. Although the higher counts of enterococci in first and third groups were detected, the faecal samples of the second group rats were not assigned any change for enterococci counts after 14 days period. For *Enterobacteriaceae* and enterococci counts, the differences among groups were not statistically significant (P>0.05).

*Lactobacillus* spp. population in first group were similar at 1<sup>st</sup> and 7<sup>th</sup> days. But on 14<sup>th</sup> day the counts of these bacteria were observed a 1 log increase. On the other hand the increase was not statistically significant (P>0.05). On 1<sup>st</sup>, 7<sup>th</sup> and 14<sup>th</sup> days, *Lactobacillus* spp. counts in second and third groups were not seen any difference.

## DISCUSSION

This study was carried out to determine the effect on faecal microorganism counts of the feeding by β-glucan. The addition of β-glucan in the diet of rats did not significantly increased (P>0.05) the numbers of coliform, enterococci, *Enterobacteriaceae*, enterococci and *Lactobacillus* spp. in the faecal samples (1, 7 and 14 days after feeding).

In the present study, coliform counts increased at least one log at first (β-glucan) and third groups. On the other hand, Turunen et al.<sup>[10]</sup> reported that faecal coliform populations were reduced in the human faeces with β-glucan feed on 30<sup>th</sup> days. We found that feed with β-glucan caused a 2 log increase in *Enterobacteriaceae* counts. A similar result was reported by Murphy et al.<sup>[20]</sup> who suggested that oat based β-glucan raised *Enterobacteriaceae* populations 2 log in the porcine gastrointestinal tract. In second (DR+ β-glucan) and third (control) groups, *Enterobacteriaceae* counts didn't change on 14<sup>th</sup> day. Enterococci populations at first and third group rats increased in the end of the feed. In another study investigated of the effect on faecal microflora of β-glucan, Lowry et al.<sup>[21]</sup> reported that β-glucan as feed additive significantly reduced the incidence of *Salmonella enterica* serovar *Enteritidis* organ invasion in immature chickens. In

this study, cause of the increase in the number of coliforms and enterococcus bacteria in the group treated with β-glucan, may be due to lack of the dose of β-glucan or administration time.

In only β-glucan feed rats, *Lactobacillus* spp. populations showed an increasing at 1 log level. There are reports showing that a β-glucan was able to increase *Lactobacillus* spp. in rat faeces<sup>[14,22]</sup>. Again, Reilly et al.<sup>[12]</sup> and Murphy et al.<sup>[20]</sup> assigned that β-glucan raised lactobacillus counts in porcine gastrointestinal tract. Also, a study conducted by Mitsou et al.<sup>[23]</sup> and Kuda et al.<sup>[13]</sup> exhibited that β-glucan significantly increased probiotic bacteria in human and rat faeces, respectively. Dietary restriction supplemented with β-glucan (second group) did not effect *Lactobacillus* spp. populations in the our study. This result could depends on the β-glucans efficacy inhibited by long time dietary restriction treatment. Therefore could not observed any increase in *Lactobacillus* spp. in second group rats. In our knowledge, the results of the present study represent the first data on the effect of dietary restriction + β-glucan on some selected faecal microorganisms in rats. On the other hand, several studies indicated that the addition of different dietary supplements such as sorbitol, heparin or heparosan at the diet increased *Lactobacillus* spp. population in gut microbiota<sup>[9,24]</sup>.

In conclusion, we found that addition of prebiotic matter β-glucan has the potential to change at rat faeces microflora, and to increase, despite not a statistically significant, both probiotic lactobacillus and other microorganisms including coliform, *Enterobacteriaceae* and enterococci populations. In the future we aim to find the exact β-glucan dose which can be used to reduce the colonization of pathogenic bacteria and to increase probiotic bacteria populations as lactobacillus in gastrointestinal tract of animals.

## REFERENCES

1. Gibson GR, Probert HM, Van Loo JAE, Roberfroid MB: Dietary modulation of the human colonic microbiota: updating the concept of prebiotics. *Nutr Res Reviews*, 17, 257-259, 2004.

- 2. Saad N, Delattre C, Urdaci M, Schmitter JM, Bressollier P:** An overview of last advances in probiotic and prebiotic field. *LWT Food Sci Technol*, 50, 1-16, 2013.
- 3. Özpınar H, Aydın İH, Klasing KC, Tekiner İH:** Interaction of manan oligosaccharide with immune system "transport of MOS in to the Lamina Propria". *Kafkas Univ Vet Fak Derg*, 18, 121-128, 2012.
- 4. Yalçın S, Yalçın S, Eser H, Şahin A, Yalçın SS, Güçer Ş:** Effects of dietary yeast cell wall supplementation on performance, on carcass characteristics, antibody production and histopathological changes in broilers. *Kafkas Univ Vet Fak Derg*, 20, 757-764, 2014. DOI: 10.9775/kvfd.2014.11088
- 5. Gülmez M, Güven A:** Probiyotik, prebiyotik ve sinbiyotikler. *Kafkas Univ Vet Fak Derg*, 8, 83-89, 2002.
- 6. Lam KL, Cheung PCK:** Non-digestible long chain beta-glucans as novel prebiotics. *Bioact Carbonhydr Dietary Fibre*, 2, 45-64, 2013.
- 7. Russo p, Lopez P, Capozzi V, de Palencia PF, Duenas MT, Spano G, Fiocco D:** Beta-glucans improve growth, viability and colonization of probiotic microorganisms. *Int J Mol Sci*, 13, 6026-6039, 2012.
- 8. Licht TR, Ebersbach T, Frøkiær H:** Prebiotics for prevention of gut infections. *Trends Food Sci Technol*, 23, 70-82, 2012.
- 9. Sarmiento-Rubiano LA, Zuniga M, Perez-Martinez G, Yebra MJ:** Dietary supplementation with sorbitol results in selective enrichment of lactobacilli in rat intestine. *Res Microbiol*, 158, 694-701, 2007.
- 10. Turunen K, Tsouvelakidou E, Nomikos TZ, Mountzouris KC, Karamanolis D, Triantafyllidis J, Kyriacou A:** Impact of beta-glucan on the fecal microbiota of polypectomised patients: A pilot study. *Anaerobe*, 17, 403-406, 2011.
- 11. Williams BA, Mikkelsen D, Le Pailh L, Gidley MJ:** *In vitro* fermentation kinetics and end-products of cereal arabinoxylans and (1,3;1,4)-b-glucans by porcine faeces. *J Cereal Sci*, 53, 53-58, 2011.
- 12. Reilly P, Sweeney T, O'Shea C, Pierce KM, Figat S, Smith AG, Gahan DA, O'Doherty JV:** The effect of cereal-derived beta-glucans and exogenous enzyme supplementation on intestinal microflora, nutrient digestibility, mineral metabolism and volatile fatty acid concentrations in finisher pigs. *Animal Feed Sci Tech*, 158, 165-176, 2010.
- 13. Kuda T, Enomoto T, Yano T:** Effects of two storage  $\beta$ -1,3-glucans, laminaran from *Eicenia bicyclis* and paramylon from *Euglena gracili*, on cecal environment and plasma lipid levels in rats. *J Funct Foods*, 1, 399-404, 2009.
- 14. Snart J, Bibiloni R, Grayson T, Lay C, Zhang H, Allison GE, Laverdiere JK, Temelli F, Vasanthan T, Bell R, Tannock GW:** Supplementation of the diet with high-viscosity beta-glucan results in enrichment for Lactobacilli in the rat cecum. *Appl Environ Microbiol*, 72 (3): 1925-1931, 2006.
- 15. Scott KP, Gratz SW, Sheridan PO, Flint HJ, Duncan SH:** The influence of diet on the gut microbiota. *Pharmacol Res*, 69, 52-60, 2013.
- 16. Aydın C, Ince E, Koparan S, Cangul IT, Naziroglu M, Ak F:** protective effects of long term dietary restriction on swimming exercise-induced oxidative stres in the liver, heart and kidney of rat. *Cell Biochem Funct*, 25, 129-137, 2007.
- 17. Aydın C, Sonat F, Sahin SK, Cangul IT, Ozkaya G:** Long term dietary restriction ameliorates swimming exercise-induced oxidative stres in brain and lung of middle-aged rat. *Indian J Exp Biol*, 47, 24-31, 2009.
- 18. Roberts D, Greenwood M:** Practical Food Microbiology. 3<sup>rd</sup> ed., 160-161, Blackwell Publishing, Massachusetts, USA, 2003.
- 19. Rubio R, Jofre A, Martin B, Aymerich T, Garriga M:** Characterization of lactic acid bacteria isolated from infant faeces as potential probiotic starter cultures for fermented sausages. *Food Microbiol*, 38, 303-311, 2014.
- 20. Murphy P, Dal Bello F, O'Doherty JV, Arendt EK, Sweeney T, Coffey A:** Effects of cereal  $\beta$ -glucans and enzyme inclusion on the porcine gastrointestinal tract microbiota. *Anaerobe*, 18, 557-565, 2012.
- 21. Lowry VK, Farnell MB, Ferro MJ, Swaggerty CL, Bahl A, Kogut MH:** Purified  $\beta$ -glucan as an abiotic feed additive up-regulates the innate immune response in immature chickens against *Salmonella enterica* serovar *Enteritidis*. *Int J Food Microbiol*, 98, 309-318, 2005.
- 22. Zhoua M, Pub C, Xiaa L, Yua X, Zhua B, Chenga R, Xua L, Zhang J:** salean diet increases short chain fatty acids and enriches beneficial microbiota in the Mouse cecum. *Carbonhydr Poly*, 102, 772-779, 2014.
- 23. Mitsou EK, Panopoulou N, Turunen K, Spiliotis V, Kyriacou A:** Prebiotic potential of barley derived  $\beta$ -glucan at low intake levels: A randomised, double-blinded, placebo-controlled clinical study. *Food Res Int*, 43, 1086-1092, 2010.
- 24. Duan R, Chen X, Wang F, Zhang T, Ling P:** Oral administration of heparin or heparosan increases the Lactobacillus population in gut microbiota of rats. *Carbonhydr Polym*, 94, 100-105, 2013.