

# Effects of Sperm from Different Bulls on Developmental Competence, Blastosist Quality and Cell Number of Bovine Embryos *In Vitro*

Uğur ŞEN <sup>1</sup> 

<sup>1</sup> Ahi Evran Üniversitesi, Ziraat Fakültesi, Tarımsal Biyoteknoloji Bölümü, TR-40100 Aşıkpaşa, Kırşehir - TÜRKİYE

Article Code: KVFD-2014-12486 Received: 21.10.2014 Accepted: 08.02.2015 Published Online: 08.02.2015

## Abstract

The aim of this study was to investigate the effects of sperm from different bulls on the developmental competence, blastosist quality and cell number of bovine embryos *in vitro*. *In vitro* matured bovine oocytes were fertilized with frozen-thawed sperm from five different bulls of Austrian Simmental Fleckvieh and then cultured in Synthetic Oviduct Fluid (SOF) medium at 38.5°C, 5% CO<sub>2</sub>, 5% O<sub>2</sub> and 90% N<sub>2</sub> atmosphere with high humidity for 8 days. In the present study, there was no significant effect of bull variations on cleavage, morula, morula/cleavage, blastocyst and blastocyst/cleavage rates of embryos. Additionally, ICM cell numbers and ICM/total cells ratio of blastocysts obtained from *in vitro* fertilized oocytes with five bull's sperm were similar. However trophectoderm and total cell numbers of blastocysts obtained from *in vitro* fertilized oocytes with bull 5 sperm were lower than other bulls (P<0.05). In the present study, blastocysts quality was affected by bull variations and the percentage of excellent or good quality blastosist (Grade I) were lower (P<0.05) in embryos from fertilized oocytes with bull 5 sperm than those of other bulls, but the percentage of low quality blastosist (Grade III) were higher (P<0.05). There was a positive correlation between quality and total cell numbers of blastocysts in experimental groups. Result of present study showed that developmental competence of embryos *in vitro* were not affected by bulls variations, but may be influence blastocyst quality and cell numbers of blastocyst.

**Keywords:** Bovine, Bull, *in vitro* fertilization, Embryo development, Blastocyst quality

## Farklı Boğa Spermalarının Sığır Embriyolarının *In Vitro* Gelişim Yetkinliği, Blastosist Kalitesi ve Hücre Sayısı Üzerine Etkileri

### Özet

Bu çalışma farklı boğa spermalarının sığır embriyolarının *in vitro* gelişim yetkinliği, blastosist kalitesi ve hücre sayısı üzerine etkilerini belirlemek amacıyla yapılmıştır. *In vitro* olgunlaştırılan sığır oositleri, 5 farklı Avusturyan Simental Fleckvieh boğasına ait dondurulmuş-çözdürülmüş boğa sperması ile fertilize edilmiş ve sonra sentetik ovidukt sıvısı medyumunda yüksek oranda nemlendirilmiş 38,5°C, %5 CO<sub>2</sub>, %5 O<sub>2</sub> ve %90 N<sub>2</sub> içeren atmosferde 8 gün boyunca kültüre alınmıştır. Sunulan çalışmada, embriyoların bölünme, morula, morula/bölünme, blastosist ve blastosist/bölünme oranları üzerine boğa çeşitliliğinin etkisi tespit edilmemiştir. Ayrıca beş farklı boğa sperması ile *in vitro* fertilize edilmiş oositlerden elde edilen blastosistlerin İHK hücre sayısı ve İHK/toplam hücre oranı benzer bulunmuştur. Ancak boğa 5 sperması ile *in vitro* fertilize edilmiş oositlerden elde edilen blastosistlerin trofektoderm ve toplam hücre sayısı diğer boğalardan daha düşük olduğu belirlenmiştir (P<0.05). Sunulan çalışmada blastosist kalitesi boğa çeşitliliği tarafından etkilenmiş olup mükemmel ve iyi kaliteli (Grade I) blastosistlerin yüzdesi boğa 5 sperması ile fertilize edilmiş oositlerden elde edilen embriyolarda diğer boğalarınkilerden daha düşük (P<0,05), fakat düşük kalitedeki (Grade III) blastosistlerin oranı ise daha yüksek olduğu belirlenmiştir (P<0.05). Bütün deneme gruplarında blastosist kalitesi ile toplam hücre sayısı arasında pozitif bir korelasyon saptanmıştır (P<0.05). Mevcut çalışmanın sonuçları boğa çeşitliliğinin *in vitro* embriyo gelişim yetkinliğini etkilemediği, ancak blastosist kalitesinin ve blastosistlerin hücre sayılarını etkileyebileceğini göstermiştir.

**Anahtar sözcükler:** Sığır, Boğa, *in vitro* fertilizasyon, Embriyo gelişimi, Blastosist kalitesi

## INTRODUCTION

*In vitro* embryo production is one of reproductive biotechnology applications and increases the speed of genetic improvement in farm animals <sup>[1]</sup>. Moreover this

reproductive biotechnology has enabled large amounts embryo production of superior breeds in various domestic animals and the production of large numbers of embryos for scientific research purposes from slaughtered and/or live animals <sup>[2,3]</sup>. Production of embryos *in vitro* also



İletişim (Correspondence)



+90 386 2804832



ugur.sen@ahievran.edu.tr

has potential for revolutionizing in cattle breeding and husbandry around the globe. Although several decades of research have gone into *in vitro* embryo production, the process involve *in vitro* embryo production is yet to be standardized and ratio of *in vitro* fertilized oocytes reaching the blastocyst stage is still low, approximately 30-40%<sup>[1]</sup>. Probably, the reason of this situation is *in vitro* embryo production practices cannot mimic *in vivo* conditions, resulting in alterations of development, morphology and gene expression in embryos<sup>[4]</sup>.

Many factors can impact efficiency of *in vitro* embryo production<sup>[5,6]</sup> and one of these factors is use of sperm from different bulls<sup>[7]</sup>. Fertility ability of sperm is one of the main factors for success of *in vitro* fertilization, and previous studies reported that there are differences in *in vitro* embryo production rates among bulls<sup>[8-10]</sup>. When success rate of bulls in in-vitro fertilization is determined, *in vitro* embryo production may be standardized and decrease the differences in rate of embryos reach to blastocyst stage among bulls. Additionally genetic potential of sperm from different bull should be considered due to genetic information transmitted by sperm to the embryo. Moreover, embryo quality and success of *in vitro* embryo production applications can be influenced by genetic heritage of sperm<sup>[11]</sup>. The aim of the study was therefore to investigate the effects of sperm from different bulls on the developmental competence, blastosist quality and cell number of bovine embryos *in vitro*

## MATERIAL and METHODS

All chemicals and media used in this study were purchased from Sigma-Aldrich (St. Louis, MO, USA), except where otherwise indicated.

### Collection and In Vitro Maturation of Oocytes

Cumulus-oocyte complexes (COCs) were obtained by aspirating antral follicles (2 to 5 mm in diameter) of bovine ovaries obtained from a local slaughterhouse. The COCs were collected in 3-4 ml Hepes-buffered Medium-199 containing Earle's salts and supplemented with 1% v/v antibiotic-antimycotic solution (10.000 IU penicillin, 10 mg streptomycin and 25 µg amphotericin B per ml). COCs were assessed morphologically and only oocytes with compact, intact cumulus cells around and homogeneous cytoplasm were selected for maturation. COCs were washed three times in Hepes-buffered Medium 199, and then twice in maturation medium. Maturation medium were prepared as reported by Cevik et al.<sup>[12]</sup>. Maturation medium was sodium bicarbonate-buffered Medium-199 containing Earle's salts and L-glutamine supplemented with sodium pyruvate (5.5 µg/ml), antibiotic-antimycotic solution (1% v/v), heat-inactivated FCS (10% v/v), LH (5.0 µg/ml), FSH (0.5 µg/ml) and EGF (10 ng/ml). The COCs were placed in 500 µl of maturation medium

(approximately 30-35 COCs per well) covered with 300 µl mineral oil in four-well dishes (Nunc, Roskilde, Denmark) and matured for 22 h in a humidified atmosphere of 5% CO<sub>2</sub> in air at 38.5°C.

### Sperm Preparation and In Vitro Fertilization

Cumulus expansion degree of COCs was assessed under a stereomicroscope at the end of the *in vitro* maturation period. COCs with full cumulus expansion considered as matured oocytes and immature COCs were discarded from experiment. The matured COCs were washed twice in Hepes-buffered Medium-199 and twice in fertilization medium. The fertilization medium was glucose-free modified TALP supplemented with 25 mM sodium bicarbonate, 22 mM Na-lactate, 1 mM Na-pyruvate, 6 mg/ml fatty acid-free BSA, 10 mg/ml heparin-sodium salt and 0.5 µl/ml antibiotic-antimycotic solution (pH 7.4 and 280-300 mOsm/kg). The matured COCs were then randomly separated and transferred into 45 µl fertilization drops (approximately 15 COCs per drop) covered with mineral oil. Frozen thawed semen from five different bulls (named; Bull 1, Bull 2, Bull 3, Bull 4 and Bull 5) of Austrian Simmental Fleckvieh was used for the *in vitro* fertilization in this study (Genovet L.T.D., S.T.I, Samsun, Turkey). Before *in vitro* fertilization frozen-thawed semen from five different bulls were separated by Percoll density gradient technique<sup>[13]</sup>. The percentage of sperm motility was visually evaluated using a phase-contrast microscope at a magnification of 400× (at least 80% progressively motile). The sperm concentration was determined by hemocytometer using a Thoma counting chamber. Sperm was then diluted to  $2 \times 10^6$  sperm/ml with fertilization medium. COCs were fertilized randomly with 5 µl diluted sperm (approximately  $13 \times 10^3$  sperm for per oocytes) from five bulls for 22 h in a humidified atmosphere of 5% CO<sub>2</sub> in air at 38.5°C.

### In Vitro Culture

After *in vitro* fertilization, the putative zygotes were washed three times in Hepes-buffered Medium 199 and vortexing for approximately 5 min, to remove the cumulus cells. The naked putative zygotes were then washed twice in Hepes-buffered Medium 199 and twice in synthetic oviduct fluid (SOF) embryo culture media. The SOF embryo culture media was supplemented with 20 µl/ml pyruvate, 8 mg/ml fatty acid-free BSA, 20 µl/ml MEM non-essential amino acids solution 100×, 10 µl/ml BME Amino Acids Solution 50× and 0.5 µl/ml antibiotic-antimycotic solution on the day of use. The naked putative zygotes were placed in 50 µl drops (approximately 15 zygotes per drop) of SOFaa media under mineral oil and cultured in a humidified atmosphere of 5% CO<sub>2</sub>, 5% O<sub>2</sub> and 90% N<sub>2</sub> in air at 38.5°C. *In vitro* fertilization was considered as 0 day. On day 3 of development the proportion of zygotes cleaved was recorded. Morula and blastocyst development the proportion of zygotes were evaluated


on days 5 and 8, respectively. Experiments were repeated 5 times for each group.

### Determination of Blastocyst Quality

The quality grading of the recovered blastocysts was done according to the morphological criteria of quality determined by International Embryo Transfer Society [14]. Blastocysts were classified as Grade I; excellent or good quality blastocysts with blastocoele filling the entire blastocyst, oval shaped and compact inner cell mass (ICM) and multicellular cohesive trophectoderm (TE), Grade II; moderate quality blastocysts with normal ICM, but non-optimal (fragmented or necrotic) TE, and Grade III poor quality blastocysts with very few cells or without ICM and with large vacuole instead of blastocoele.

### Differential Staining of ICM and TE Cells

After determination of blastocyst quality, blastocysts were differential stained as described by Van Soom et al. [15] with some modifications. Briefly, the blastocysts were washed in PBS supplemented with 0.1 mg/ml polyvinylalcohol and then incubated with picrylsulphonic acid diluted to with  $\text{Ca}^{2+}$ -free PBS (10 mM) for 5 min in the refrigerator (4°C). The blastocysts were then washed in PBS supplemented with 0.1 mg/ml polyvinylalcohol and incubated for 30 min at 38.5°C in anti-dinitrophenyl antibody diluted to 30% (v/v) with  $\text{Ca}^{2+}$ -free PBS. The blastocysts were repeat washed in PBS supplemented with 0.1 mg/ml polyvinylalcohol and transferred into guinea pig complement (55852, ICN biochemicals, Irvine, CA, USA) diluted to 20% (v/v) with  $\text{Ca}^{2+}$ -free PBS



**Fig 1.** The picture of a blastocyst after differential staining (400× magnification)

**Şekil 1.** Diferansiyel boyama sonrası bir blastosist resmi (400× büyütme)

supplemented with 50 µg/ml propidium iodide and 12.5 µg/ml bisbenzamide for 30 min at 38.5°C. Finally, blastocysts were fixed in 2% paraformaldehyde for 1-2 min at room temperature. Before mounting on slides with a 10 µl drop of 0.2 M 1.4 diazabicyclo-octanes in glycerol (50%, v/v) in  $\text{Ca}^{2+}$ -free PBS as an anti-fading solution. Blastocysts were examined under a fluorescence microscope (Nikon Invert Microscope Eclipse Ti-FL, 340-380 nm excitation and 430 nm suppression). ICM nuclei labelled with bisbenzamide appeared blue and TE nuclei labelled with both bisbenzamide and propidium iodide appeared pink to red (Fig. 1). The numbers of ICM and TE nuclei were counted directly under the fluorescence microscope using a 345 nm ultraviolet light filter.

### Statistical Analysis

Data were analyzed by one-way ANOVA after appropriate transformation where necessary (percentage of cleaved zygotes, morula and blastocyst yields, arcsine-transformation; cell numbers of blastocyst,  $\log_{10}$  transformation). Data involved in blastocyst quality were analyzed by chi-square ( $\chi^2$ ) test. Relationships between quality and total cell numbers of blastocysts were determined with a Pearson correlation analysis at the 95% confidence interval. Significant differences between means were tested using Duncan's test with SPSS 20.0. The results are presented as untransformed mean  $\pm$  SE values, and statistical significance was determined at the level of 0.05.

## RESULTS

In the present study, total number of 2389 bovine cumulus-oocyte complexes (COCs) were used and matured in standard maturation medium. Approximately 92% of COCs matured following *in vitro* maturation. *In vitro* matured oocytes were subjected to the *in vitro* fertilization procedure with sperm from various bulls and total numbers of 2198 bovine embryos were cultured in SOF media.

Developmental characteristics of *in vitro* produced bovine embryos obtained from *in vitro* fertilized oocytes with different bulls' sperm are presented in the Table 1. There were no significant differences between bulls in terms of cleavage, morula and blastocyst formation rates. Similarly, there was no significant effect of bull variations on morula/cleavage and blastocyst/cleavage rates of embryos.

Inner cell mass, trophectoderm and total cell numbers of *in vitro* produced bovine blastocysts obtained from *in vitro* fertilized oocytes with different bulls' sperm are presented in the Table 2. ICM cell numbers of blastocysts obtained from *in vitro* fertilized oocytes with five bull's sperm were similar, but trophectoderm and total cell

**Table 1.** Developmental characteristics of *in vitro* produced bovine embryos obtained from *in vitro* fertilized oocytes with different bulls' sperm. Results are presented as untransformed mean values ( $\pm$  SEM) per groups for 5 replicates

**Tablo 1.** Farklı boğaların sperması ile *in vitro* fertilize edilen oositlerden elde edilen *in vitro* üretilmiş siğir embriyolarının gelişim özellikleri. Sonuçlar 5 tekrarlanmış grup başına dönüştürülmemiş ortalama değerler ( $\pm$  SEM) olarak sunulmuştur

Bulls	Oocytes Fertilized (n)	Developmental Competence of Bovine Embryos (%)				
		Cleavage	Morula	Morula/Cleavage	Blastocyst	Blastocyst/Cleavage
1	420	73.8 $\pm$ 4.03	34.2 $\pm$ 4.23	46.3 $\pm$ 4.03	25.37 $\pm$ 3.79	34.02 $\pm$ 4.71
2	498	69.5 $\pm$ 2.08	44.6 $\pm$ 2.84	64.7 $\pm$ 5.54	29.23 $\pm$ 1.73	42.34 $\pm$ 3.03
3	476	73.9 $\pm$ 1.69	39.3 $\pm$ 4.18	52.5 $\pm$ 5.23	29.26 $\pm$ 4.63	39.48 $\pm$ 6.02
4	424	74.8 $\pm$ 4.11	34.1 $\pm$ 3.47	45.4 $\pm$ 3.17	26.76 $\pm$ 3.95	34.93 $\pm$ 3.86
5	380	73.1 $\pm$ 2.86	37.4 $\pm$ 1.63	51.8 $\pm$ 3.87	25.38 $\pm$ 0.91	34.92 $\pm$ 1.62

**Table 2.** Inner cell mass, trophectoderm and total cell numbers of *in vitro* produced bovine blastocysts obtained from *in vitro* fertilized oocytes with different bulls' sperm. Results are presented as mean values ( $\pm$  SEM) per groups for 5 replicates

**Tablo 2.** Farklı boğaların sperması ile *in vitro* fertilize edilen oositlerden elde edilen *in vitro* üretilmiş siğir blastosistlerinin iç hücre kitlesi, trofektoderm ve toplam hücre sayıları. Sonuçlar 5 tekrarlanmış grup başına ortalama değerler ( $\pm$  SEM) olarak sunulmuştur

Bulls	Cell Numbers		
	Inner Cell Mass	Trophectoderm	Total
1	37.0 $\pm$ 1.70	67.4 $\pm$ 1.91 <sup>a</sup>	104.4 $\pm$ 2.54 <sup>a</sup>
2	34.7 $\pm$ 1.15	62.2 $\pm$ 1.71 <sup>a</sup>	96.9 $\pm$ 2.26 <sup>a</sup>
3	34.4 $\pm$ 1.18	63.7 $\pm$ 2.48 <sup>a</sup>	98.1 $\pm$ 3.06 <sup>a</sup>
4	35.5 $\pm$ 0.94	62.5 $\pm$ 1.46 <sup>a</sup>	98.0 $\pm$ 2.03 <sup>a</sup>
5	34.8 $\pm$ 1.53	51.0 $\pm$ 2.40 <sup>b</sup>	85.8 $\pm$ 2.74 <sup>b</sup>


<sup>a,b</sup> Means in column with different superscripts are significantly different at  $P < 0.05$

**Table 3.** The percentages of blastocyst quality of *in vitro* produced bovine blastocyst obtained from *in vitro* fertilized oocytes with different bulls' sperm. Results are presented as mean values per ( $\pm$  SEM) groups for 5 replicates

**Tablo 3.** Farklı boğaların sperması ile *in vitro* fertilize edilen oositlerden elde edilen *in vitro* üretilmiş siğir blastosistlerinin blastosist kalite oranları. Sonuçlar 5 tekrarlanmış grup başına ortalama değerler ( $\pm$  SEM) olarak sunulmuştur

Bulls	Blastocyst Quality		
	Grade I	Grade II	Grade III
1	36.5 <sup>a</sup>	60.4 <sup>b</sup>	3.1 <sup>c</sup>
2	24.4 <sup>b</sup>	73.5 <sup>a</sup>	2.1 <sup>c</sup>
3	21.7 <sup>b</sup>	72.4 <sup>a</sup>	6.9 <sup>b</sup>
4	26.9 <sup>b</sup>	71.4 <sup>a</sup>	1.7 <sup>c</sup>
5	12.5 <sup>c</sup>	78.3 <sup>a</sup>	9.2 <sup>a</sup>

<sup>a,b</sup> Means in column with different superscripts are significantly different at  $P < 0.05$



**Fig 2.** ICM/total cells number ratio of *in vitro* produced bovine blastocyst obtained from *in vitro* fertilized oocytes with different bulls' sperm

**Şekil 2.** Farklı boğaların sperması ile *in vitro* fertilize edilen oositlerden elde edilen *in vitro* üretilmiş siğir blastosistlerinin İHK/toplam hücre sayısı oranları

numbers of blastocysts obtained from *in vitro* fertilized oocytes with bull 1 sperm were lower than those of other bulls ( $P < 0.05$ ). ICM/total cells number ratio of *in vitro* produced bovine blastocyst obtained from *in vitro* fertilized oocytes with different bulls' sperm are presented in the Fig. 2. There were no significant differences between bulls in terms of ICM/total cells ratio of blastocysts.

**Table 4.** Pearson correlation coefficients (95% confidence intervals) between quality and total cell numbers of *in vitro* produced bovine blastocyst obtained from *in vitro* fertilized oocytes with different bulls' sperm

**Tablo 4.** Farklı boğaların sperması ile *in vitro* fertilize edilen oositlerden elde edilen *in vitro* üretilmiş siğir blastosistlerinin kalitesi ve toplam hücre sayısı arasındaki Pearson korelasyon katsayısı

Bulls	Correlation Coefficients
1	.521*
2	.493*
3	.624*
4	.454*
5	.620*

\*  $P < 0.05$

The percentages of blastocyst quality of *in vitro* produced bovine blastocyst obtained from *in vitro* fertilized oocytes with different bulls' sperm are presented in the Table 3. Blastocyst quality was dependent on bull variations. The percentage of excellent or good quality blastocyst (Grade I) were higher in embryos from fertilized oocytes with sperm of bull 1 than those of other bulls ( $\chi^2 = 0.39$ ;  $P < 0.05$ ), but moderate quality blastocyst

(Grade II) were lower in same bull than those of other bulls ( $\chi^2 = 0.37$ ;  $P < 0.05$ ). The percentage of poor quality blastocyst (Grade III) were higher in embryos from fertilized oocytes with sperm of bull 3 and 5 than those of bull 1, 2 and 4 ( $\chi^2 = 0.32$ ;  $P < 0.05$ ).

Pearson correlation coefficients (95% confidence intervals) between quality and total cell numbers of *in vitro* produced bovine blastocyst obtained from *in vitro* fertilized oocytes with different bulls' sperm are presented in the Table 4. There were positive correlations between quality and total cell numbers of blastocysts from *in vitro* produced bovine embryos obtained from *in vitro* fertilized oocytes with five bulls sperm ( $P < 0.05$ ).

## DISCUSSION

The present study demonstrates that the use of semen from different bull for *in-vitro* fertilization did not affect developmental competence of embryos, but bull variations influenced trophoctoderm and total cell numbers and quality of blastocyst derived *in vitro*. Additionally, excellent or good quality blastocysts had more total cell number than moderate or low quality blastocyst in the experimental groups.

The use of spermatozoa from different bulls during *in vitro* fertilization results in variable fertility rates<sup>[16]</sup>. Saeki et al.<sup>[17]</sup>, Marquant-Le Guienne et al.<sup>[18]</sup> and Thara and Nair<sup>[19]</sup> showed that spermatozoa from different bulls have different rates of fertilization on *in vitro* matured bovine oocytes. Similarly, Totey et al.<sup>[20]</sup> and Jamil et al.<sup>[21]</sup> reported that fertilization rate was significantly different in buffalo oocytes inseminated with sperm from different buffalo bulls. Unfortunately, fertilization rates of *in vitro* fertilized oocytes with sperm from various bulls were not investigated in the present study, but Ciray et al.<sup>[22]</sup> indicated that early cleavage is a biological indicator of fertilization. Thus, it may be mentioned that the fertilization rates were not affected by sperm from various bulls in the present study.

Shamsuddin and Larsson<sup>[23]</sup>, Zhang et al.<sup>[9]</sup> and Sudano et al.<sup>[16]</sup> reported that the use of sperm from various bulls or bulls with different fertility levels during *in vitro* fertilization leads to different cleavage rates of bovine embryos. Additionally, Al Naib et al.<sup>[10]</sup> found that *in vitro* fertilized oocytes with sperm from high fertility Holstein Friesian bulls had a higher rate of early cleavage compared with oocytes fertilized by sperm from low fertility Holstein Friesian bulls. Contrary to these studies, in the present study the cleavage rates were not affected by bull variations and average cleavage rate were 73% in the experimental groups. These finding is in agreement with the arguments of Schneider et al.<sup>[8]</sup> who reported that mean cleavage rates were similar in oocytes inseminated with semen from different bulls with different fertility

levels. Also Galli and Lazzari<sup>[24]</sup> have compared cleavage rates of bulls and they have not seen a difference in the rates of cleavage between bulls. The differences in effects of bull variations on cleavage or fertilization rates with the above-mentioned studies may possibly be due to differences in age and breeds of bull, sufficient capabilities in *in-vitro* fertilization and metabolic characteristics of sperm cells.

Generally blastocyst yield is approximately 25-30% in *in-vitro* embryo production depending on embryo culture media and protocol<sup>[25]</sup>. In the present study, blastocyst development rates were approximately 27% in the experimental groups. Additionally developmental competence of embryos, obtained from *in vitro* fertilized oocytes with five bull's sperm, until the morula and blastocyst stage were similar in the present study. Moreover morula/cleavage and blastocyst/cleavage rates were similar among bulls. These results agree with the work of Shamsuddin and Larsson<sup>[23]</sup> who have demonstrated that the use of different bulls during *in vitro* fertilization have similar effect on embryo development until the morula and blastocyst stage. Similarly Sudano et al.<sup>[16]</sup> reported that although the cleavage rates observed between different bulls changed, differences were not observed among bulls in terms of blastocyst formation rates. On the contrary Akyol et al.<sup>[7]</sup> show that blastocyst development rate had a wide range among bulls. These differences may be explained with the variation at phenotypic traits and breed type of bulls.

Morphological observations are most widely used as indicator of blastocyst quality<sup>[26]</sup>. However the cell number of the blastocyst is a valid indicator of the viability of preimplantation embryos, while morphological criteria alone are poor indicators<sup>[27,28]</sup>. Additionally, Jiang et al.<sup>[29]</sup> reported that the cell number of *in vitro* derived blastocyst decreased with decreasing quality of the blastocyst. The analysis of distribution of ICM and trophoctoderm cells by differential staining has been used as a technique to evaluate embryo quality in several species. Additionally ICM cells/total cell numbers ratio is an indicator of quality of blastocysts. In bovine embryos, poor quality is associated with blastocysts with low numbers of ICM or total cells<sup>[25,27,30]</sup>. In the present study blastocysts obtained from *in vitro* fertilized oocytes with five bulls's sperm had same cell numbers in ICM and the ICM/total cells ratios. However, trophoctoderm and total cell numbers of blastocysts were different between bulls. Moreover quality of blastocysts obtained from *in vitro* fertilized oocytes with five bull's sperm quality of blastocysts varied between bulls. Generally, excellent or good quality blastocysts had more total cell number than moderate or low quality blastocyst in all bulls. Additionally the positive correlations observed between blastocysts quality and total cell number of blastocysts in all bulls. This result supports the view that total cell number influence blastocysts quality

and better quality blastocysts have more cell number than poor quality blastocysts.

The results of this study indicate that the use of sperm from various bulls in *in-vitro* fertilization does not have any effect on embryonic development. However, use of different bulls can influence the quality and cell numbers of the embryos reached the blastocyst stage in similar culture conditions. These results show that effects of bull have to be determined with preliminary studies in *in-vitro* researches and successful bulls should be selected for *in vitro* fertilization. Thus, the effects of bull on the success of *in vitro* embryo production may be eliminated. Moreover, determining the success rate after transfer of embryos obtained from *in vitro* fertilized oocytes with different bulls' sperm will help to identify the bull effect on *in vitro* embryo production more clearly.

## REFERENCES

1. **Camargo LSA, Viana JHM, Sa WF, Ferreira AM, Ramos AA, Vale Filho VR:** Factors influencing *in vitro* embryo production. *Anim Reprod*, 3 (1): 19-28, 2006.
2. **Rajesh N, Shankar MB, Deecaraman M:** Effect of vitamin A supplementation at different gaseous environments on *in vitro* development of pre-implantation sheep embryos to the blastocyst stage. *Animal*, 4, 1884-1890, 2010. DOI: 10.1017/S1751731110001187
3. **Cevik M, Kocyigit A, Sen U, Kuran M:** Can commercial human embryo culture media be used in bovine embryo culture? *Kafkas Univ Vet Fak Derg*, 20, 149-153, 2014. DOI: 10.9775/kvfd.2013.9676
4. **Loneragan P, Fair T, Corcoran D, Evans AC:** Effect of culture environment on gene expression and developmental characteristics in IVF-derived embryos. *Theriogenology*, 65, 137-152, 2006. DOI: 10.1016/j.theriogenology.2005.09.028
5. **Bucak NM, Satilmis M, Kizil HS, Karasahin T, Akyol N:** Sığır embriyolarının *in vitro* gelişiminde kültür medyumlarına katılan antioksidanların etkisi. *Kafkas Univ Vet Fak Derg*, 16, 69-74, 2010. DOI: 10.9775/kvfd.2009.413
6. **Kizil HS, Akyol N, Karasahin T, Satilmis M:** Etilen glikol ile direkt transfer metoduna göre dondurulan *in vivo* sığır embriyolarının transferi. *Kafkas Univ Vet Fak Derg*, 17, 721-724, 2011. DOI: 10.9775/kvfd.2011.4176
7. **Akyol N, Kızıl SH, Satılmış M, Karasahin T:** Investigation of bull effect on *in vitro* embryo production. *Kafkas Univ Vet Fak Derg*, 20, 561-564, 2014. DOI: 10.9775/kvfd.2013.10582
8. **Schneider CS, Ellington JE, Wright Jr RW:** Relationship between bull field fertility and *in vitro* embryo production using sperm preparation methods with and without somatic cell co-culture. *Theriogenology*, 51, 1085-1098, 1999. DOI: 10.1016/S0093-691X(99)80013-2
9. **Zhang M, Lu KH, Seidel GE:** Development of bovine embryos after *in vitro* fertilization of oocytes with flow cytometrically sorted, stained and unsorted sperm from different bulls. *Theriogenology*, 60, 1657-1663, 2003. DOI: 10.1016/S0093-691X(03)00177-8
10. **Al Naib A, Hanrahan JP, Loneragan P, Fair S:** *In vitro* assessment of sperm from bulls of high and low field fertility. *Theriogenology*, 76, 161-167, 2011. DOI: 10.1016/j.theriogenology.2010.10.038
11. **Leibfried-Rutledge ML:** Factors determining competence of *in vitro* produced cattle embryos. *Theriogenology*, 51, 473-485, 1999. DOI: 10.1016/S0093-691X(98)00241-6
12. **Cevik M, Sen U, Kocyigit A, Soydan E, Kuran M:** Effects of serum, gonadotropins, epidermal growth factor and estradiol 17-Beta on cumulus expansion and nuclear maturation of bovine oocytes. *Kafkas Univ Vet Fak Derg*, 17, 1009-1014, 2011. DOI:10.9775/kvfd.2011.4963
13. **Parrish JJ, Krogenaes A, Susko-Parrish JL:** Effect of bovine sperm separation by either swim-up or Percoll method on success of *in vitro* fertilization and early embryonic development. *Theriogenology*, 44, 859-869, 1995. DOI: 10.1016/0093-691X(95)00271-9
14. **Wright JM:** Photographic illustrations of embryo developmental stage and quality codes. In, Stringfellow DA, Seidel SM (Eds): Manual of the International Embryo Transfer Society (IETS), Illinois, USA, 167-170, 1998.
15. **Van Soom A, Boerjan M, Ysebaert MT, Kruijff AD:** Cell allocation to the inner cell mass and the trophectoderm in bovine embryos cultured in two different media. *Mol Reprod Dev*, 45, 171-182, 1996. DOI: 10.1002/(SICI)1098-2795(199610)45:2<171::AID-MRD10>3.0.CO;2-4
16. **Sudano MJ, Crespilho AM, Fernandes CB, Junior AM, Papa FO, Rodrigues J, Machado R, Landim-Alvarenga Fda C:** Use of bayesian inference to correlate *in vitro* embryo production and *in vivo* fertility in zebu bulls. *Vet Med Int*, 14, 1-6, 2011. DOI: 10.4061/2011/436381
17. **Saeki K, Nagao Y, Hoshi M, Nagai M:** Effects of heparin, sperm concentration and bull variation on *in vitro* fertilization of bovine oocytes in a protein-free medium. *Theriogenology*, 43, 751-759, 1995. DOI: 10.1016/0093-691X(95)00017-3
18. **Marquant-Le Guenne B, Humblot P, Thibier M, Thibault C:** Evaluation of bull semen fertility by homologous *in vitro* fertilization tests. *Reprod Nutr Dev*, 30, 259-266, 1990. DOI: 10.1051/rnd:19900212
19. **Thara KM, Nair SPS:** Sire effect on *in vitro* fertilizability of matured cattle oocytes. *Indian J Biotechnol*, 6, 421-422, 2007.
20. **Totey SM, Pawshe CH, Singh GP:** Effects of bull and heparin and sperm concentrations on *in vitro* fertilization of buffalo (*Bubalus bubalis*) oocytes matured *in vitro*. *Theriogenology*, 39, 887-898, 1993.
21. **Jamil H, Samad HA, Qureshi ZI, Rehman N, Lodhi LA:** Effect of bull and sperm preparation method on *in vitro* fertilization of buffalo oocytes. *Pak Vet J*, 27, 29-34, 2007.
22. **Ciray HN, Karagenç L, Ulug U, Bener F, Bahceci M:** Use of both early cleavage and day 2 mononucleation to predict embryos with high implantation potential in intracytoplasmic sperm injection cycles. *Fertil Steril*, 84, 1411-1416, 2005. DOI: 10.1016/j.fertnstert.2005.05.021
23. **Shamsuddin M, Larsson B:** *In vitro* development of bovine embryos after fertilization using semen from different donors. *Reprod Domes Anim*, 28, 77-84, 1993. DOI: 10.1111/j.1439-0531.1993.tb00727.x
24. **Galli C, Lazzari G:** Practical aspects of IVM/IVF in cattle. *Anim Reprod Sci*, 42, 371-379, 1996. DOI: 10.1016/0378-4320(96)01530-8
25. **Gordon I:** Laboratory Production of Cattle Embryos. 2<sup>nd</sup> ed., CABI Publishing, Wallingford, 2003.
26. **Mori M, Otoi T, Suzuki T:** Correlation between the cell number and diameter in bovine embryos produced *in vitro*. *Reprod Domest Anim*, 37, 181-18, 2002. DOI: 10.1046/j.1439-0531.2002.00354.x
27. **Van Soom A, Ysebaert MT, Kruijff AD:** Relationship between timing of development, morula morphology, and cell allocation to inner cell mass and trophectoderm in *in vitro*-produced bovine embryos. *Mol Reprod Dev*, 47, 47-56, 1997.
28. **Stojkovic M, Büttner M, Zakhartchenko V, Brem G, Wolf E:** A reliable procedure for differential staining of *in vitro* produced bovine blastocysts: comparison of tissue culture medium 199 and Menezes's B2 medium. *Anim Reprod Sci*, 50, 1-9, 1998.
29. **Jiang HS, Wang L, Lu KH, Gordon I, Polge C:** Examination of cell numbers of blastocysts derived from IVM, IVF and IVC of bovine follicular oocytes. *Theriogenology* 37, 229, 1992.
30. **Kuran M, Robinson JJ, Brown DS, McEvoy TG:** Development, amino acid utilization and cell allocation in bovine embryos after *in vitro* production in contrasting culture systems. *Reproduction*, 124, 155-165, 2002.