

Analysis of FecB, BMP15 and CAST Gene Mutations in Sakiz Sheep ^[1]

Deniz DİNÇEL ¹ Sena ARDIÇLI ¹ Bahadır SOYUDAL ¹ Mehlika ER ¹
Fazlı ALPAY ² Hale ŞAMLI ¹ Faruk BALCI ¹ ✉

^[1] This study was supported by Uludag University Scientific Research Projects Unit (Project Number: V-2004/4) and the State Planning Organization (Project Number: DPT-2005/2)

¹ Uludag University, Faculty of Veterinary Medicine, Department of Genetics, TR-16059 Nilufer, Bursa - TURKEY

² Uludag University, Faculty of Veterinary Medicine, Department of Zootechnics, TR-16059 Nilufer, Bursa - TURKEY

KVFD-2014-12680 Received: 24.11.2014 Accepted: 13.03.2015 Published Online: 19.03.2015

Abstract

Fertility traits, such as the ovulation rate and the number of offspring at birth, are genetically regulated by fecundity genes. This study was performed to identify DNA polymorphisms in Booroola (*FecB*), Galway (*FecX^G*), Inverdale (*FecX^I*) and Calpastatin (*CAST*) genes in Sakiz sheep. A total of 71 ewes were genotyped for gene or allelic polymorphisms in the genes listed above using the PCR-RFLP method. The results obtained from this study indicated that all of the Sakiz ewes sampled were non-carriers for *FecB*, *FecX^G* or *FecX^I* mutations. However, genotypic frequencies in the *CAST* gene were 0.59, 0.36 and 0.05 for AA, AB and BB, respectively. A significant deviation from Hardy-Weinberg equilibrium for the *CAST* gene was not observed in the investigated breed ($P>0.05$). As a result, more extensive screening is required as tests for newly discovered mutations are developed. Additionally, this study is the first to report a genetic polymorphism in the *CAST* gene in Sakiz sheep.

Keywords: Sakiz sheep, Fertility, *FecB*, *BMP15*, *CAST*

Sakiz Irkı Koyunlarda FecB, BMP15 ve CAST Genlerindeki Mutasyonların Analizi

Özet

Ovulasyon oranı ve bir batındaki yavru sayısı gibi dölerim özellikleri fekondite genleri tarafından regüle edilmektedir. Bu çalışma Sakiz ırkı koyunlarda, Booroola (*FecB*), Galway (*FecX^G*), Inverdale (*FecX^I*) ve Calpastatin (*CAST*) genlerindeki DNA polimorfizmlerini belirlemek amacıyla yapılmıştır. Araştırılan gen veya aleller açısından toplam 71 koyun PCR-RFLP metoduyla genotiplendirilmiştir. Çalışmadan çıkan sonuçlar, Sakiz koyunların *FecB*, *FecX^G* ya da *FecX^I*'ye ait mutasyonları taşımadığını göstermiştir. Ancak *CAST* geni açısından AA, AB ve BB genotiplerinin sırasıyla 0.59, 0.36 ve 0.05 frekans gösterdiği tespit edilmiştir. *CAST* geni açısından çalışılan sürünün Hardy-Weinberg dengesine uyduğu görülmüştür ($P>0.05$). Sonuç olarak yeni keşfedilen mutasyonların geliştirilmesi gibi daha yoğun tarama çalışmaları gerekmektedir. Ek olarak Sakiz koyunlarında *CAST* genindeki polimorfizmin varlığı ilk defa bu çalışmada bildirilmektedir.

Anahtar sözcükler: Sakiz Koyunu, Fertilite, *FecB*, *BMP15*, *CAST*

INTRODUCTION

The incorporation of a major gene for prolificacy into a flock using marker-assisted selection (MAS) promotes increased selection pressure on other traits, leading to increased genetic gain ^[1]. It is hypothesized that MAS using both the bone morphogenetic protein receptor 1B (*BMPR-1B*) and the bone morphogenetic protein 15 (*BMP15*) genes is warranted to increase litter size in sheep and will be of considerable economic value to sheep breeders.

Sakiz is a dairy sheep breed of considerable economic interest to Turkish farmers, mainly due to its high prolificacy

and milk production. These sheep are raised in Cesme, the provinces of Izmir and Aydin, and coastal locations in the Marmara and Aegean regions. The average litter size is 1.7 to 2.3 lambs, and the ovulation rate is 2.9 to 3.3 ^[2,3].

In recent years, a number of natural genetic mutations have been associated with ovulation rates in sheep breeds, including one mutation in the *BMPR-1B* gene (*FecB*) and ten different mutations in the bone morphogenetic protein 15 gene (*FecX^I*, *FecX^H*, *FecX^G*, *FecX^B*, *FecX^R*, *FecX^L*, *FecX^{Gr}*, *FecX^{TT}*, *FecX^O* and *FecX^W*) ^[4-6]. These mutations are significantly associated with the ovulation rate of different sheep breeds.



İletişim (Correspondence)



+90 532 4140333; Fax: +90 224 294 12 02



fbalci@uludag.edu.tr

The *FecB* gene is a dominant autosomal gene responsible for the fecundity of Booroola Merino sheep, and its additive effect on the ovulation rate was first identified in the 1980s [4,7]. The *FecB* gene maps to chromosome 6 in sheep [8]. A point mutation that significantly increases the ovulation rate of ewes is located at base 746 of the coding region (746 A → G) in the highly conserved intracellular kinase signaling domain of the *BMPR-1B*; this mutation causes a glutamine to arginine amino acid substitution [9]. The gene dosage effect of the mutation is additive for the ovulation rate with an increase of 1.5 for each gene copy. These extra ovulations subsequently increased litter size by approximately 1.0 and 1.5, respectively [4,10]. *FecB* gene mutations have been reported in many other sheep breeds, such as Booroola Merino [7], Belclare, Cambridge, China Small Tailed Han, Hu [5], Romney [11], Kendrapada [12], Garole and Javanese sheep [13]. In contrast, *FecB* mutations were not detected in Thoka, Woodlands, Olkuska, Lacaune [13] and Sangsari sheep [14].

The *BMP15* located on the chromosome X [15]. Each *BMP15* gene significantly affects prolificacy [11]. Ewes with two inactive copies of the *BMP15* gene (homozygous animals) are sterile [15,16] and exhibit a similar ovarian phenotype. Ewes with a single inactive *BMP15* gene (heterozygous animals) are fertile and exhibit an increased ovulation rate and an increased incidence of twin or triplet births [4,5,13-18]. In general, heterozygous ewes with mutations in both *FecB* and *FecX^G* exhibited increased fertility compared with ewes harboring a mutation in only one of these genes [6]. The Galway mutation has been identified in Cambridge [16], Garole, Kendrapada [12] and Small Tailed Han sheep [19]. Hanrahan et al. [16] estimated the effect of *FecX^G* to be 0.77 and 1.18 in Belclare and Cambridge ewes, respectively. The *FecX^I* mutation was studied in Inverdale, Belclare, Cambridge, Romney, Hanna and Rasa Aragonese, Cambridge, Small Tailed Han, Sakiz [13,16,19] sheep. Recently, Gursel et al. [20] report that the Inverdale mutation was identified in Sakiz sheep in Turkey.

The *CAST* gene, which is located on sheep chromosome 5, acts on growth characteristics, such as muscle development, birth weight of the lamb, weight at weaning, and postmortem meat qualities, such as tenderness [21-24]. In addition, the *CAST* gene affects cataract development and the reproductive performance of the animals [23-25]. In a study by Garcia et al. [26], calpastatin exerted a significant impact on fertility and longevity on cattle. Based on these results, the use of *CAST* markers together with increased predicted transfer ability (PTA) potential in selection programs for dairy parameters is projected to increase milk yield together with fertility [26].

FecB, *FecX^G* and *FecX^I* mutations have been studied in local sheep breeds, such as Akkaraman, Morkaraman, Daglic, Awassi, Tuj, Karakas and Bafra (Sakiz x Karayaka

cross) in Turkey as well as in Greek Sakiz ewes; however, these mutations were not identified [10,27,28]. In addition, the *CAST* gene has not been studied in Sakiz or other Turkish domestic sheep breeds.

We tested the hypothesis that a major gene was segregating in the Sakiz breed. Therefore, the aim of this study was to investigate *FecB*, *FecX^G*, *FecX^I* and *CAST* gene mutations in the Sakiz sheep breed and the potential introgression of these genes into these breeds to enhance their reproductive rate.

MATERIAL and METHODS

Animals

A total of 71 ewes were used in the study. Ewes were randomly selected from five unrelated herds. These herds were managed within the scope of the Protection of Indigenous Genetic Resources Project under the supervision of the Ministry of Agriculture.

DNA Isolation

Initially, wool samples were prepared by cleaning with EtOH (ethanol). The CTAB (hexa-decyltrimethylammonium bromide) method of DNA isolation was performed using wool samples obtained from selected animals [29]. The amount of DNA obtained and its purity were determined by spectrophotometry (Nanodrop-2000c, Thermo Scientific). The quantified DNA was stored at -80°C until PCR-RFLP (Polymerase chain reaction and Restriction fragment length polymorphism) was performed.

PCR-RFLP Genotyping

DNA was amplified by PCR using the primer sets provided in Table 1. The amplification products were cut with restriction enzymes, and the alleles or mutations were determined. The primer sets used for the amplification of *FecB*, *FecX^G*, *FecX^I*, and *CAST*; the size of the PCR products; the restriction enzymes used and their product sizes and references for PCR-RFLP are provided in Table 1.

For the PCR amplification of *FecB*, *FecX^G*, *FecX^I* and *CAST*, 33.5 µl of dH₂O, 5 µl of 10X buffer, 5 µl of MgSO₄, 1 µl of dNTPs (2.5 mM), 2.5 U Taq DNA polymerase (Biomatik Corp., Cambridge, Canada) and 1 µl of (0.025 µM) forward primer and 1 µl of (0.025 µM) reverse primer from each primer sets presented in Table 1 were used. The final volume was 50 µl after the addition of 3 µl of 100 ng/mL DNA sample.

The PCR conditions for *FecB* [5,13], *FecX^G* [16,27], *FecX^I* [5,15,27] and *CAST* [21,30] were as follows: initial denaturation at 94 to 95°C for 0.30 s to 5 min followed by 33 to 35 cycles of denaturation at 94 to 95°C for 0.30 s to 3 min, annealing for 40 s to 1 min at 60 to 61°C, elongation at

Table 1. Primer sets and enzymes for PCR-RFLP of *FecB*, *BMP15* (*FecX^G*, *FecX^I*) and *CAST* genes**Tablo 1.** PCR-RFLP için *FecB*, *BMP15* (*FecX^G*, *FecX^I*) ve *CAST* genlerinin primer set ve enzimleri

Gene-Allele	Primer Sets*	PP (bp)	RE	RP (bp)		References
<i>FecB</i>	F:5'-CCAGAGGACAATAGCAAAGCAA-3' R:5'-CAAGATGTTTCATGCCTCATCAACAGGTC-3'	190	<i>Avall</i>	AA	190	[5,13]
				BB	159, 31	
				AB	190, 159, 31	
<i>FecX^G</i>	F:5'-CACTGTCTTCTTGTTACTGTATTCAATGAGAC-3' R:5'-GATGCAATACTGCCTGCTTG-3'	141	<i>Hinf I</i>	AA	141	[16,27]
				BB	112, 29	
				AB	141, 112, 29	
<i>FecX^I</i>	F:5'-GAAGTAACCAGTGTCCCTCCACCCTTTTCT-3' R:5'-CATGATTGGGAGAATTGAGACC-3'	154	<i>XbaI</i>	AA	154	[5,15,27]
				BB	124, 30	
				AB	154, 124, 30	
<i>CAST</i>	F:5'-TGGGGCCCAATGACGCCATCGATG-3' R:5'-GGTGGAGCAGCACTTCTGATCACC-3'	622	<i>MspI</i>	AA	336, 268	[21,30]
				BB	622	
				AB	622, 336, 268	

* GenBank accession no: *BMP15*(*FecX^G*, *FecX^I*)(AH009593), *FecB* (AF312016), *CAST* (AF016006-8); (PP: PCR Product, RE: Restriction Enzyme, RP: Restriction Product)





Fig 1. The *FecB* gene cut with the *Avall* restriction enzyme. The products were electrophoresed on a 2.5% agarose gel. No mutations were detected in any of the individuals (M: Marker, Columns 1 to 6: 190 bp band, NC: Negative Control)

Şekil 1. *FecB* geninin *Avall* restriksiyon enzimi ile kesimi (%2.5'lik agaroz jel- tüm bireylerde mutasyon yok) (M: Markör, Sütun1-6: 190 bç'lik bant, NK: Negatif Kontrol)

Fig 2. The *FecX^G* allele cut with the *HinfI* restriction enzyme. The products were electrophoresed on a 4% agarose gel. No mutations were detected in any of the individuals (M: Marker, Columns 1 to 6: 29 and 112 bp bands, NC: Negative Control)

Şekil 2. *FecX^G* allelinin *Hinf I* restriksiyon enzimi ile kesimi (%4'lük agaroz jel- tüm bireylerde mutasyon yok) (M: Markör, Sütun1-6: 29 ve 112 bç'lik bant, NK: Negatif Kontrol)






Fig 3. The *FecX'* allele cut with the *XbaI* restriction enzyme. The products were electrophoresed on a 3.5% agarose gel. No mutations were detected in any of the individuals (M: Marker, Columns 1-5: 154 bp band, NC: Negative Control)

Şekil 3. *FecX'* allelinin *XbaI* restriksiyon enzimi ile kesimi (%3.5'lik agaroz jel- tüm bireylerde mutasyon yok.) (M: Markör, Sütun1-5: 154 bç'lik bant, NK: Negatif Kontrol)




Fig 4. The 622-bp PCR product from the *CAST* gene cut by the *MspI* enzyme. The products were electrophoresed on a 2.5% agarose gel. AA, AB and BB genotypes were detected (M: Marker, Columns 1, 2, 4, 5, 7, 8, 9, 13, 14, 16: 268 and 336 bp bands, Columns 3, 6, 10, 11, 12, 15: 268, 336 and 622 bp bands, Column 17: 622 bp band, NC: Negative Control)

Şekil 4. *CAST* geninin 622 bç'lik PCR ürününün *MspI* enzimi ile kesimi (%2.5'lik agaroz jel görüntüsü - AA, AB, BB genotipi) (M: Markör, Sütun 1,2,4,5,7,8,9,13,14,16: 268 ve 336 bç'lik bant, Sütun3,6,10,11,12,15: 268-336-622 bç'lik bant, Sütun17: 622 bç'lik bant, NK: Negatif Kontrol)

70 to 72°C for 30 s to 2 min, and a final extension at 72°C for 4 to 8 min.

The PCR products for the amplification of the *FecB* gene (product size 190 bp), *FecX^G* allele (product size 141 bp), *FecX'* allele (product size 154 bp) and *CAST* gene (product size 622 bp) were subjected to electrophoresis using 2%, 3%, 2.5% and 1.5% agarose gels, respectively.

For RFLP, 15 µl of the PCR products were incubated with 4 µl buffer, 4 µl dH₂O and 15 U of restriction enzymes at 37°C for 16 h. The restriction digestion products (*FecB*, *FecX^G*, *FecX'* and *CAST*) were visualized in agarose gels stained with 2.5%, 4%, 3.5% and 2.5% ethidium bromide, respectively. The PCR and RFLP product bands were by visualized using the DNr Minilumi imaging system (Fig. 1, 2, 3, 4).

Statistical Analysis

Data were processed by POPGENE V1.32 software to calculate genotypic frequencies and Hardy-Weinberg equilibrium [31].

RESULTS

In the present study, *FecB*, *FecX^G*, *FecX'* and *CAST* mutations were investigated in Sakiz sheep breeds. The agarose gel images of PCR-RFLP products are presented in Fig. 1, 2, 3, 4 respectively. Mutations in the *FecB* gene and at the *FecX^G* and *FecX'* loci in the *BMP15* gene were assessed in the samples. Based on the PCR-RFLP analysis, no digested fragments were observed for *FecB*, *FecX^G* and *FecX'* at 190, 141 and 154 bp, respectively, if the herd did not carry the mutations [5]. All of the 71 individuals were homozygous for *FecB* and *BMP15*. None of the samples harbored the *FecB* mutation, and similarly no *FecX^G* and *FecX'* mutations were noted in the *BMP15* gene. Hardy-Weinberg equilibrium and χ^2 -values were not estimated due to the genotypic frequency.

A total of 71 Sakiz individuals were genotyped. Of these sheep, 42 were AA, 25 were AB and 4 were BB for the *CAST* gene. Genotypic frequencies were estimated as 0.59, 0.36 and 0.05 for AA, AB and BB, respectively, for the *CAST* gene (Table 2). Significant deviation from Hardy-Weinberg

Table 2. Genotypic frequencies of *FecB*, *BMP15* (*FecX^G*, *FecX^I*) and *CAST* genes**Tablo 2.** *FecB*, *BMP15* (*FecX^G*, *FecX^I*) ve *CAST* genlerinin genotipik frekansları

Regions	Allele	n	Frequency	Chi ²	P
<i>FecB</i>	AA	71	1.00	NE	NE
	AB	0	0.00		
	BB	0	0.00		
<i>FecX^G</i>	AA	0	0.00	NE	NE
	AB	0	0.00		
	BB	71	1.00		
<i>FecX^I</i>	AA	71	1.00	NE	NE
	AB	0	0.00		
	BB	0	0.00		
<i>CAST</i>	AA	42	0.59	0.01	0.91
	AB	25	0.36		
	BB	4	0.05		

NE: Not Estimated

equilibrium was not observed for the *CAST* gene in the investigated breed ($P > 0.05$).

DISCUSSION

The results of our study indicate that *FecB*, *FecX^G* and *FecX^I* mutations, which have a major effect on litter size, are not present in Sakiz sheep.

These results are consistent with reports in Romanov, Finn, Thoka, Woodlands, Olkuska, Lacaune [13], Sangsari [14], Akkaraman, Morkaraman, Daglıc, Tuj, Karakas [10], Awassi, Imroz, Kivircik [20] and Bafra [28] sheep breeds wherein *FecB* mutant alleles do not segregate in these breeds. In contrast, *FecB* mutations are reported in several sheep breeds, such as Booroola Merino [7], Belclare, Cambridge, China Small Tailed Han, Hu [5], Garole, Javanese [13], Kendrapada [12] and Romney sheep [17].

The *FecX^G* mutation was analyzed in Garole, Kendrapada [12], Malpura, Deccani, Baluchi [32] and Sangsari sheep [14], however, none of these sheep breeds carried the *FecX^G* mutation in the *BMP15* gene. Additionally, the *FecX^G* mutation was identified in Small Tailed Han, Belclare and Cambridge sheep [1,33]. In contrast with results from Gursel et al. [20], our results indicate that the point mutation in *FecX^G* might not serve as a major gene that influences prolificacy in Sakiz sheep. Galway and Inverdale gene polymorphisms were analyzed as likely candidate genes influencing high prolificacy in Sakiz breeds of Greece, but no polymorphism has been identified [27]. The *FecX^I* mutation was first identified in the Romney sheep breed, which exhibits high fertility traits [1]. Studies indicate that the Small Tailed Han, Hu [33], Olkuska, Garole, Javanese, Woodland, Lacaune [13], Egyptian [34], East Friesian, Finn, Romanov [5] and Greek Sakiz [27] sheep do not carry the *FecX^I*

alleles. Our results demonstrate that the *FecX^I* mutation we examined was not present in Sakiz sheep. In addition our results are consistent with reports by Gursel et al. [20].

In this study, we analyzed a *CAST* gene polymorphism. The genotypic frequencies of the AA, AB and BB alleles for the *CAST* gene were calculated as 0.59, 0.35 and 0.56, respectively. These frequencies are similar to those reported in Arabic [21], Iranian Zel [23] and Makoi sheep [24]. The *CAST* gene, which encodes calpastatin, regulates the activity of calpain as a protease inhibitor [35]. In addition to its role in determining the quality of meat, the *CAST* gene has also effects on reproductive activity [23]. However, studies on reproductive activities are very limited. Garcia et al. [26] reported that the *CAST* gene was strongly correlated with fertility and longevity. However, Byun et al. [35] reported that the *CAST* gene did not affect fertility or longevity in Romney, Corriedale, Merino, Polwarth, Kelso and Coopworth ewes. Proof of the relationship between the *CAST* genes and reproductive activity may explain the variety of litter sizes in Sakiz ewes.

These results suggest that the high prolificacy of the Sakiz breed does not result from *FecB*, *FecX^G* and *FecX^I* mutations. Further investigation should be directed at other loci of the *BMP15* gene or other genes and involve larger sample sizes. Furthermore, the effect of *CAST* gene alleles on reproductive parameters should be investigated. Studies on the genomic aspects of fertility, while improving the accuracy of selection, will allow for more economical and efficient use of resources.

REFERENCES

- Davis GH:** Major genes affecting ovulation rate in sheep. *Genet Sel Evol*, 37 (Suppl. 1): S11-S23, 2005. DOI: 10.1051/gse:2004026
- Işık S:** Bafra koyununun (Sakiz × Karayaka G1) Kazım Karabekir Tarım İşletmesi şartlarında dölverimi, yaşama gücü ve büyüme özellikleri. *Doktora Tezi*, Kafkas Univ. Sağlık Bil. Enst, 2010.
- Ulker H, Bas S:** Koyunlarda major etkili fekonidite genleri. 2- Fekonditeye etki eden mutasyonların henüz belirlenemediği genler. *Hayvansal Üretim*, 50 (1): 68-73, 2009.
- Davis GH:** Fecundity genes in sheep. *Anim Reprod Sci*, 82-83, 247-253, 2004. DOI: 10.1016/j.anireprosci.2004.04.001
- Davis GH, Balakrishnan L, Ross IK, Wilson T, Galloway SM, Lumsden BM, Hanrahan JP, Mullen M, Maoe XZ, Wang GL, Zhaoe ZS, Zenge YQ, Robinson JJ, Mavrogenis AP, Papachristoforou C, Peter C, Baumungi R, Cardyn P, Boujenane I, Cockett NE, Eythorsdottir E, Arranz JJ, Notter DR:** Investigation of the Booroola (*FecB*) and Inverdale (*FecXI*) mutations in 21 prolific breeds and strains of sheep sampled in 13 countries. *Anim Reprod Sci*, 92, 87-96, 2006. DOI: 10.1016/j.anireprosci.2005.06.001
- Mishra C:** Genetic basis of prolificacy in sheep. *Int J Livest Res*, 4, 46-57, 2014. DOI: 10.5455/ijlr.20131227083421
- Piper LR, Bindon BM, Davis GH:** The single gene inheritance of the high litter size of the Booroola Merino. In, Land RB, Robinson DW (Eds): *Genetics of Reproduction in Sheep*. 115-125, Butterworths, London, 1985.
- Montgomery GW, Lord EA, Penty JM, Dodds KG, Broad TE, Cambridge L, Sunden ALF, Stone RT, Crawford AM:** The Booroola fecundity (*FecB*) gene maps to sheep chromosome 6. *Genomics*, 22, 148-153, 1994. DOI: 10.1006/geno.1994.1355

- 9. Mulsant P, Lecerf F, Fabre S, Schibler L, Monget P, Lanneluc I, Pisselet C, Riquet J, Monniaux D, Callebaut I, Cribiu E, Thimonier J, Teyssieri J, Bodin L, Cognie Y, Chitour N, Elsen J:** Mutation in bone morphogenetic protein receptor-1B is associated with increased ovulation rate in Booroola Merino ewes. *Proc Natl Acad Sci U S A*, 98, 5104-5109, 2001. DOI: 10.1073/pnas.091577598
- 10. Karsli T, Balcioğlu MS:** Türkiye'de yetiştirilen altı yerli koyun ırkında BMPR-1B (Booroola) geninde *FecB* allel varlığının PCR-RFLP yöntemiyle araştırılması. *Kafkas Univ Vet Fak Derg*, 16, 1033-1036, 2010. DOI: 10.9775/kvfd.2010.2333
- 11. Wilson T, Wu XY, Juengel JL, Ross, IK, Lumsden JM, Lord, EA, Dodds KG, Walling GA, Mcewan JC, O'connel AR, Mcnatty KP, Montgomery GW:** Highly prolific Booroola sheep have a mutation in the intracellular kinase domain of Bone Morphogenetic Protein 1B Receptor (ALK-6) that is expressed in both oocytes and granulosa cells. *Biol Reprod*, 64, 1225-1235, 2001.
- 12. Kumar S, Mishra AK, Kolte AP, Dash SK, Karim SA:** Screening for Booroola (*FecB*) and Galway (*FecX^G*) mutations in Indian sheep. *Small Rumin Res*, 80, 57-60, 2008. DOI: 10.1016/j.smallrumres.2008.09.007
- 13. Davis GH, Galloway SM, Ross IK, Gregan SM, Ward J, Nimbkar BV, Ghalsasi PM, Nimbkar C, Gray DG, Inounu I, Tiesnamurti B, Martyniuk E, Eythorsdottir E, Mulsant P, Lecerf F, Hanrahan JP, Bradford GH, Wilson T:** DNA tests in prolific sheep from eight countries provide new evidence on origin of the Booroola (*FecB*) mutation. *Biol Reprod*, 66, 1869-1874, 2002.
- 14. Kasiriyani MM, Hafezeyan H, Sayahzadeh H, Jamshidi R, Asgrahi SR, Irajyan GH, Buesagh H:** Genetic polymorphism *FecB* and *BMP15* genes and association with litter size in Sangsari sheep breed of Iran. *J Anim Vet Adv*, 8 (5): 1025-1031, 2009.
- 15. Galloway, SM, Mcnatty KP, Cambridge LM, Laitinen MPE, Juengel JL, Jokiranta TS, McLaren RJ, Luiro K, Dodds KG, Montgomery GW, Beattie AE, Davis GH, Ritvos O:** Mutations in an oocyte-derived growth factor gene (*BMP15*) cause increased ovulation rate and infertility in a dosage-sensitive manner. *Nat Genet*, 25, 279-283, 2000. DOI: 10.1038/77033
- 16. Hanrahan JP, Gregan SM, Mulsant P, Mullen M, Davis GH, Powell R, Galloway SM:** Mutations in the genes for oocyte-derived growth factors *GDF9* and *BMP15* are associated with both increased ovulation rate and sterility in Cambridge and Belclare sheep (*Ovis aries*). *Biol Reprod*, 70, 900-909, 2004. DOI: 10.1095/biolreprod.103.023093
- 17. Davis GH, McEwan JC, Fennessy PF, Dodds KG, Farquhar PA:** Evidence for the presence of a major gene influencing ovulation rate on the x-chromosome of sheep. *Biol Reprod*, 44, 620-624, 1991. DOI: 10.1095/biolreprod44.4.620
- 18. Monteagudo LV, Ponz R, Tejedor MT, Lavina A, Sierra I:** A 17 bp deletion in the Bone Morphogenetic Protein 15 (*BMP15*) gene is associated to increased prolificacy in the Rasa Aragonesa sheep breed. *Anim Reprod Sci*, 110, 139-146, 2009. DOI: 10.1016/j.anireprosci.2008.01.005
- 19. Chu M, Liu ZH, Jiao CL, He YQ, Fang L, Ye SC, Chen GH, Wang JY:** Mutations in *BMPR-1B* and *BMP15* genes are associated with litter size in Small Tailed Han sheep (*Ovis aries*). *J Anim Sci*, 85, 598-603, 2006. DOI: 10.2527/jas.2006-324
- 20. Esen Gursel F, Akis I, Durak H, Mengi A, Oztabak K:** Determination of *BMP15*, *BMPR-1B* and *GDF-9* gene mutations of the indigenous sheep breeds in Turkey. *Kafkas Univ Vet Fak Derg*, 17, 725-729, 2011. DOI: 10.9775/kvfd.2011.4256
- 21. Mohammadi M, Beigi Nasiri MT, Alami-Saeid KH, Fayazi J, Mamooe M, Sadr AS:** Polymorphism of calpastatin gene in Arabic sheep using PCR-RFLP. *Afr J Biotechnol*, 7 (15): 2682-2684, 2008.
- 22. Chung H, Davis M:** PCR-RFLP of the ovine Calpastatin and its association with growth. *Asian J Anim Vet Adv*, 7, 641-652, 2012. DOI: 10.3923/ajava.2012.641.652
- 23. Dehnavi E, Azari MA, Hasani S, Nassiry MR, Mohajer M, Ahmadi ARK:** Genetic variability of calpastatin and calpain genes in Iranian Zel sheep using PCR-RFLP and PCR-SSCP methods. *Iranian J Biotech*, 10 (2): 136-139, 2012.
- 24. Ranjbari M, Hashemi A, Mardani K, Darvishzadeh R:** Allelic polymorphism of Makoei sheep Calpastatin gene identified by polymerase chain reaction and single strand conformation polymorphism. *J Agr Sci Tech*, 14, 533-538, 2012.
- 25. Szkudlarek-Kowalczyk M, Wiśniewska E, Mroczkowski S:** Polymorphisms of calpastatin gene in sheep. *JCEA*, 12, 425-432, 2011. DOI: 10.5513/JCEA01/12.3.934
- 26. Garcia MD, Michal JJ, Gaskins CT, Reeves JJ, Ott TL, Liu Y, Jiang Z:** Significant association of the calpastatin gene with fertility and longevity in dairy cattle. *Anim Genet*, 37, 304-305, 2006. DOI: 10.1111/j.1365-2052.2006.01443.x
- 27. Michaidilis G, Avdi M, Pappa V:** Reproductive performance and investigation of *BMPR-1B* and *BMP15* gene mutations in Greek Chios and Florina sheep breeds. *Arch Zootech*, 11 (1): 24-31, 2008.
- 28. Kurar E, Ozsensoy Y, Bulut Z, Guzeloglu A, Nizamlioglu M:** Investigation of *BMP15* and *BMPR-1B* gene mutations in prolific Sakiz sheep. *Reprod Domest Anim*, 46 (3): 121, 2011.
- 29. Doyle JJ, Doyle JL:** A rapid DNA isolation procedure for small quantities of fresh leaf tissue. *Phytochem Bull*, 19, 11-15, 1987.
- 30. Killefer J, Koohmarai M:** Bovine skeletal muscle calpastatin: Cloning, sequence analysis, and steady-state mRNA expression. *J Anim Sci*, 72, 606-614, 1994.
- 31. Yeh FC, Yang RC, Boyle TBJ, Ye ZH, Mao JX:** Popgene (v.1.32) the user-friendly shareware for population genetic analysis, 2000. <http://www.ualberta.ca/~fyeh/>, Accessed: 01.01.2015.
- 32. Moradband F, Rahimi G, Gholizadeh M:** Association of polymorphisms in fecundity genes of *GDF9*, *BMP15* and *BMP15-1B* with litter size in Iranian Baluchi sheep. *Asian-Aust J Anim Sci*, 24, 1179-1183, 2011. DOI: 10.5713/ajas.2011.10453
- 33. Roy J, Polley S, De S, Mukherjee A, Batabyal S, Pan S, Brahma B, Datta TK, Goswami SL:** Polymorphism of fecundity genes (*FecB*, *FecX*, and *FecG*) in the Indian Bonpala sheep. *Anim Biotechnol*, 22, 151-162, 2011. DOI: 10.1080/10495398.2011.589239
- 34. Abulyazid I, Sharada HM, Abdalla MS, Saleh HM, Hassanin WF:** *FecX¹* gene as a monitor for the production of twins in the Egyptian sheep. *Roavs*, 1 (4): 217-219, 2011.
- 35. Byun SO, Zhou H, Frampton CM, Hickford JGH:** No association between variation in the ovine calpastatin gene and either longevity or fertility in sheep. *Anim Sci*, 41, 222-224, 2009. DOI: 10.1111/j.1365-2052.2009.01982.x