

Effects of Organic and Inorganic Manganese Supplementation on Bone Characteristics, Immune Response to Vaccine and Oxidative Stress Status in Broiler Reared Under High Stocking Density ^[1] ^[2]

Zehra BOZKURT ¹ Tuba BÜLBÜL ² Mehmet Fatih BOZKURT ³
Aziz BÜLBÜL ⁴ Gökhan MARALCAN ⁵ Koray ÇELİKELOĞLU ¹

^[1] This study was supported by the Scientific Research Project Committee of Afyon Kocatepe University, Afyonkarahisar, Turkey (BAPK-07.VF.12); ^[2] This study was presented as a poster in VII. National Animal Nutrition Congress, 26-27 September 2013 - Ankara, Turkey

¹ Afyon Kocatepe University, Faculty of Veterinary Medicine, Department of Animal Science, TR-03200 Afyonkarahisar - TURKEY; ² Afyon Kocatepe University, Faculty of Veterinary Medicine, Department of Animal Nutrition and Nutritional Diseases, TR-03200 Afyonkarahisar - TURKEY; ³ Afyon Kocatepe University, Faculty of Veterinary Medicine, Department of Pathology, TR-03200 Afyonkarahisar - TURKEY; ⁴ Afyon Kocatepe University, Faculty of Veterinary Medicine, Department of Physiology, TR-03200 Afyonkarahisar - TURKEY; ⁵ Afyon Kocatepe University, Faculty of Medicine, Department of Orthopedics and Traumatology, TR-03200 Afyonkarahisar - TURKEY

Article Code: KVFD-2014-12655 Received: 24.11.2014 Accepted: 05.05.2015 Published Online: 12.05.2015

Abstract

This study was carried out to determine the effects of organic and inorganic manganese (Mn) at different levels supplemented to diets of the broilers under stocking density stress on bone characteristics, immune response to vaccine and oxidative stress. A total of 1000, day-old broiler chicks were divided into one control group and nine treatment groups, each consisting of 100 chicks. They were randomly located at a density of 0.064 kg/m² until reaching seven days of age and in the compartments with 2 m² space at a density of 60 kg/m² until they are 49 days old. The control group was fed the basal diet only, whereas experimental groups were fed with basal diet supplemented with organic Mn (6.25, 12.5 and 25 mg/kg of Mn-methionine), inorganic Mn (12.5, 25 and 50 mg/kg Mn-oxide) and organic + inorganic Mn (3.125+6.25, 6.25+12.5 and 12.5+25 mg/kg Mn-methionine + Mn-oxide). There were no changes on days 25 and 49 in terms of physical properties such as humerus, femur and tibia as well as tibia fracture resistance and tibial metaphyseal chondral tissue length in all experimental groups (P>0.05). Tibia bone elasticity was higher in the 50 mg/kg Mn-oxide group compared to the control and other groups on day 49 (P<0.05). The serum calcium level on day 25 increased in the 6.25 and 12.5 mg/kg Mn-methionine and 6.25+12.5 mg/kg Mn-methionine + Mn-oxide groups (P<0.01). Antibody titre against Newcastle Disease Virus in Mn-methionine group on day 25 and titer against Infectious Bursal Disease in Mn-methionine and Mn-oxide groups on day 49 were higher compared to the control group (P<0.01). Serum levels of malondialdehyde and nitric oxide on day 49 decreased in all experimental groups, whereas serum glutathione level increased in the same groups (P<0.01). As a result, dietary Mn supplementation to broilers hosted at 60 kg/m² of stocking density increased immune response and antioxidant activity by reducing the impact of density stress; yet, did not affect the physical properties of the bone. The effects of organic and inorganic forms of Mn on the examined properties were concluded to be similar.

Keywords: Stocking density stress, Organic Mn, Inorganic Mn, Bone characteristic, Immune response, Oxidative stress, Broiler

Sıklığa Maruz Bırakılmış Broilerlerde Rasyona Organik ve İnorganik Manganez İlavesinin Kemik Özellikleri, Aşılar Karşı İmmun Yanıt ve Oksidatif Stres Durumuna Etkileri

Özet

Araştırma sıklık stresi altındaki broilerlerde rasyona farklı düzeylerde organik ve inorganik manganez (Mn) ilave edilmesinin kemik özellikleri, aşılar karşı immün yanıt ve oksidatif stres üzerine etkilerinin incelenmesi amacıyla yapıldı. Araştırmada toplam 1000 adet günlük yaşta broiler civciv her biri 100 civcivden oluşan 1 kontrol ve 9 deneme grubuna ayrıldı. Civcivler 7 günlük yaşa kadar 0.064 kg/m²lik ve 49 günlük yaşa kadar 2 m²lik bölmelerde 60 kg/m² sıklıkta barındırıldı. Kontrol grubu temel rasyonla, deneme grupları ise temel rasyona sırasıyla 6.25, 12.5 ve 25 mg/kg düzeylerinde organik Mn (Mn-metiyonin), 12.5, 25 ve 50 mg/kg inorganik Mn (Mn-oksit), 3.125+6.25, 6.25+12.5 ve 12.5+25 mg/kg organik + inorganik Mn (Mn-metiyonin + Mn-oksit) ilave edilerek beslendi. Araştırmanın 25. ve 49. günlerinde broilerlerde humerus, femur ve tibiaya ait incelenen fiziksel özellikler ile tibia kırılma mukavemetinin ve tibial metafizyal kırıldak doku uzunluğutüm deneme gruplarında değişmediği belirlendi (P>0.05). Tibia kemiği elastikiyeti 49. günde 50 mg/kg Mn-oksit grubunda kontrol ve diğer gruplara göre yüksek bulundu (P<0.05). Serum kalsiyum düzeyinin 25. günde 6.25 ve 12.5 mg/kg Mn-metiyonin ile 6.25+12.5 mg/kg Mn-metiyonin + Mn-oksit gruplarında arttığı belirlendi (P<0.01). Newcastle hastalığı virusuna karşı antikor titresinin 25. günde Mn-metiyonin, Enfeksiyöz Bursal hastalık virusuna karşı antikor titresinin ise 49. günde Mn-metiyonin ve Mn-oksit gruplarında kontrol grubuna göre yüksek olduğu belirlendi (P<0.01). Serumda 49. gündeki malondialdehit ve nitrik oksit düzeyleri tüm deneme gruplarında düşük iken, glutasyon düzeyinin aynı gruplarda arttığı belirlendi (P<0.01). Sonuç olarak 60 kg/m² yerleşim sıklığına barındırılan broilerlerde rasyona Mn ilavesinin sıklık stresinin etkisini azaltarak immün yanıtı ve antioksidan aktiviteyi artırdığı, fiziksel kemik özelliklerini etkilemediği belirlendi. Organik ve inorganik Mn formlarının incelenen özellikler üzerine etkisinin benzer olduğu kanaatine varıldı.

Anahtar sözcükler: Sıklık stresi, Organik Mn, İnorganik Mn, Kemik özelliği, İmmun yanıt, Oksidatif stress, Broiler



İletişim (Correspondence)



+90 272 281312/159, Fax: +90 272 2281349



tbulbul@aku.edu.tr

INTRODUCTION

In the second half of the twentieth century, global poultry meat production has developed more rapidly than any farm animal meat production ^[1]. The poultry meat production which occurred as 103 million tons in 2010 is predicted to reach 122.5 million tons in 2020 ^[2]. This growth springs from modern broiler genotypes obtained from intense selections applied for juvenile growth rate, breast-meat yield and feed efficiency ^[3-5], and intensive production systems where high-tech is used ^[6].

According to European Commission Directive (2007/43/EC) optimum stocking density is 33-39 kg/m² when provided that the owner or keeper complies with the requirements set out in this legislation. In addition, this density can be increased by a maximum of 3 kg/m² when further requirements are provided in the same legislation. However, when the stocking density increases the movement of broiler chickens is restricted largely in the overcrowded conditions and the opportunity to exercise required for bone development is reduced ^[2]. Due to the genetic rehabilitation, bones of broilers develop more slowly than the muscle tissue, and the lack of exercise resulting from high stocking density affects locomotor system reversely. Thus, major foot and leg problems and welfare loss owing to the increase in stocking density were reported ^[7-9]. Especially, the impacts of high stocking density in broilers on the foot health, lameness or gait disturbance, tibial dyschondroplasia, gait scores, carcass bruising and scratching were reported in many researches. At finally, the leg abnormalities result in culling, mortality and economic losses ^[6,10,11].

It is quite clear that, in the perspective of cost-benefit analysis, stocking density is of vital importance in producing more broiler chickens per unit area ^[7,11-13]. In this context, the findings that the environmental conditions are more effective on the health and welfare of the broiler chickens than the stocking density are of vital importance for the economic future of white meat industry ^[10,14-16]. Therefore, reducing the reverse impacts of high stocking density by the modification of environmental conditions, and increasing the upper limit set by the relevant regulations for the stocking density form a significant option ^[17].

In the researches conducted on the feed compositions, the importance of dietary Mn for the commercial broiler sector was especially expressed ^[17-21]. One of the problems caused by the stocking density is the increasing lipid peroxidation in the chickens ^[22,23]. It specifically involves in MnSOD structure of the Manganese. In the case of oxidative stress, MnSOD steps in to protect cell mitochondria from the degrading effects of free oxygen radicals ^[24]. The opportunities of benefiting from minerals taking part in enzymatic activities (Orjinas, piruvate, carboxilase and Mn superoxide dismutase etc.) are being examined in order to reduce these problems ^[10]. The effects of dietary nickel ^[25],

calcium, copper, zinc and iron ^[26,27] in broiler diets have been examined. The importance of manganese (Mn) has been gradually increasing, since it is significant for bone formation, enzyme function and amino acid metabolism in the poultry ^[27,28].

In recent years, many supplementary organic Mn sources have been produced and used in feeding animals; yet, economic reasons overweigh in determining dietary Mn forms being supplemented in conventional broiler diets. The notifications about the sources of Mn additives or the doses administered are conflicting. According to Berta et al. ^[29] and El-Husseiny et al. ^[30] organic forms of Mn are superior to inorganic forms of Mn in terms of utility, whereas there is no difference between the forms of Mn according to Bertechini and Hossain ^[30] and Conly et al. ^[20]. The importance of manganese (Mn) has been gradually increasing since it is significant for bone formation, enzyme function and amino acid metabolism in the poultry ^[28,31].

Therefore, the objective of this study was to determine the effects of organic and inorganic Mn supplementation to the diet of broilers grown under stocking density stress on bone characteristics, immune response and oxidative stress status.

MATERIAL and METHODS

Birds and Management: This study was conducted from May to July at the Afyon Kocatepe University Animal Research and Application Center with the approval of the University Ethics Committee (AKUHADYK-37-2008). A total of 1000, day-old broiler chicks were used in this study. They were divided into one control group and nine treatment groups, each consisting of 100 chicks. Each group was subdivided into four subgroups (replicates) with 25 chicks in each replicate; all subgroups were randomly placed in cages with 1 m² space. Birds were raised until 49 days of age under the same housing conditions and at a stocking density of 60 kg/m². Thus, a total 40 cages were used having 25 birds in each in conformity according to the management guide for Hubbard broilers ^[21]. Feed and water were provided *ad libitum*. A vaccination program for broilers was designed as day 0 with inactive Infectious Bursal Disease (IBD) + Newcastle Disease (ND) vaccine (Gumbopest, Merial RTA, subcutaneously), day 7 with live ND vaccine (Nobilis ND Lasota, Intervet, in drinking water) and infectious bronchitis vaccine (Nobilis, Intervet, in drinking water), day 14 with live IBD vaccine (Bursine Plus, Ford Dodge-Reform, in drinking water), and day 21 with live ND vaccine (Nobilis ND Lasota, Intervet, in drinking water) vaccines.

Dietary Treatments: The control group was fed a basal diet including pre-dominantly of corn, soybean meal and full fat soybean (*Table 1*). The experimental groups were fed basal diets containing three different diet treatments

(organic Mn, inorganic Mn and organic plus inorganic combination) with three different dose levels for each treatment. Each dietary treatment was comprised of 100 birds which were further sub-divided into four replicates, each replicate containing 25 birds. The organic Mn groups were fed diets supplemented with 6.25, 12.5 and 25 mg/kg Mn-methionine, inorganic Mn groups were offered diets containing 12.5, 25 and 50 mg/kg Mn-oxide and organic + inorganic Mn groups were fed diets containing 3.125 mg/kg Mn-methionine + 6.25 mg/kg Mn-oxide, 6.25 mg/kg Mn-methionine + 12.5 mg/kg Mn-oxide and 12.5 mg/kg Mn-methionine + 25 mg/kg Mn-oxide. The nutrient composition of the basal diet (Table 1), including crude protein and calcium was determined according to

the AOAC [32]. The metabolisable energy (ME) of the basal diet was estimated using the Carpenter and Clegg equation [33]. The available phosphorus, lysine, methionine and threonine levels of the basal diet were calculated according to the diet guide for Hubbard broilers [34].

Bone Characteristics: At the age of 25 and 49 days, 8 birds were slaughtered (4 male and 4 female) from each group. The fresh weights of the left tibia, femur and humerus bones (after muscle, cartilage and membranes were removed) of the chicken were measured with 0.01 g precise digital scales; vertical diameters were measured using digital compass; their volumes were determined. Also, fracturing resistance of left tibia bones was determined via three-point break test performed in accordance to Park et al. [35]. Bone density was determined via the values collected with the help of a measuring cylinder and bone weight values. The Tibia weight/length index was calculated by dividing the tibia weight by its length [36].

In order to evaluate the tibial metaphyseal chondral tissue length, right tibial bones of 8 slaughtered birds were separated for histopathological examinations. By opening articulation genus in rear extremities, muscle and other tissues were removed. Cutting tibias from the center, extremities proximalises were taken into buffered formalin solution. After being detected fixed for 72 h, the tissues were taken into Decastro (buffered 3% nitric acid) solution and decalcification was projected. Solutions were renewed every three days. Once the tissues softened, they were washed in running water for 48 h and embedded in paraffin after routine tissue monitoring. They were cut with microtome and stained with hematoxylin eosin, were examined in the light microscope. Slides were examined via DP20 camera set up on Olympus B51 microscope. Via Cell A program, from epiphyseal plates of the bones towards distal (metaphysis), the cartilage tissue was measured upto the area where it transforms into bone tissue (Fig. 1). The length of the cartilage tissue were scored as <1000 μ =1; 1001-1500 μ =2; 1501-2000 μ =3; >2001 μ =4. The ratio of leg abnormalities in the groups were determined using percentage of severely lame birds that clinically lameness or unable to walk.

Serum Parameters: On days 25 and 49 of this study, serums obtained from the blood samples of 8 birds (4 male and 4 female) from each group were stored at -18°C for determination of serum malondialdehyde (MDA), glutathione (GSH), nitric oxide (NO), calcium (Ca) and phosphorus (P) levels and antibody titers against Infectious Bursal Disease (IBD) and Newcastle Disease (ND). The levels of serum MDA, GSH and NO were determined by the methods described by Drappier et al. [37], Beutler et al. [38] and Miranda et al. [39], respectively. Total specific antibody production in serum against IBD and ND was performed by using commercial ELISA kits and in accordance to Puthongsiriporn and Scheideler [40]. Serum Ca and P levels were determined with 300 Alobat Alcycon-auto analyzer.

Table 1. Composition of the basal diets used in different periods (%)

Tablo 1. Farklı dönemlerde kullanılan temel rasyonun bileşimi (%)

| Ingredients | Starter (0 to 12 day) | Grower (12 to 28 day) | Finisher (28 to 49 day) |
|-----------------------------|--------------------------|--------------------------|----------------------------|
| Corn | 47.37 | 50.44 | 51.85 |
| Soybean meal | 24.65 | 17.70 | 14.82 |
| Full fat soybean | 15.00 | 18.00 | 17.00 |
| Poultry meal | 4.00 | 4.00 | 5.00 |
| Meat bone meal | 2.50 | 4.00 | 4.00 |
| Vegetable oil | 3.50 | 3.52 | 4.91 |
| Limestone | 0.73 | 0.59 | 0.61 |
| Dicalcium phosphate | 0.86 | 0.46 | 0.35 |
| DL-Methionine | 0.36 | 0.33 | 0.34 |
| Sodium bicarbonate | 0.30 | 0.25 | 0.31 |
| Vitamin premix* | 0.20 | 0.20 | 0.20 |
| Mineral premix** | 0.10 | 0.10 | 0.15 |
| Salt | 0.14 | 0.12 | 0.07 |
| L-Lysine | 0.13 | 0.13 | 0.23 |
| Enzyme*** | 0.10 | 0.10 | 0.10 |
| Anticoccidial | 0.06 | 0.06 | 0.06 |
| Chemical composition | | | |
| ME**** (kcal/kg) | 3050 | 3200 | 3360 |
| Crude protein, % | 23.20 | 22.70 | 21.80 |
| Calcium, % | 1.00 | 1.00 | 1.00 |
| Phosphorus (available), % | 0.49 | 0.49 | 0.49 |
| Lysine, % | 1.50 | 1.40 | 1.40 |
| Methionine, % | 0.75 | 0.70 | 0.70 |
| Threonin, % | 0.997 | 0.884 | 0.902 |

* Vitamin premix (Rovimix 124-F) provides per 2.5 kg of diets: 15.000.000 IU vitamin A, 1.500.000 IU vitamin D, 50.000 vitamin E, 5.000 mg vitamin K₃, 3.000 mg vitamin B₁, 6.000 mg vitamin B₂, 25.000 mg niacin, 12.000 mg Ca-D Pantothenate, 5.000 mg vitamin B₆, 30 mg vitamin B₁₂, 1.000 mg folic acid, 125 mg D-biotin, 300.000 mg choline chloride, 300.000 L-lysine;
** Mineral premix provides per kg of diets: 80.000 mg manganese, 30.000 mg ferric, 60.000 mg zinc, 5.000 mg copper, 500 mg cobalt, 2.000 mg iodine, 235.680 mg calcium carbonate; *** Enzyme (Optimise M): 5.000 BU/g endo-1.3 beta-glucanase 5.000 BXU/g endo-1.4-ksilinase, 500 FYT/g phytase; ****Estimated according to the Carpenter and Clegg equation




Fig 1. The line in the picture shows that the cartilage tissue was measured for metaphyseal chondral tissue in tibia, HE

Şekil 1. Resimdeki çizgi tibiadaki metafizial kırıldak doku için ölçülen doku uzunluğunu göstermektedir, HE

Statistical Analysis: Differences among group values with regard to all parameters were examined by analysis of variance for all values and the significance of differences between groups tested by Duncan's test. Besides, Mn-methionine, Mn oxide, mixture groups and the control group each of which composing of three groups were considered as single dimensions, and the Contrast test was applied in order for each dimension to be contrasted to the block of other dimensions. A value of $P < 0.05$ was considered as the limit for statistical significance.

RESULTS

Bone Characteristics: On days 25 and 49 of the research, physical properties such as weight and length of humerus, weight and length of femur; weight, length, thickness, density and volume of tibia and tibiotarsi index (Table 2) as well as tibia fracture resistance (Table 3) and tibial metaphyseal chondral tissue length (Table 4) did not change ($P > 0.05$) between the control and experimental groups. Tibia bone elasticity was found higher in the group supplemented with 50 mg/kg Mn-oxide compared to the control and other Mn supplemented groups on day 49 ($P < 0.05$; Table 3).

Serum Parameters: In this study, It was determined that antibody titre against ND in Mn-methionine group on day 25 and titer against IBD in Mn-methionine and Mn-oxide groups on day 49 were higher compared to the control group ($P < 0.01$). At the same time, ND titer on day 25 and IBD titer on day 49 increased in the 12.5 mg/kg Mn-oxide group and in the 12.5+25 mg/kg Mn-methionine + Mn-oxide groups, respectively ($P < 0.01$). Serum Ca level increased in the 6.25 and 12.5 mg/kg Mn-methionine and 6.25+12.5 mg/kg Mn-methionine + Mn-oxide groups compared to the control group on day 25 ($P < 0.01$), whereas Ca level on day 49 and P level on day 25 and 49 in serum did not change in all experimental groups ($P > 0.05$; Table 5).

According to the Contrast test results; In terms of ND antibody level, there were significant differences between Mn-methionine group and the other groups ($P < 0.01$; 25 and 49 day-old chicks); Mn-methionine + Mn-oxide group and the other groups ($P < 0.001$; 25 day-old chicks); the control group and the other groups ($P < 0.001$; 25 and 49 day-old chicks). In terms of IBD antibody level, there were significant differences between Mn-methionine group and the other groups ($P < 0.05$; 49 day-old chicks); the control group and the other groups ($P < 0.01$; 49 day-old chicks). In terms of Serum Ca level, there were significant differences between Mn-methionine group and the other groups ($P < 0.01$; 25 day-old chicks); Mn-oxide group and the other groups ($p < 0.05$; 25day-old chicks), the control group and the other groups ($P < 0.01$; 25 day-old chicks).

Serum GSH and NO levels on days 25 and 49 as well as serum MDA level on day 49 showed significant variations between the experimental groups. Serum levels of MDA and NO on day 49 decreased in the group supplemented with Mn, whereas serum GSH level increased in the same groups ($P < 0.01$). Serum NO level on day 25 was lower ($P < 0.05$) in the 25 mg/kg Mn-oxide and Mn-methionine + Mn-oxide groups compared to the control group (Table 6).

According to the Contrast test results; In terms of MDA level there were significant differences between Mn-methionine + Mn-oxide group and the other groups ($P < 0.001$; 49 day-old chickens), control group and the other groups ($P < 0.01$; 49 day-old chickens). In terms of GSH level, the differences were significant between Mn-oxide group and the other groups ($P < 0.01$; 25 day-old chickens), Mn-methionine + Mn-oxide group and the other groups ($P < 0.01$; 49 day-old chickens), the control group and the other groups ($P < 0.001$; 25 and 49 day-old chickens). In terms of NO level, the differences were significant between Mn-oxide group and the other groups ($P < 0.05$; 25 day-old chickens), Mn-methionine + Mn-oxide

Table 2. Effects of organic and inorganic Mn supplementation to diets of the broilers under stocking density stress on some bone characteristics**Table 2.** Sıklık stresine maruz bırakılmış broyler rasyonlarına organik ve inorganik Mn ilavesinin bazı kemik özelliklerine etkisi

| Parameters | Days | Control | Mn-methionine (mg/kg) | | | Mn-oxide (mg/kg) | | | Mn-methionine + Mn-oxide (mg/kg) | | | SE | P |
|---------------------------------|------|---------|-----------------------|--------|--------|------------------|--------|--------|----------------------------------|-----------|-----------|------|------|
| | | | 6.25 | 12.5 | 25.0 | 12.5 | 25 | 50 | 3.125+6.25 | 6.25+12.5 | 12.5+25.0 | | |
| Humerus weight (g) | 25 | 3.41 | 3.30 | 2.73 | 3.05 | 3.04 | 3.27 | 3.40 | 3.11 | 4.05 | 3.24 | 0.09 | 0.24 |
| | 49 | 6.55 | 7.84 | 7.93 | 8.58 | 7.42 | 7.89 | 7.77 | 7.60 | 7.42 | 6.36 | 0.21 | 0.36 |
| Humerus length (mm) | 25 | 50.35 | 52.37 | 48.84 | 49.19 | 49.97 | 51.11 | 51.68 | 49.01 | 51.91 | 49.26 | 0.36 | 0.23 |
| | 49 | 67.97 | 69.55 | 70.62 | 71.21 | 69.83 | 69.33 | 70.85 | 68.97 | 68.75 | 67.58 | 0.51 | 0.82 |
| Femur weight (g) | 25 | 5.66 | 5.31 | 5.26 | 5.98 | 5.64 | 5.15 | 5.73 | 6.13 | 5.58 | 5.31 | 0.09 | 0.38 |
| | 49 | 10.06 | 10.57 | 12.42 | 11.28 | 11.59 | 10.20 | 11.56 | 9.98 | 12.13 | 9.97 | 0.32 | 0.57 |
| Femur length (mm) | 25 | 47.77 | 54.71 | 54.48 | 55.75 | 55.07 | 54.51 | 55.87 | 55.54 | 54.70 | 54.51 | 0.67 | 0.26 |
| | 49 | 71.33 | 73.26 | 75.09 | 73.38 | 72.72 | 72.29 | 73.90 | 71.82 | 74.03 | 73.46 | 0.47 | 0.85 |
| Tibia weight (g) | 25 | 8.03 | 7.72 | 7.28 | 8.57 | 7.71 | 7.79 | 8.76 | 8.72 | 8.25 | 7.98 | 0.13 | 0.18 |
| | 49 | 14.20 | 14.26 | 14.95 | 16.50 | 16.35 | 14.38 | 17.19 | 12.57 | 15.65 | 13.33 | 0.48 | 0.56 |
| Tibia length (mm) | 25 | 75.83 | 75.67 | 74.99 | 77.99 | 76.53 | 75.44 | 77.78 | 77.81 | 76.49 | 76.01 | 0.31 | 0.34 |
| | 49 | 101.35 | 101.60 | 103.36 | 102.64 | 104.28 | 117.04 | 103.81 | 97.23 | 102.97 | 111.46 | 1.87 | 0.62 |
| Tibia thickness (mm) | 25 | 6.01 | 6.50 | 6.56 | 6.12 | 5.92 | 6.11 | 5.74 | 5.97 | 5.82 | 5.92 | 0.06 | 0.08 |
| | 49 | 7.88 | 8.24 | 8.25 | 8.36 | 8.27 | 7.90 | 8.33 | 8.70 | 8.35 | 8.02 | 0.09 | 0.72 |
| Tibia density | 49 | 1.58 | 1.99 | 1.67 | 2.08 | 2.21 | 1.73 | 1.80 | 1.92 | 2.09 | 1.75 | 0.08 | 0.80 |
| Tibia volume (cm ³) | 49 | 7.61 | 7.42 | 8.60 | 7.79 | 6.92 | 7.08 | 8.48 | 6.81 | 7.03 | 7.76 | 0.30 | 0.94 |
| TibioWt/length index, mg/mm | 25 | 105.92 | 101.82 | 97.05 | 109.50 | 100.75 | 102.98 | 112.48 | 111.79 | 107.83 | 104.56 | 1.41 | 0.26 |
| | 49 | 137.79 | 138.81 | 143.11 | 158.39 | 155.92 | 127.00 | 164.50 | 128.78 | 150.71 | 124.34 | 3.94 | 0.30 |

Table 3. Effects of organic and inorganic Mn supplementation to diets of the broilers under stocking density stress on fracture characteristics of tibia bone**Table 3.** Sıklık stresine maruz bırakılmış broyler rasyonlarına organik ve inorganik Mn ilavesinin tibia kemiği kırılma özellikleri üzerine etkisi

| Parameters | Days | Control | Mn-methionine (mg/kg) | | | Mn-oxide (mg/kg) | | | Mn-methionine + Mn-oxide (mg/kg) | | | SE | P |
|--|------|--------------------|-----------------------|--------------------|--------------------|-------------------|--------------------|-------------------|----------------------------------|-------------------|-------------------|-------|------|
| | | | 6.25 | 12.5 | 25.0 | 12.5 | 25 | 50 | 3.125+6.25 | 6.25+12.5 | 12.5+25.0 | | |
| Maximum fracture height (N) | 25 | 208.98 | 218.46 | 265.64 | 259.70 | 190.80 | 223.95 | 215.20 | 226.05 | 208.51 | 222.41 | 5.68 | 0.13 |
| | 49 | 231.79 | 288.00 | 283.96 | 332.44 | 283.69 | 269.00 | 285.82 | 356.66 | 270.85 | 250.76 | 11.18 | 0.39 |
| Tibia bone elasticity (mm/s) | 25 | 1.69 | 1.72 | 1.67 | 1.68 | 1.57 | 1.70 | 1.73 | 1.51 | 1.76 | 1.96 | 0.03 | 0.23 |
| | 49 | 1.97 ^{bc} | 2.06 ^{bc} | 2.07 ^{bc} | 2.40 ^{ab} | 1.89 ^c | 1.98 ^{bc} | 2.64 ^a | 2.01 ^{bc} | 2.14 ^c | 1.88 ^c | 0.05 | 0.04 |
| Tibia fracture resistance (N/mm) MPa | 25 | 73.62 | 67.35 | 72.91 | 87.00 | 71.37 | 76.61 | 82.96 | 79.88 | 80.74 | 83.91 | 1.72 | 0.15 |
| | 49 | 60.36 | 65.81 | 64.39 | 67.50 | 63.93 | 68.75 | 61.07 | 67.85 | 58.55 | 63.05 | 1.64 | 0.96 |
| Tibia fracture resistance Kgf (N/9.81) | 25 | 21.30 | 22.27 | 27.08 | 26.47 | 19.44 | 22.83 | 21.94 | 23.04 | 21.26 | 22.67 | 0.58 | 0.13 |
| | 49 | 23.62 | 29.36 | 28.95 | 33.89 | 28.92 | 27.42 | 29.13 | 36.36 | 27.61 | 25.57 | 1.14 | 0.40 |

Letters (a, b, c) in the same line indicate significant differences between different letters; Tibia bone flexibility (mm/second): The collapse amount of the bone during the fracturing process; millimeter/second; Tibia bone resistance (kgf (N/9.81): Maximum force that the bone could endure during the fracturing; kgf, N/mm

group and the other groups ($P < 0.05$; 49 day-old chickens), the control group and the other groups ($P < 0.05$; 25 and 49 day-old chickens).

DISCUSSION

In this study, physical properties of the bones such as humerus, femur and tibia and Ca levels in serum (other than Ca level on day 25), as well as tibia fracturing properties (other than tibia elasticity on day 49) were not affected from Mn supplementation. It was observed that

the values of tibia fracturing resistance were found to have numerically increased in Mn-methionine groups. Also, tibia fracturing resistance was detected to be the lowest in the control group which Mn supplementation was not applied. The fact that a statistically significant difference was not observed in the present study may be due to the high standard error resulting from small sampling size (n). As a matter of fact our finding corresponds with the research results asserting that Mn contributes to bone properties. Ruff and Hughes ^[41] and El Hussein et al. ^[30] also reported similar results and the researchers indicated that dietary

Table 4. Effects of organic and inorganic Mn supplementation to diets of the broilers under stocking density stress on tibial metaphyseal chondral tissue length, distribution of tibial metaphyseal chondral tissue length score and ratio of leg abnormalities**Tablo 4.** Sıklık stresine maruz bırakılmış broyler rasyonlarına organik ve inorganik Mn ilavesinin tibialarda metafizial kıkırdak doku uzunluğu, metafizial kıkırdak doku skorlarının dağılımı ve bacak problemleri oranı üzerine etkisi

| Groups | | 25 th day | | | | 49 th day | | | | | | |
|-------------------|------------|--------------------------------------|-------------------------------|-------|---|----------------------|--------------------------------------|--|---|-------|---|------------------------|
| | | Cartilage Tissue Length Mean (µm) | Cartilage Tissue Length Score | | | | Cartilage Tissue Length Mean (µm) | Cartilage Tissue Length Severity Score | | | | Leg Abnormalities % |
| | | | 1 | 2 | 3 | 4 | | 1 | 2 | 3 | 4 | |
| Control | | 2154.22 | | 1 | 1 | 2 | 3020.85 | | 3 | 2 | 3 | 5.60 |
| Mn-meth | 6.25 | 2382.90 | | 1 | | 6 | 1676.16 | 1 | 2 | 3 | 2 | 2.59 |
| | 12.5 | 2109.89 | | 2 | | 4 | 1934.09 | 1 | 6 | | 1 | 2.73 |
| | 25 | 3204.90 | | | 1 | 6 | 1261.89 | 1 | 4 | 3 | | 1.26 |
| Mn-oxide | 12.5 | 2032.29 | 2 | | | 5 | 1536.79 | | 2 | 5 | | 9.70 |
| | 25 | 2020.12 | | | 2 | 2 | 1613.41 | 1 | 4 | 1 | 2 | 4.17 |
| | 50 | 2499.51 | | 1 | | 5 | 1567.28 | | 5 | 2 | 2 | 2.63 |
| Mn-meth +Mn-oxide | 3.125+6.25 | 2320.49 | | 1 | 2 | 4 | 1687.73 | | 4 | 2 | 2 | 2.50 |
| | 6.25+12.5 | 2422.62 | | | 2 | 4 | 1646.31 | 1 | 1 | 4 | 2 | 0 |
| | 12.5+25.0 | 2541.83 | | 1 | | 4 | 1411.41 | 1 | 4 | 3 | | 2.70 |
| SEM | | 140.73 | | 0.11 | | | 143.94 | | | 0.01 | | |
| P | | 0.767 | | 0.919 | | | 0.328 | | | 0.538 | | |

Severity score of tibial metaphyseal chondral tissue: <1000µ=1; 1001-1500µ=2; 1501-2000µ=3; >2001µ=4

Table 5. Effects of organic and inorganic Mn supplementation to diets of the broilers under stocking density stress on ND and IBD antibody titers and Ca (mg/dl) ve P (mg/dl) levels in serum**Tablo 5.** Sıklık stresine maruz bırakılmış broyler rasyonlarına organik ve inorganik Mn ilavesinin serum ND ve IBD antikor titresi ile Ca (mg/dl) ve P (mg/dl) düzeyi üzerine etkisi

| Parameters | Days | Control | Mn-methionine (mg/kg) | | | Mn-oxide (mg/kg) | | | Mn-methionine+Mn-oxide (mg/kg) | | | SE | P |
|------------|------|--------------------|-----------------------|--------------------|--------------------|---------------------|---------------------|---------------------|--------------------------------|---------------------|---------------------|-------|------|
| | | | 6.25 | 12.5 | 25.0 | 12.5 | 25 | 50 | 3.125+6.25 | 6.25+12.5 | 12.5+25.0 | | |
| ND | 25 | 2308 ^c | 7287 ^b | 11352 ^a | 7390 ^b | 7829 ^b | 5259 ^{bc} | 5313 ^{bc} | 2320 ^c | 2164 ^c | 2638 ^c | 457.2 | 0.00 |
| | 49 | 9266 ^b | 11177 ^{ab} | 15591 ^a | 14873 ^a | 11221 ^{ab} | 14095 ^{ab} | 9080 ^b | 10391 ^{ab} | 12260 ^{ab} | 11672 ^{ab} | 540.5 | 0.06 |
| IBD | 25 | 15956 | 16675 | 16207 | 13973 | 13868 | 14168 | 17738 | 13737 | 12304 | 11646 | 668.4 | 0.55 |
| | 49 | 10492 ^c | 15195 ^a | 17417 ^a | 15903 ^a | 14758 ^a | 14663 ^{ab} | 11175 ^{ab} | 11782 ^c | 16041 ^{cb} | 15511 ^a | 397.9 | 0.00 |
| Ca | 25 | 4.89 ^d | 6.11 ^b | 7.19 ^a | 5.22 ^{cd} | 4.90 ^d | 5.07 ^{cd} | 5.37 ^{bcd} | 4.76 ^d | 5.91 ^{bc} | 5.16 ^{cd} | 0.11 | 0.00 |
| | 49 | 5.91 | 7.73 | 5.96 | 7.54 | 7.50 | 7.76 | 6.65 | 5.98 | 7.00 | 6.16 | 0.22 | 0.28 |
| P | 25 | 4.24 | 4.07 | 4.19 | 4.18 | 3.80 | 3.92 | 3.72 | 3.57 | 3.70 | 4.22 | 0.07 | 0.33 |
| | 49 | 5.98 | 7.85 | 5.99 | 6.84 | 6.29 | 7.05 | 5.82 | 5.51 | 6.16 | 5.89 | 0.21 | 0.34 |

Letters (a, b, c, d) in the same line indicate significant differences between different letters; Data are means of 8 replicate cages consisting of 12 birds per replicate cage

Table 6. Effects of organic and inorganic Mn supplementation to diets of the broilers under stocking density stress on oxidative stress status**Tablo 6.** Sıklık stresine maruz bırakılmış broyler rasyonlarına organik ve inorganik Mn ilavesinin oksidatif stres durumu üzerine etkisi

| Parameters | Days | Control | Mn-methionine (mg/kg) | | | Mn-oxide (mg/kg) | | | Mn-methionine+Mn-oxide (mg/kg) | | | SE | P |
|------------|------|--------------------|-----------------------|----------------------|----------------------|---------------------|---------------------|----------------------|--------------------------------|---------------------|---------------------|------|------|
| | | | 6.25 | 12.5 | 25.0 | 12.5 | 25 | 50 | 3.125+6.25 | 6.25+12.5 | 12.5+25.0 | | |
| MDA | 25 | 280.62 | 252.25 | 258.50 | 250.00 | 247.25 | 260.37 | 240.25 | 212.62 | 227.75 | 225.25 | 6.09 | 0.40 |
| | 49 | 3.71 ^a | 2.89 ^{bcd} | 3.06 ^{bcd} | 2.62 ^d | 3.28 ^{abc} | 2.91 ^{bcd} | 2.77 ^{cd} | 3.03 ^{bcd} | 2.88 ^{bcd} | 3.35 ^a | 0.05 | 0.01 |
| GSH | 25 | 22.47 ^c | 24.76 ^b | 25.87 ^{ab} | 24.87 ^b | 25.75 ^{ab} | 26.77 ^a | 25.06 ^{ab} | 25.14 ^{ab} | 26.01 ^{ab} | 26.04 ^{ab} | 0.20 | 0.00 |
| | 49 | 17.43 ^d | 20.32 ^c | 23.24 ^{ab} | 25.22 ^a | 22.33 ^c | 25.09 ^{ab} | 23.67 ^{ab} | 23.74 ^{ab} | 24.93 ^{ab} | 26.07 ^a | 0.38 | 0.00 |
| NO | 25 | 13.62 ^a | 13.06 ^{ab} | 11.88 ^{abc} | 11.73 ^{abc} | 12.91 ^{ab} | 11.03 ^{bc} | 11.56 ^{abc} | 11.35 ^{bc} | 11.44 ^{bc} | 10.63 ^c | 0.22 | 0.04 |
| | 49 | 12.80 ^a | 11.28 ^b | 11.39 ^b | 10.06 ^{bc} | 9.85 ^c | 10.10 ^{bc} | 10.40 ^{bc} | 11.20 ^b | 10.22 ^{bc} | 9.39 ^c | 0.18 | 0.00 |

Letters (a, b, c, d) in the same line indicate significant differences between different letters

manganese improved bone features. Also, we found the largest tibial metaphyseal chondral tissue in control group at 49 days of age (Table 4). In general, it has been known that increase in the length of tibial metaphyseal chondral tissue is related to a predisposing factor for tibial dyschondroplasia^[42,43]. It can be argued that regarding to those results, Manganese supplementation into diet decreased the incidence of tibial dyschondroplasia in broilers reared under high stock density.

The positive effect of manganese on tibia in this study has suggested that it enhanced Ca accumulation in tibia. In this study, Ca levels in the groups Mn was supplemented were lower than the control group without Mn, and that difference reflected on the Contrast test. Sunder et al.^[44] also reported that 100 ppm of dietary Mn supplementation contributed to foot-leg health. However, these differences in calcium did not reflect upon P level. Therefore, it must be considered that dietary Mn supplementation especially during the growth period might disrupt the ratio of Ca and P. Furthermore, while the number of the chickens with clinical lameness or foot discomfort was at acceptable levels (2-3%) in the groups other than the group with 12.5 mg/kg of inorganic Mn and the control group despite the bone parameters and increasing serum Ca level, this rate was found to be numerically higher (5.6%) in the control group.

The present study revealed that broilers supplemented with dietary Mn responded to ND and IBD vaccines by producing antibodies at higher levels than chickens fed on the control feed with no Mn supplementation. Both ND and IBD antibody levels elevated more particularly in Mn methionine groups when compared to other groups (Table 5). These findings suggest that Mn enhances the immune system in broilers. Especially organic manganese has a more powerful impact on that. Similarly, Sunder et al.^[44] reported that dietary Mn supplementation at the level of 100 mg/kg enhanced immunity. Gajula et al.^[45] also asserted that immune response became swifter as dietary Mn level increased. Moreover, as the stress caused by the exposure to stocking density was reduced to some extent by the feed with Mn supplementation, the immunity of the chickens in the control group with no Mn supplementation may have been affected from this stress more. Similarly, Bozkurt et al.^[46] observed that the *bursa of fabricius* weight did not change in the groups supplemented with organic and inorganic Mn.

In this study, serum MDA and NO levels of the broilers fed with dietary Mn supplementation were found to be lower compared to the control group (Table 6). Similarly, Bulbul et al.^[47] reported that organic or inorganic dietary Mn supplementation reduced serum MDA. However, Lu et al.^[48] reported that dietary Mn supplementation did not affect serum MDA level in broilers. In this study, high MDA level in the control group may be related to abdominal fat amount. The deterioration in oxidant-antioxidant balance in favor of antioxidants in the control group may have led

to high MDA and NO levels causing high feed consumption and fat accumulation, consequently. In fact, it was observed that less abdominal fat was numerically or statistically detected in broilers fed with Mn supplemented feed than in the control group except for the group feeding with 25 mg/kg Mn-oxide^[46]. It has also been reported that redundant free fatty acid intake exceeding antioxidant capacity may increase oxidative stress by leading to lipid peroxidation and also bring molecular features of fat tissue into the forefront as a major cause for oxidative stress^[49]. Lu et al.^[48] also reported that 100 mg/kg of dietary Mn supplementation lowered abdominal fat amount. Furthermore, Mn supplementation to the diet above the rate determined for broilers by NRC may suggest that Mn has inhibited the activities of enzymes involving in lipid peroxidation.

As a result, it was observed that dietary Mn supplementation enhanced immune response and antioxidant activity by reducing the effect of density stress for broilers having been hosted at 60 kg/m² stocking density; slightly increased ossification in tibia while it did not affect the physical properties of the bone or bone resistance. Considering the levels used at the study, organic and inorganic forms of Mn have similar impacts on examined properties; yet, further research on the impacts of higher doses of Mn on ossification is required.

ACKNOWLEDGEMENTS

We are grateful to the Afyon Kocatepe University, Animal Husbandry Research and Application Center.

REFERENCES

- 1. Chang HS:** Overview of the world broiler industry: Implications for the Philippines. *AJAD*, 4, 67-82, 2007.
- 2. OECD/FAO:** OECD-FAO Agricultural Outlook 2011-2020, Chapter 7: Meat. Paris/Rome: Organisation for Economic Co-operation and Development and Food and Agriculture Organization of the United Nations (OECD-FAO AGRICULTURAL OUTLOOK 2011-2020 © OECD/FAO 2011), 2012.
- 3. Bessei W:** Welfare of broilers: A review. *World Poultry Sci J*, 62, 455-466, 2006. DOI: 10.1017/S0043933906001085
- 4. Arnould C, Leterrier C:** Welfare of chickens reared for meat production. *Prod Anim*, 20, 41-45, 2007.
- 5. Estevez I:** Density allowances for broilers: Where to set the limits? *Poult Sci*, 86, 1265-1272, 2007. DOI: 10.1093/ps/86.6.1265
- 6. Dawkins MS, Layton R:** Breeding for better welfare: genetic goals for broiler chickens and their parents. *Animal Welfare*, 21, 147-155, 2012.
- 7. Kestin SC, Knowles TG, Tinch AE, Gregory N:** Prevalence of leg weakness in broiler chickens and its relationship with genotype. *Vet Rec*, 131, 190-194, 1992. DOI: 10.1136/vr.131.9.190
- 8. Martrechar A, Boilletot E, Huonnic D, Pol F:** Risk factors for on ascitis incidence in broiler chickens. *Poult Sci*, 74, 1586-1590, 2002.
- 9. Lynch M, Thorp BH, Whitehead CC:** Avian tibial dyschondroplasia as a cause of bone deformity. *Avian Pathol*, 21, 275-285, 1992. DOI: 10.1080/03079459208418842
- 10. Dawkins MS, Donnelly CA, Jones TA:** Chicken welfare is influenced more by housing conditions than by stocking density. *Nature*, 22, 342-

344, 2004. DOI: 10.1038/nature02226

11. Bradshaw RH, Kirkden RD, Broom DM: A review of the aetiology and pathology of leg weakness in broilers in relation to welfare. *Avian Poultry Biol Rev*, 13, 45-103, 2002.

12. Sanotra GS, Lund JD, Ersboll AK, Petersen JS, Vestergaard KS: Monitoring leg problems in broilers: A survey of commercial broiler production in Denmark. *World Poult Sci J*, 57, 55-67, 2001. DOI: 10.1079/WPS20010006

13. Knowles TG, Kestin SC, Haslam SM, Brown SN, Green LE, Butterworth A, Pope SJ, Pfeiffer D, Nicol CJ: Leg disorders in broiler chickens: Prevalence, risk factors and prevention. *PLoS One*, 3, e1545, 2008. DOI: 10.1371/journal.pone.0001545

14. Jones TA, Donnelly CA, Dawkins MS: Environmental and management factors affecting the welfare of chickens on commercial farms in the United Kingdom and Denmark stocked at five densities. *Poult Sci*, 84, 1155-1165, 2005.

15. Blokhuis HJ: Intensive production units and welfare: Domestic fowl. *Rev Sci Tec*, 13, 67-78, 1994.

16. Elson HA: Poultry housing and husbandry. *Br Poult Sci*, 51, 23-34, 2010. DOI: 10.1080/00071668.2010.497300

17. European Commission: Council Directive 2007/43/EC of 28 June 2007 laying down minimum rules for the protection of chickens kept for meat production. *Official Journal of the European Union*, 2007.

18. Henry PR, Ammerman CB, Littell RC: Relative bioavailability of manganese from a manganese-methionine complex and inorganic sources for ruminants. *J Dairy Sci*, 75, 3473-3478, 1992.

19. Henry PR, Ammerman CB, Miles RD: Relative bioavailability of manganese in a manganese-methionine complex for broiler chicks. *Poult Sci*, 68, 107-112, 1989. DOI: 10.3382/ps.0680107

20. Conly AK, Poureslami R, Koutsos EA, Batal AB, Jung B, Beckstead R, Peterson DG: Tolerance and efficacy of tribasic manganese chloride in growing broiler chickens. *Poult Sci*, 91, 1633-1640, 2012. DOI: 10.3382/ps.2011-02056

21. Yang XJ, Sun XX, Li CY, Wu XH, Yao JH: Effects of copper, iron, zinc, and manganese supplementation in a corn and soybean meal diet on the growth performance, meat quality, and immune responses of broiler chickens. *J Appl Poult Res*, 20, 263-271, 2011. DOI: 10.3382/japr.2010-00204

22. Simsek UG, Dalkilic B, Ciftci M, Yuce A: The influences of different stocking densities on some welfare indicators, lipid peroxidation (MDA) and antioxidant enzyme activities (GSH, GSH-Px, CAT) in broiler chickens. *JAVA*, 8, 1568-1572, 2009.

23. Simitzis PE, Kalogeraki E, Goliomytis M, Charismiadiou MA, Triantaphyllopoulos K, Ayoutanti A, Niforou K, Hager-Theodorides AL, Deligeorgis SG: Impact of stocking density on broiler growth performance, meat characteristics, behavioural components and indicators of physiological and oxidative stress. *Br Poult Sci*, 53, 721-30, 2012. DOI: 10.1080/00071668.2012.745930

24. Lu L, Luo XG, Ji C, Liu B, Yu SX: Effect of manganese supplementation and source on carcass traits, meat quality, and lipid oxidation in broilers. *Anim Sci*, 85, 812-822, 2007. DOI: 10.2527/jas.2006-229

25. Wilson JH, Wilson EJ, Ruzler PL: Dietary nickel improves male broiler (*Gallus domesticus*) bone strength. *Biol Trace Elem Res*, 83, 239-249, 2001. DOI: 10.1385/BTER:83:3:239

26. Guinotte F, Nys Y, Monredon F: The effects of particle size and origin of calcium carbonate on performance and ossification characteristics in broiler chicks. *Poult Sci*, 70, 1908-1920, 2001. DOI: 10.3382/ps.0701908

27. Bao YM, Choct M, Iji PA, Bruerton K: Effect of organically complexed copper, iron, manganese, and zinc on broiler performance, mineral excretion, and accumulation in tissues. *J Appl Poult Res*, 16, 448-455, 2007. DOI: 10.1093/japr/16.3.448

28. Scott ML, Nesheim MC, Young RJ: Nutrition of the chicken. M.L. Ithaca, NY, 1976.

29. Berta E, Andrásófszky E, Bersényi A, Glávits R, Gáspárdy A, Fekete SG: Effect of inorganic and organic manganese supplementation on the performance and tissue manganese content of broiler chicks.

Acta Vet Hung, 52, 199-209, 2004. DOI: 10.1556/AVet.52.2004.2.8

30. El-Husseiny OM, Hashish SM, Ali RA, Arafa SA: Effects of feeding organic zinc, manganese and copper on broiler growth, carcass characteristics, bone quality and mineral content in bone, liver and excreta. *Int J Poult Sci*, 11, 368-377, 2012. DOI: 10.3923/ijps.2012.368.377

31. Ji F, Luo XG, Lu L, Liu B, Yu SX: Effect of manganese source on manganese absorption by the intestine of broilers. *Poult Sci*, 85, 1947-1952, 2006. DOI: 10.1093/ps/85.11.1947

32. AOAC (Association of Official Analytical Chemists): Official Methods of Analysis. 17th ed., AOAC International, Maryland, USA, 2000.

33. Leeson S, Summers JD: Nutrition of the Chicken. University Books, Guelph, 2001.

34. Hubbard LLC: Management guide for broiler. Hubbard L.L.C., Duluth GA 30096-USA, 2004.

35. Park SY, Birkhold SG, Kubena LF, Nisbet DJ, Ricke SC: Effect of storage condition on bone breaking strength and bone ash in laying hens at different stages in production cycles. *Poult Sci*, 82, 1688-1691, 2003.

36. Seedor JG, Quarruccio HA, Thompson DD: The bisphosphonate alendronate (MK-217) inhibits bone loss due to ovariectomy in rats. *J Bone Miner Res*, 6, 339-346, 1991. DOI: 10.1359/jbmr.2005.20.2.354

37. Draper HH, McGirr LG, Hadley M: The metabolism of malandialdehyde. *Lipids*, 21, 305-307, 1986. DOI: 10.1007/BF02536418

38. Beutler E, Gelbart T, Pegelow C: Erythrocyte glutathione synthetase deficiency leads not only to glutathione but also to glutathione-S-transferase deficiency. *J Clin Invest*, 77, 38-41, 1986. DOI: 10.1172/JCI112298

39. Miranda KM, Espey MG, Wink AD: A rapid, simple spectrophotometric method for simultaneous detection of nitrate and nitrite. *Nitric Oxide*, 5, 62-71, 2001. DOI: 10.1006/niox.2000.0319

40. Puthongsiripors U, Scheideler SE: Effects of dietary ratio of linoleic to linolenic acid on performance, antibody production, and *in vitro* lymphocyte proliferation in two strains of leghorn pullet chicks. *Poult Sci*, 84, 846-857, 2005.

41. Ruff CR, Hughes BL: Bone strength of height-restricted broilers as affected by levels of calcium, phosphorus, and manganese. *Poult Sci*, 64, 1628-1636, 1985. DOI: 10.3382/ps.0641628

42. Edwards HM: Studies on the etiology of tibial dyschondroplasia in chickens. *J Nutr*, 114, 1001-1013, 1984.

43. Leach RM, Nesheim MC: Nutritional genetic and morphological studies of an abnormal cartilage formation in young chicks. *J Nutr*, 86, 236-244, 1965.

44. Sunder GS, Panda AK, Gopinath NCS, Mantena VLN, Savaram RR, Chalasani VK: Effect of supplemental manganese on mineral uptake by tissue and immune response in broiler chickens. *J Poult Sci*, 43, 371-377, 2006. DOI: 10.2141/jpsa.43.371

45. Gajula SS, Chelasani VK, Panda AK, Mantena VL, Savaram RR: Effect of supplemental inorganic Zn and Mn and their interactions on performance of broiler chicken, mineral bioavailability and immune response. *Biol Trace Elem Res*, 139, 177-187, 2011. DOI: 10.1007/s12011-010-8647-8

46. Bozkurt Z, Bulbul T, Bulbul A, Tekerli M, Köken M. Effects of dietary organic and inorganic manganese supplementation on growth performance and carcass traits in broiler reared under stocking density stress. VII. *National Animal Nutrition Congress*, 158-163, 26-27 September 2013 - Ankara, Turkey.

47. Bulbul A, Bulbul T, Kucukersan S, Sireli M, Eryavuz A: Effects of dietary supplementation of organic and inorganic Zn, Cu and Mn on oxidant/antioxidant balance in laying hens. *Kafkas Univ Vet Fak Derg*, 14, 19-24, 2008. DOI: 10.9775/kvfd.2007.36-A

48. Lu L, Ji C, Luo XG, Liu B, Yu SX: The effects of supplemental manganese in broiler diets on abdominal fat deposition and meat quality. *Anim Feed Sci Technol*, 129, 49-59, 2006. DOI: 10.1016/j.anifeedsci.2005.12.005

49. Vincent HK, Taylor AG: Biomarkers and potential mechanisms of obesity-induced oxidant stress in humans. *Int J Obes (Lond)*, 30, 400-18, 2006. DOI: 10.1038/sj.ijo.0803177