

Characterization and Distribution of *Salmonella* spp. Isolated from Poultry Slaughterhouse-processing in Shandong Province ^[1]

Wenyan GAI ¹ Junwei WANG ¹ Juan WANG ¹ Zhigang CUI ²
Zhina QU ¹ Jun HONG ¹ Jinghua CUI ² Xiaoli DU ²
Xiumei HUANG ¹ Xumin CAO ¹ Jianmei ZHAO ¹

^[1] This work was supported by the Project of Ministry of Agriculture Agricultural Product Quality and Safety Supervision of China (GJFP2014007)

¹ China Animal Health and Epidemiology Center, Laboratory of Quality & Safety Risk Assessment for Animal Products of Ministry of Agriculture, 266032 Qingdao - CHINA

² Chinese Center for Disease Control and Prevention, PulseNetChina, 102206 Beijing - CHINA

Article Code: KVFD-2016-14974 Received: 06.01.2016 Accepted: 07.03.2016 Published Online: 07.03.2016

Abstract

Salmonella spp. is the most important food-borne pathogens of public health interest incriminated in poultry meat worldwide. The purpose of this study to estimate the prevalence *Salmonella* spp. contamination in poultry products from 12 different located geographical areas from among the 12 poultry slaughterhouses authorized in China and to characterize all the isolates by serotypes, PFGE patterns, MLST patterns and antimicrobial susceptibility. The prevalence of *Salmonella* spp. in the poultry slaughterhouse was 10.4%. All 100 strains of *Salmonella* spp. comprising 13 majority serotypes were identified. *S. Enteritidis* was the most frequently isolated from samples. The isolates displayed resistance to sulfisoxazole (SF) (79.0%), doxycycline (DOX) (68.0%), tetracycline (TE) (65.0%), florfenicol (FFC) (64.0%), ampicillin (AM) (50.0%), gentamicin (GM) (48.0%), trimethoprim-sulfamethoxazole (SXT) (30.0%), spectinomycin (SPT) (30.0%), enrofloxacin (ENR) (10.0%), ofloxacin (NOR) (10.0%), amoxicillin potassium clavulanate (AC) (1.0%) and polymyxin (PME) (4.0%). The *Salmonella* spp. isolates were not resistant to cefotaxime (EFT). Each isolates were multi-drug resistant as they were resistant to at least 2 groups of antimicrobials. Four clusters and 38 fingerprint-patterns generated by PFGE were identified among strains recovered from various locations, providing information on associations among the strains as well as evidence of the existence of persistent strains in some areas. MLST analysis of isolates identified the 10 STs, the 7 housekeeping genes had the different variation.

Keywords: *Salmonella* spp., Poultry slaughterhouse-processing, Serotyping, PFGE, MLST, Antimicrobial susceptibility

Shandong Eyaletindeki Kanatlı Hayvan Kesimhanelerinden İzole Edilen *Salmonella* spp.'nin Karakterizasyonu ve Yaygınlığı

Özet

Salmonella spp. dünya çapında kanatlı hayvan etlerinden köken alan halk sağlığı bakımından en önemli gıda kaynaklı patojendir. Bu çalışmanın amacı Çin'de 12 değişik bölgede yer alan kesimhanelerde kesilen kanatlı hayvanlara ait ürünlerde *Salmonella* spp. prevalansını tespit etmek ve tüm izolatları serotipleri, PFGE ve MLST görüntüleri ile antimikrobiyal duyarlılıklarına göre karakterize etmektir. Çalışmada *Salmonella* spp. prevalansı kanatlı kesimhanelerinde %10.4 olarak belirlendi. 13 serotipi içeren 100 *Salmonella* suşu tespit edildi. Örneklerden en çok izole edilen etken *S. Enteritidis* olarak kaydedildi. İzolatlar belirtilen antibiyotiklere direnç gösterdi; sulfisozaksol (SF) (%79.0), doksisisilin (DOX) (%68.0), tetrasiklin (TE) (%65.0), florfenikol (FFC) (%64.0), ampisillin (AM) (%50.0), gentamisin (GM) (%48.0), trimethoprim-sulfamethoksazol (SXT) (%30.0), spektinomisin (SPT) (%30.0), enrofloksasin (ENR) (%10.0), ofloksasin (NOR) (%10.0), amoksisilin potasyum klavulanat (AC) (%1.0) ve polimiksin (PME) (%4.0). *Salmonella* spp. izolatları cefotaxime (EFT) karşı dirençli değildi. Her bir izolat en az 3 grup antimikrobiyal madde olmak üzere çoklu ilaç direnci gösterdi. PFGE ile elde edilen dört küme ve 38 parmakizi-görüntüsü değişik bölgelerden toplanan suşların arasından belirlendi. Suşlar arası asosiyasyon ve bazı bölgelerde persiste suşların varlığı tespit edildi. İzolatların MLST analizi 10 ST belirlendi. Yedi housekeeping gen farklı varyasyonlara sahipti.

Anahtar sözcükler: *Salmonella* spp., Kanatlı hayvan kesimhane-işleme, Serotiplendirme, PFGE, MLST, Antimikrobiyal duyarlılık



İletişim (Correspondence)



+86 0532 85633936



yffs2000@sina.com

INTRODUCTION

Salmonella spp. is an important group of bacterial zoonotic pathogens which can cause acute food-borne diseases in humans. Each year, approximately 90 million cases of gastroenteritis due to *Salmonella* spp. occur globally [1]. It multiplies mainly in poultry intestinal tract where it can be detected within two hours after infection [2]. A great increase in human food-borne infections caused by *Salmonella* spp. including *Salmonella* Enteritidis and *Salmonella* Typhimurium has been noted in the United States, Europe [3,4]. *Salmonella* spp. infections in humans often result from the ingestion of contaminated foods, such as poultry, beef, pork, eggs and produce. Estimates from the Centers for Disease Control and Prevention (CDC) reported that more than a million people have *Salmonella* spp. poisoning every year from a variety of causes. Poultry products have consistently been identified as important sources of *Salmonella* spp. infection in humans, because vertical transfer of infection from breeding hens to progeny is an important aspect of the epidemiology of *Salmonella* spp. infection within the poultry industry [5]. Other study showed that a great number of *Salmonella* spp. contaminate poultry products and have somewhat different genetic types according to the origin of the integrated broiler operation [6]. Nevertheless, there have been no studies following the dissemination of *Salmonella* spp. in the entire poultry industry in China.

The aim of this study was to estimate the prevalence *Salmonella* spp. contamination in poultry products from 12 different located geographical areas from among the 12 poultry slaughterhouses authorized in China and to characterize all the isolates by serotypes, pulse field gel electrophoresis (PFGE) patterns, multi-locus sequence typing (MLST) patterns and antimicrobial susceptibility to investigate possible sources of infection and to provide information which could help strengthen salmonellosis control programs, and minimize the risk of *Salmonella* spp. exposure in the slaughtering line, which can reduce the contamination pressure downstream at retail shops as well as for end consumers.

MATERIAL and METHODS

The Sample Collection and Isolation

The present study was performed in 12 representative slaughterhouses (A ~ L) in 12 different areas in Shandong province from 2014 to 2015. A total of 960 swabs samples were randomly collected from poultry slaughterhouse-processing, every location the sample number is 80. *Salmonella* spp. contamination samples were collected in five critical steps in slaughterhouse-processing.

All samples were shipped in an icebox to the Microbiology Laboratory of Quality and Safety Risk Assessment

for Animal Products of Ministry of Agriculture (CAHEC) in Qingdao, China, for *Salmonella* spp. isolation within 24 h. The procedure for the detection and isolation of *Salmonella* spp. was carried out according to the techniques recommended by the International Organization for Standardization [7]. All suspicious *Salmonella* colonies with black center were confirmed biochemically using lysine and triple sugar iron agars and API 20E (bioMérieux, France). The *invA* (F: 5'-GAATCCTCAGTTTTCAGTTTC-3'; R: 5'-TAGCCGTAACAACCAATACAAATG-3') gene of *Salmonella* was used to further confirm the identity of the presumptive *Salmonella* spp. [8].

Serotyping

Salmonella-positive isolates were randomly serotyped according to the White-Kauffmann-Le Minor scheme by slide agglutination with O and H antigen specific sera (Sifin Diagnostics, Berlin, Germany) [9].

Pulsed-field Gel Electrophoresis

PFGE was performed for *Salmonella enterica* using the Pulse Net protocol procedures described previously [10]. The bacteria genomic DNA was digested with 50 U of *Xba* I (Takara, China) at 37°C for 3 h. The Pulse Net "Universal" standard strain *Salmonella enterica* serovar Braenderup H9812 was used as a reference marker and *Xba* I (Takara, China) was used as the digestion enzyme. PFGE was repeated twice to determine reproducibility. For untypable isolates, 50 µ Mthiourea (Sigma, USA) was added to the 0.5 XTBE buffer prior to PFGE run as described by Römling and Tümmler [11]. CHEF-Mapper (Bio-Rad, USA) was used for electrophoresis (AutoAlgorithm: 30 kb-low MW, 700 kb-high MW, 19 h). The gel was stained with Gelred (Biofilm) and visualized using a gel imaging system (Bio-Rad, Gel DocXR, USA). PFGE patterns were analyzed using BioNumerics version 5.10; and a dendrogram was constructed using the Dice coefficient and un-weighted pair group methods with the arithmetic mean algorithm (UPGMA). PFGE banding patterns with a similarity index >80% were grouped in the same genotype cluster.

Multi-locus Sequence Typing

The multilocus sequence typing method was performed according to the recommendations of the *Salmonella enterica* MLST website (<http://mlst.warwick.ac.uk/mlst/dbs/Senterica>). Seven housekeeping genes, *aroC*, *dnaN*, *hemD*, *hisD*, *purE*, *sucA* and *thrA*, were amplified using the recommended primers. DNA Taq premix was used with the amplification procedure: 94°C 5 min; followed by 30 cycles: 94°C 1 min, 55°C 30 s, 72°C 1 min; and a final extension at 72°C 10 min. The amplification products were sent for bidirectional sequencing to Takara, China; and the sequences were analyzed using DNASTAR and MEGA4 software. We submitted the sequences to the UCC database for their allele and ST assignments. A minimum

spanning tree based on these STs was generated with BioNumerics version 5.10.

Antimicrobial Susceptibility Tests

Susceptibility of the isolates to antimicrobial agents was evaluated according to Clinical Laboratory Standards Institute (CLSI, 2012) guidelines using the disc diffusion method on Mueller-Hinton agar (Becton Dickinson, USA) [12]. Following 13 antimicrobial agents were tested, including chloramphenicol (florfenicol), penicillins (ampicillin and amoxicillin potassium clavulanat), sulfonamides (sulfisoxazole and trimethoprim-sulfamethoxazole), cepheims (cefotaxime), aminoglycosides (gentamicin and spectinomycin), tetracyclines (tetracycline and doxycycline), fluoroquinolones (enrofloxacin and ofloxacin), and other (polymyxin). Results were interpreted using the Clinical and Laboratory Standards Institute (CLSI, 2012) breakpoints when available. *E. coli* ATCC 25922 was used as quality control [13].

RESULTS

The prevalence of *Salmonella* spp. from each of the 12 poultry slaughterhouses is presented in Table 1. In this study, 100 (10.42%) *Salmonella* spp. was isolated from poultry slaughterhouses, the 12 (12.0%) *Salmonella* isolated

from duck slaughterhouses and the 88 (88.0%) isolated from chicken slaughterhouses. All of the samples from 2 duck slaughterhouses were contaminated with *Salmonella* spp.. The major prevalent serotypes in the duck slaughterhouses were *Salmonella* Thompson, and 5 different serotypes (*Salmonella* Agona, *Salmonella* Montevideo, *S. Typhimurium*, *Salmonella* Indiana and *Salmonella* Albany). In the 10 chicken slaughterhouses, the major of *Salmonella* spp. serotypes were *S. Enteritidis* and *S. Indiana*. Other different serotypes, also were found in chicken slaughterhouses (Fig.1). The characteristics of 100 *Salmonella* spp. strains comprising thirteen majority serotypes were identified. *S. Enteritidis* was the most frequently isolated serotype from samples (43.0%, 43 of 100) (Fig.1). *Salmonella* group *S. Enteritidis* (n = 43), *S. Indiana* (n = 14), *Salmonella* Derby (n = 11), *Salmonella* Infantis (n = 8), *S. Agona* (n = 8), *S. Salmonella* Agona, *Salmonella* Montevideo, *S. Typhimurium*, *Salmonella* Indiana and *Salmonella* Albany (n = 7), *Salmonella* Essen (n = 2), *S. Typhimurium* (n = 2), *S. Albany* (n = 1), *Salmonella* Othmarschen (n = 1), *Salmonella* Schwarzengrund (n = 1), *Salmonella* Abortus equi (n = 1) and *S. Montevideo* (n = 1) were identified during this study (Fig.1).

The results of the antimicrobial susceptibility analysis of the *Salmonella* spp. isolates were summarized in Fig. 2. All 100 *Salmonella* spp. isolates were observed to be susceptible to the 13 antimicrobial agents tested in this




Fig 1. The distinction of serotype of 100 *Salmonella* spp. Isolate

Şekil 1. 100 adet *Salmonella* spp. izolatının serotiplendirmesi

Fig 2. The drug resistance rate of all isolates for 13 kind of antimicrobial agents. Statistical test was only performed in all isolated. Antibiotics abbreviations are: SF: Sulfisoxazole, DOX: Doxycycline, TE: Tetracycline, FFC: Florfenicol, AM: Ampicillin, GM: Gentamicin, SPT; Spectinomycin, SXT: Trimethoprim-sulfamethoxazole, ENR: Enrofloxacin, NOR: Ofloxacin, PME: Polymyxin, EFT: Cefotaxime, AC: Amoxicillin potassium clavulanat

Şekil 2. Tüm izolatların 13 antimikrobiyal madde için ilaç direnci oranları. İstatistiksel test bütün izolatlar için uygulandı. Antibiyotiklerin kısaltmaları; SF: Sulfisozaksol, DOX: Doksisisilin, TE: Tetrasiklin, FFC: Florfenikol, AM: Ampisillin, GM: Gentamisin, SPT; Spektinomisin, SXT: Trimethoprim-sulfamethoksazol, ENR: Enrofloksasin, NOR: Ofloksasin, PME: Polimiksin, EFT: Sefotaksim, AC: Amoksisilin potasyum klavulanat




Table 1. Origin and characterization of *Salmonella* isolated from twelve poultry slaughterhouse**Tablo 1.** Oniki kanatlı hayvan kesimhanesinden izole edilen *Salmonellaların* orjin ve karakterizasyonu

Number of Isolates (n)	Serotype	Location	Animal	MLST Pattern	PFGE Pattern	Prevalence (%)		
A(6)	Enteritidis(3)	A	Chicken	ST-11	0031	7.5		
	Enteritidis(3)	A	Chicken	ST-11	0032			
B (6)	Agona(3)	B	Chicken	ST-13	0003	7.5		
	Indiana(1)	B	Chicken	ST-17	0005			
	Indiana(1)	B	Chicken	ST-17	0017			
	Enteritidis(1)	B	Chicken	ST-11	0031			
C(19)	Abortusequi(1)	C	Chicken	ND	0018	23.8		
	Schwarzengrund(1)	C	Chicken	ND	0024			
	Enteritidis(2)	C	Chicken	ST-11	0031			
	Othmarschen(1)	C	Chicken	ND	0031			
	Enteritidis(1)	C	Chicken	ST-11	0032			
	Derby(8)	C	Chicken	ST-40	0037			
	Essen(2)	C	Chicken	ND	0031			
	Derby(3)	C	Chicken	ND	0024			
D(9)	Indiana(2)	D	Chicken	ST-17	0004	11.3		
	Indiana(1)	D	Chicken	ST-17	0005			
	Indiana(1)	D	Chicken	ST-17	0014			
	Indiana(2)	D	Chicken	ST-17	0025			
	Enteritidis(1)	D	Chicken	ST-17	0028			
	Indiana(2)	D	Chicken	ST-17	0030			
E(13)	Enteritidis(1)	E	Chicken	ST-11	0008	16.3		
	Enteritidis(1)	E	Chicken	ST-11	0009			
	Enteritidis(1)	E	Chicken	ST-17	0010			
	Enteritidis(1)	E	Chicken	ST-11	0011			
	Indiana(1)	E	Chicken	ST-17	0012			
	Agona(1)	E	Chicken	ST-13	0012			
	Enteritidis(1)	E	Chicken	ST-11	0013			
	Agona(1)	E	Chicken	ST-13	0015			
	Agona(1)	E	Chicken	ST-13	0016			
	Indiana(1)	E	Chicken	ST-17	0028			
	Enteritidis(3)	E	Chicken	ST-11	0031			
	F(7)	Agona(1)	F	Chicken	ST-13		0001	8.8
		Infantis(2)	F	Chicken	ST-32		0022	
Indiana(1)		F	Chicken	ST-17	0031			
Enteritidis(2)		F	Chicken	ST-11	0033			
Indiana(1)		F	Chicken	ST-17	0027			
G(9)	Typhimurium(1)	G	Chicken	ST-34	0006	11.3		
	Infantis(1)	G	Chicken	ST-32	0021			
	Infantis(1)	G	Chicken	ST-32	0023			
	Enteritidis(3)	G	Chicken	ST-11	0032			
	Enteritidis(2)	G	Chicken	ST-11	0034			
	Enteritidis(1)	G	Chicken	ST-11	0036			
H(7)	Infantis(1)	H	Chicken	ST-32	0021	8.8		
	Infantis(2)	H	Chicken	ST-32	0022			
	Enteritidis(1)	H	Chicken	ST-11	0029			
	Enteritidis(3)	H	Chicken	ST-11	0032			
I(4)	Agona(1)	I	Duck	ST-13	0003	5.0		
	Montevideo(1)	I	Duck	ST-305	0007			
	Typhimurium(1)	I	Duck	ST-19	0019			
	Indiana(1)	I	Duck	ST-17	0020			
J(6)	Enteritidis(2)	J	Chicken	ST-11	0026	7.5		
	Enteritidis(3)	J	Chicken	ST-11	0031			
	Enteritidis(1)	J	Chicken	ST-11	0032			
K(6)	Enteritidis(5)	K	Chicken	ST-11	0033	7.5		
	Enteritidis(1)	K	Chicken	ND	0035			
L(8)	Thompson(7)	L	Duck	ST-26	0002	10.0		
	Albany(1)	L	Duck	ST-45	0038			

Table 2. Sequence types (STs) and allele profile of each isolate**Table 2.** Herbir izolatin sekans tipi ve allel profili

STs	<i>acroC</i>	<i>dnaN</i>	<i>hemD</i>	<i>hisD</i>	<i>purE</i>	<i>SucA</i>	<i>thrE</i>	Number of Isolates
11	5	2	3	7	6	6	11	41
13	3	3	7	3	3	3	7	8
17	8	8	11	11	5	11	15	16
19	10	7	12	9	5	9	2	1
26	14	13	18	12	5	18	1	7
32	17	18	22	17	5	21	19	7
34	10	19	12	9	5	9	2	1
40	19	20	3	20	5	22	22	8
45	104	100	54	78	104	1	48	1
305	43	41	16	42	34	13	23	1

study. The isolates were resistant to sulfisoxazole (SF) (79.0%), doxycycline (DOX) (68.0%), tetracycline (TE) (65.0%), florfenicol (FFC) (64.0%), ampicillin (AM) (50.0%), gentamicin (GM) (48.0%), trimethoprim-sulfamethoxazole (SXT) (30.0%), spectinomycin (SPT) (30.0%), enrofloxacin (ENR) (10.0%), ofloxacin (NOR) (10.0%), amoxicillin potassium clavulanat (AC) (1.0%) and polymyxin (PME) (4.0%). The *Salmonella* spp. isolates were not resistant to cefotaxime (EFT). Each isolates were multi-drug resistant as they were resistant to at least 2 groups of antimicrobials.

The 100 *Salmonella* spp. isolates were analyzed by PFGE using enzymes *XbaI* resulted in 38 distinguishable patterns demonstrating a high level of genetic diversity among the isolates. An UPGMA dendrogram was constructed (Fig. 3), and described in Table 1. Ten *Salmonella* spp. isolates were untypeable by PFGE. After the addition of thiourea (Sigma, USA) to the running buffer, all isolates remained typeable. The 100 isolates could be divided into four clusters. All serotypes were divided into groups based on their PFGE patterns. In this study, the most common pattern was 0031, which included 13 strains of *S. Enteritidis*, followed by 0032 which was composed of 11 strains of *S. Enteritidis*, and by 0037 which was composed of 8 strains of *S. Derby*. The origins and characteristics of *Salmonella* spp. Isolates identified in this study are outlined in (Table 1). Looking at the strains in more detail, most pattern were within a single serotype and single source except 0003-pattern, 0005-pattern, 0021-pattern, 0022-pattern, 0031-pattern, 0032-pattern and 0033-pattern. The 0031-pattern was found to be composed of 13 *S. Enteritidis*, all recovered from various processing steps at six slaughterhouse, which were A(3), B(1), C(2), E(3), F(1), and J(3); the 0032-pattern was found to be composed of 11 *S. Enteritidis*, which were from H(3), G(3), A(3), J(1) and C(1) five slaughterhouse. The results showed that different places had the same isolates, these strains maybe come from the same farms, or the slaughterhouse circulation caused by cross contamination.

Ten discrete STs were identified among the 91 *Salmonella*

spp. isolates, indicating a degree of genotypic diversity, other 9 isolates were not done. Of these 10 STs, 4 were represented by single isolates, 6 were represented by more than one isolate ($n = 7$ to 41). The predominant STs were ST11, ST17, ST13 and ST26 containing 41 (45.1%), 16 (17.6%), 8 (8.8%) and 8 (8.8%) isolates respectively (Table 2). Isolates characterized as the same STs did not necessarily have the same PFGE pattern. For example, the 41 isolates characterized as ST11 had 11 distinct PFGE patterns (Fig. 3). All of the isolates that shared a PFGE pattern had the same STs. Meanwhile, isolates that shared a PFGE pattern had the same STs (Fig. 3). Showing the detail of the strains in more detail, most STs were within a single serotype.

DISCUSSION

Poultry products have consistently been identified as important sources of *Salmonella* infection in humans, because vertical transfer of infection from breeding hens to progeny is an important aspect of the epidemiology of *Salmonella* spp. infection within the poultry industry. In many countries all over the world including USA, Europe, Korea and China, a wide range of different *Salmonella* serotypes have been found to contaminate the broiler houses, flocks and carcasses of the poultry industry [14]. In this study, 10.4% of the carcasses sampled from 12 poultry slaughterhouses were contaminated with *Salmonella* spp. and showed a lower prevalence than poultry carcasses originating from other countries like Korea (42.1%), Spain (17.9%), Canada (21.2%) and Ireland (26.4%) [15-18]. Although different sampling procedures, sample sizes and bacterial isolation and identification methods could affect the prevalences of *Salmonella* spp., this elevated level of contamination indicates a potential breakdown of hygiene at various stages at poultry farms and processing plants [19]. *S. Enteritidis*, which is responsible for most *Salmonella* infections in humans, is the major serotypes in this study, and this result supports that contaminated carcasses are the major source of infection in human salmonellosis [20].




Fig 3. Dendrogram of PFGE profiles

PFGE patterns and the corresponding dendrogram for 100 isolates obtained in the present study are depicted. The 4PFGE clusters were marked on the node as I to IV. The different clusters observed are designated on the left side of the figure. Displayed on the right hand side are number, serotype, PFGE-Pattern, sequence types (STs), ST complex and source (number)

Şekil 3. PFGR profillerinin dendrogramı

Çalışmadaki PFGE görüntüleri ve ilgili 100 izolatin dendogramı. 4PFGE kümeleri I'den IV'e kadar işaretlenmiştir. Elde edilen farklı kümeler şeklin sol tarafında belirtilmiştir. Sağ tarafta numara, serotip, PFGR-görüntüsü, sekans tipleri (STs), ST kompleks ve kaynakları (numara)

Each isolates showed antimicrobial resistance to at least one antimicrobial agent. The rate of resistance of *Salmonella* spp. to antimicrobial agents was higher than that previously observed^[21,22], but sulfisoxazole, doxycycline and tetracycline were showed the higher resistance in 13 antimicrobial agents. The most of the isolates displayed a high level of susceptibility to most of the antimicrobial agents tested in this study, the antimicrobial susceptibility must will present a challenge for veterinary medicine and farm animal husbandry, and could also pose a threat to public health. Fig. 2 shows *Salmonella* spp. resistance pattern of ofloxacin (NOR) at 10% is a reflection that it is least resistant to the organism, whereas the fluoroquinolones generally have been previously known to have a good sensitivity to *Salmonella* spp.. 100% resistance of *Salmonella* spp. from poultry sources to amoxicillin potassium clavulanat is in line with previous works^[23], even though amoxicillin potassium clavulanat has been known to be of variable resistance. Tetracycline have repeatedly shown high levels of resistance of 60%-65% to which the work was lowed^[24].

Some previous studies showed that the PFGE patterns of *S. Enteritidis* in China are similar even when the isolates have different phage types. The present investigation also confirmed that 18 *S. Enteritidis* isolates with different MLST pattern displayed different PFGE patterns in agreement with previous reports^[25,26]. The PFGE results showed that different places had the same isolates, these strains maybe come from the same farms, or the slaughterhouse circulation caused by cross contamination.

This study shows that the domestic serotypes are *S. Enteritidis* and *S. Indian* in chicken and duck slaughterhouses in Shandong province, and provides detailed information about *Salmonella* spp. isolates from poultry. In addition, the appearance of resistant *Salmonella* spp. isolates from poultry suggests the need for a more prudent use of antibiotics and the importance of controlling this pathogen in poultry products. As human salmonellosis has been repeatedly related to the consumption of products worldwide, continuous research must be conducted to minimize the *Salmonella* spp. contamination in poultry slaughterhouse.

ACKNOWLEDGMENTS

We thank Xianxian Liu, Lu Sun, Yuqing Li, Sijun Zhao and Xiaoliang Sun for their help.

REFERENCES

1. Majowicz SE, Musto J, Scallan E, Angulo FJ, Kirk M, O'Brien SJ, Jones TF, Fazil A, Hoekstra RM: The global burden of nontyphoidal *Salmonella* gastroenteritis. *Clin Infect Dis*, 50, 882-889, 2010. DOI: 10.1086/650733
2. Rostagno MH, Callaway TR: Pre-harvest risk factors for *Salmonella* Enterica in pork production. *Food Res Int*, 45, 634-640, 2012. DOI: 10.1016/j.foodres.2011.04.041
3. Brooks JT, Matyas BT, Fontana J, DeGroot MA, Beuchat LR, Hoekstra M, Friedman CR: An outbreak of *Salmonella* serotype Typhimurium infections with an unusually long incubation period. *Foodborne Pathog Dis*, 9, 245-248, 2012. DOI: 10.1089/fpd.2011.0992
4. Janmohamed K, Zenner D, Little C, Lane C, Wain J, Charlett A, Adak B, Morgan D: National outbreak of *Salmonella* Enteritidis phage type 14b in England, September to December 2009: Case-control study. *Euro Surveill*, 16, Pii: 19840, 2011.
5. Gast RK: Understanding *Salmonella* Enteritidis in laying chickens: the contributions of experimental infections. *Int J Food Microbiol*, 21, 107-116, 1994. DOI: 10.1016/0168-1605(94)90204-6
6. Kim A, Lee YJ, Kang MS, Kwag SI, Cho JK: Dissemination and tracking of *Salmonella* spp. in integrated broiler operation. *J Vet Sci*, 8, 155-161, 2007. DOI: 10.4142/jvs.2007.8.2.155
7. International Organization for Standardization (ISO): Microbiology of Food and Animal Feeding stuffs-Horizontal Method for the Detection of *Salmonella* spp. International Organization for Standardization: Geneva, Switzerland, 2007.
8. Kim JS, Lee GG, Park JS, Jung YH, Kwak HS, Kim SB, Nam YS, Kwon ST: A novel multiplex PCR assay for rapid and simultaneous detection of five pathogenic bacteria: *Escherichia coli* O157: H7, *Salmonella*, *Staphylococcus aureus*, *Listeria monocytogenes*, and *Vibrio parahaemolyticus*. *J Food Prot*, 70, 1656-1662, 2007.
9. Grimont PAD, Weill FX: Antigenic formulae of the *Salmonella* Serovars. Paris: World Health Organization, Institut Pasteur, 166p, 2007.
10. Wu W, Wang H, Lu J, Wu J, Chen M, Xu Y, Lu Y: Genetic diversity of *Salmonella* enterica serovar Typhi and Paratyphi in Shenzhen, China from 2002 through 2007. *BMC Microbiol*, 10, 32, 2010. DOI: 10.1186/1471-2180-10-32
11. Römling U, Tümmler B: Achieving 100% typeability of *Pseudomonas aeruginosa* by pulsed-field gel electrophoresis. *J Clin Microbiol*, 38, 464-465, 2000.
12. CLSI: Performance Standards for Antimicrobial Susceptibility Testing, Twenty-Second Informational ament Wayne Pennsylvania: Clinical and Laboratory Standards Institute; 2012.
13. Gai W, Wang J, Wang J, Cui Z, Qu Z, Cui J, Du X, Huang X, Zhao J: Molecular classification and drug resistance analysis of *Escherichia coli* isolated from poultry in China. *Int J Clin Exp Med*, 8, 836-844, 2015.
14. Song YK, Shin DH, Byun JW, Jeon KY, Jung BY: Seroprevalence of *Salmonella* Pullorum and Gallinarum in grandparent poultry stock farms during the 2006-2007. *Kor J Vet Publ Hlth*, 33, 131-136, 2009.
15. Bae DH, Dessie HK, Baek HJ, Kim SG, Lee HS, Lee YJ: Prevalence and characteristics of *Salmonella* spp. isolated from poultry slaughterhouses in Korea. *Vet Med Sci*, 75, 1193-1200, 2013.
16. Arsenault J, Letellier A, Quessy S, Boulianne M: Prevalence and risk factors for *Salmonella* and *Campylobacter* spp. carcass contamination in broiler chickens slaughtered in Quebec, Canada. *J Food Prot*, 70, 1820-1828, 2007.
17. Capita R, Alonso-Calleja C, Prieto M: Prevalence of *Salmonella* enterica serovars and genovars from chicken carcasses in slaughterhouses in Spain. *J Appl Microbiol*, 103, 1366-1375, 2007.
18. Duffy G, Cloak OM, O'Sullivan MG, Guillet A, Sheridan JJ, Blair IS, McDowell DA: The incidence and antibiotic resistance profiles of *Salmonella* spp. on Irish retail meat products. *Food Microbiol*, 16, 623-631, 1999. DOI: 10.1006/fmic.1999.0278
19. Jørgensen F, Bailey R, Williams S, Henderson P, Wareing DR, Bolton FJ, Frost JA, Ward L, Humphrey TJ: Prevalence and numbers of *Salmonella* and *Campylobacter* spp. on raw, whole chickens in relation to sampling methods. *Int J Food Microbiol*, 76, 151-164, 2002. DOI: 10.1016/S0168-1605(02)00027-2
20. Cheong HJ, Lee YJ, Hwang IS, Kee SY, Cheong HW, Song JY, Kim JM, Park YH, Jung JH, Kim WJ: Characteristics of non-typhoidal *Salmonella* isolates from human and broiler-chickens in southwestern Seoul, Korea. *J Korean Med Sci*, 22, 773-778, 2007.
21. Kuang X, Hao H, Dai M, Wang Y, Ahmad I, Liu Z, Zonghui Y: Serotypes and antimicrobial susceptibility of *Salmonella* spp. isolated

from farm animals in China. *Front Microbiol*, 6, 602, 2015. DOI: 10.3389/fmicb.2015.00602

22. Morar A, Sala C, Imre K: Occurrence and antimicrobial susceptibility of *Salmonella* isolates recovered from the pig slaughter process in Romania. *J Infect Dev Ctries*, 9, 99-104, 2015. DOI: 10.3855/jidc.5236

23. Ezekiel CN, Olanmoye AO, Oyinloye JMA, Olaoye OB, Edun AO: Distribution, antibiogram and multidrug resistance in *Enterobacteriaceae* from commercial poultry feeds in Nigeria. *Afr J Microbiol Res*, 5, 294-301, 2011.

24. Sakaridis I, Soutos N, Iossifidou E, Koidis P, Ambrosiadis I: Prevalence and antimicrobial resistance of *Salmonella* serovars from chicken

carcasses in northern Greece. *J Food Saf*, 2011. DOI: 10.1111/j.1745-4565.2010.00286.x

25. Bai L, Lan R, Zhang X, Cui S, Xu J, Guo Y, Li F, Zhang D: Prevalence of *Salmonella* isolates from chicken and pig slaughterhouses and emergence of ciprofloxacin and cefotaxime co-resistant *S. enterica* serovar Indiana in Henan, China. *PLoS One*, 10, e0144532, 2015. DOI: 10.1371/journal.pone.0144532

26. Wang Y, Liu C, Zhang Z, Hu Y, Cao C, Wang X, Xi M, Xia X, Yang B, Meng J: Distribution and molecular characterization of *Salmonella enterica* hypermutators in retail food in China. *J Food Prot*, 78, 1481-1487, 2015. DOI: 10.4315/0362-028X.JFP-14-462