

RESEARCH ARTICLE

Host Preferences of Vector *Culicoides* (Diptera, Ceratopogonidae, *Culicoides* Latreille) Species in Türkiye

Ahmet DENİZ¹ (*)  Çağla KORKMAZ²  Hilal BEDİR¹  Zati VATANSEVER³ 
Aysel İTİK EKİNCİ⁴  Osman SEZER⁵  Ufuk EROL⁶  Selim TUNCER⁷ 

¹ Department of Parasitology, Faculty of Medicine, Kafkas University, TR-36100 Kars - TÜRKİYE

² Veterinary Control Central Research Institute, Department of Parasitology, TR-06020 Ankara - TÜRKİYE

³ Department of Parasitology, Faculty of Veterinary Medicine, Kafkas University, TR-36100 Kars - TÜRKİYE

⁴ Elazığ Veterinary Control Institute, Department of Parasitology, TR-23200 Elazığ - TÜRKİYE

⁵ Adana Veterinary Control Institute, Department of Parasitology, TR-01170 Adana - TÜRKİYE

⁶ Department of Parasitology, Faculty of Veterinary Medicine, Cumhuriyet University, TR-58140 Sivas - TÜRKİYE

⁷ Ankara Food Control Laboratory, Molecular Biology Laboratory Unit, TR-06790 Ankara - TÜRKİYE

ORCID: A.D. 0000-0003-4561-675X, C.K. 0000-0002-4308-4035, H.B. 0000-0002-1583-3385, Z.V. 0000-0003-3460-3849, A.İ.E. 0000-0002-2730-7657, O.S. 0000-0002-4938-9761, U.E. 0000-0002-6766-1335, S.T. 0000-0002-6753-4379

Article ID: KVFD-2022-28539 Received: 26.09.2022 Accepted: 14.01.2023 Published Online: 27.01.2023

Abstract: Determination of host preferences of blood-feeding arthropods is an important criterion for how ectoparasite-host interactions, host selection, and feeding behaviour affects pathogen transmission. This study aimed to examine the host's blood from engorged *Culicoides* species collected from cattle farms in Turkey to shed light on their host preferences. For this purpose, we investigated the blood of 5 different hosts (cattle, sheep, dog, horse, and human) in 7 *Culicoides* species/species complexes by multiplex-PCR analysis, considering the domestic animals on the farms. Engorged *Culicoides* specimens were collected from places in Turkey with different geographical and ecological characteristics. A total of 1225 female *Culicoides* belonging to 7 different species/species complexes namely *C. obsoletus* complex (*C. obsoletus* and *C. scoticus*) (n:450), *C. schultzei* complex (n:234), *C. imicola* (n:208), *C. punctatus* (n:162), *C. newsteadi* (n:144), *C. lupicaris* (n:24), and *C. pulicaris* (n:3) were analysed for host blood identification. Abdomens of the engorged midges were separated from their body and pooled according to the date of collection and species. A total of 69 pools consisting of 1-28 specimens were analysed by multiplex-PCR and only cattle blood was detected in all pools. This study presents the first data on the identification of host preference of some *Culicoides* species in Turkey.

Keywords: Blood meal preferences, *Culicoides*, multiplex-PCR, Türkiye.

Türkiye'de Vektör *Culicoides* (Diptera, Ceratopogonidae, *Culicoides* Latreille) Türlerinin Konak Tercihlerinin Belirlenmesi

Öz: Kanla beslenen artropodların konak tercihlerinin tanımlanması, ektoparazit-konak etkileşimleri, eklembacaklıların konak seçimi ve beslenme davranışının patojen bulaşmasını nasıl etkilediği konularında önemli bir kriterdir. Bu çalışmanın amacı, Türkiye'deki sığır çiftliklerinden toplanan doymuş *Culicoides* türlerinden konakçı kanını inceleyerek konak tercihlerine ışık tutmaktır. Bu amaçla çiftliklerde bulunan evcil hayvanlar göz önünde bulundurularak 7 *Culicoides* tür/tür kompleksinde 5 farklı konağın (sığır, koyun, köpek, at ve insan) kanını multiplex-PCR testi ile araştırdık. Doymuş *Culicoides* örnekleri Türkiye'de farklı coğrafi ve ekolojik özelliklere sahip yerlerden toplanmıştır. *Culicoides obsoletus* complex (including *C. obsoletus* and *C. scoticus*) (n:450), *C. schultzei* complex (n:234), *C. imicola* (n:208), *C. punctatus* (n:162), *C. newsteadi* (n:144), *C. lupicaris* (n:24) ve *C. pulicaris* (n:3) olarak belirlenen 7 farklı tür/tür kompleksine ait toplam 1225 adet *Culicoides* konak kanı tespiti için analiz edildi. Doymuş dişilere ait abdomenler gövdelerinden ayrılarak toplanma tarihlerine ve türlerine göre havuzlar oluşturuldu. Toplam 1-28 sinekten oluşan 69 havuz multiplex-PCR ile analiz edildi ve tüm örneklerde sadece sığır kanı tespit edildi. Çalışmamız Türkiye'de *Culicoides* türlerinin konak tercihinin belirlenmesi konusunda ilk verileri sunmaktadır.

Anahtar sözcükler: Kan emme tercihi, *Culicoides*, multiplex-PCR, Türkiye.

How to cite this article?

Deniz A, Korkmaz C, Bedir H, Vatansever Z, İtik Ekinci A, Sezer O, Erol U, Tuncer S: Host preferences of vector *Culicoides* (Diptera, Ceratopogonidae, *Culicoides* latreille) species in Türkiye. *Kafkas Univ Vet Fak Derg*, 29 (1): 63-69, 2023.
DOI: 10.9775/kvfd.2022.28539

(*) Corresponding Author

Cellular phone: +90 532 7454171

E-mail: ametd@hotmail.com (A. Deniz)



This article is licensed under a Creative Commons Attribution-NonCommercial 4.0 International License (CC BY-NC 4.0)

INTRODUCTION

Culicoides (Diptera: Ceratopogonidae) are among the smallest hematophagous flies in the world, measuring 1-3 mm in size. The genus is of great interest for its role as biological vector of pathogens of medical and veterinary importance. In addition to several nematode and protozoan species, more than 50 arboviruses have been isolated from *Culicoides* species^[1-3]. Their role in transmitting diseases such as bluetongue virus (BTV), epizootic haemorrhagic disease virus (EHDV), African horse sickness virus (AHSV), Akabane virus, and Schmallenberg virus which causes economic losses in the livestock industry is the main reason for large scale investigation of this genus^[4,5]. They are found on nearly all major landmasses from sea level to 4000 m^[3]. A total number of 1347 species have been described so far^[6]; 72 species have been reported from Turkey^[7,8].

Culicoides species can cause severe itchy skin reactions in humans and animals, forcing the hosts to relocate or to flee indoors^[1]. Their ability to transmit the Oropouche virus (OROV), the etiological agent of the febrile illness Oropouche fever, among humans has been the most important role of *Culicoides* species in public health to date^[3]. A wide variety of insights can be obtained by examining the blood on which arthropods feed, such as the evolution of host specificity between vertebrates and their ectoparasites, ectoparasite-host interactions, the ecology of infectious diseases, how arthropod host selection and feeding behaviour affect pathogen transmission, and the economic and wellness effects of ectoparasite infestation on domestic animals and wildlife^[9].

The vast majority of *Culicoides* species feed on mammals and birds; however, reptiles and amphibians are also in the host spectrum of these flies^[1,10-12]. Although feeding frequency varies by species and environmental conditions, host availability plays an important role in the feeding behaviour of biting midges in general. Most of the *Culicoides* species are mammalophilic and ornithophilic, although some species are known to feed on reptiles and frogs^[1,13-16]. As a result of the molecular identification of vertebrate hosts of *Culicoides* species in Europe, 45 different host species, including 33 bird species and 12 mammal species, were identified^[17].

While serological methods were used to determine host preferences in the past^[18,19], these have now been replaced by molecular methods. Advances in polymerase chain reaction (PCR)-based assays and molecular techniques for blood-meal analysis using direct sequencing of the cytochrome b (*cytb*) gene allow for species-level identification of hosts with a much higher degree of accuracy than has been achieved with previous serological assays^[9-12,20-26]. This study aimed to identify the host's blood in engorged *Culicoides* species collected from cattle

farms with other domesticated animals to understand their feeding preferences. This is the first study in Turkey to analyse the host's blood based on a segment of the mitochondrial gene, *cytb*, in *Culicoides* species.

MATERIAL AND METHODS

Study Area

The stations where *Culicoides* specimens were collected were selected from places in Turkey with different geographical and ecological characteristics (*Table 1*).

Collection of *Culicoides* Specimens

In this study, Onderstepoort Veterinary Institute (OVI) type light traps working with 220V, 8 Watt black fluorescent light and downward fan with a 4 mm net around to prevent the entry of large arthropods were used. Light traps were placed approximately 1.5-2 m above ground out of animals' reach and operated one hour before sunset until one hour after sunrise in cattle farms, where there were at least 20 cattle. There were few sheep, dogs, and a few horses in and around these farms. The samples were obtained from 24 cattle farms in 20 provinces by running traps once a month for 14 months between April-2019 and October-2020 (*Table 1*). Collected *Culicoides* specimens were morphologically identified under the stereomicroscope using the identification key for Palaearctic *Culicoides* species^[27]. *Culicoides schultzei* complex consists of several species (*C. oxystoma*, *C. schultzei*, *C. subschultzei*, *C. kingi*, *C. rhizophorensis*, *C. enderleini*, *C. nevillei*, and *C. neoschultzei*)^[28]. Since there is no detailed identification key on the complex, specimens were identified as *C. schultzei* complex. In this study, the abdomens of engorged fully females were separated and pooled according to their species and date of collection, DNA extraction was performed and the DNA samples were stored at -20°C until use.

DNA Extraction and Validation with Blood Samples

Blood of from domestic hosts (cattle, sheep, dog, and horse) was obtained from the animal hospital of Kafkas University Faculty of Veterinary Medicine and human blood obtained from the University Research Hospital. DNA extraction was conducted from each blood sample by using QIAmp DNA Mini Kit (Qiagen, GmbH, Hilden, Germany) in order to obtain positive controls in the PCR assay. These hosts were chosen for the assay because they were potential targets where traps were placed. Abdomens of engorged *Culicoides* females were separated from their bodies and pooled (1-28 per tube) in separate tubes according to date and species. Each pool was digested in 200 mL lysis buffer in tubes with steel balls and extracted according to the kit procedure (Analytik Jena AG, AJ Innuscreen GmbH, Berlin, Germany).

Table 1. Summary of locations where blood-fed *Culicoides* were collected

Province	District/Village	Species	Latitude	Longitude	Altitude (m)	Type of Animals Around the Trap	Blood-Fed Midges, n	Trap Localisations
Artvin	Ardanuç	<i>C. newsteadi</i>	41.114820	42.066060	489	Cattle, dog	40	Indoor
		<i>C. obsoletus</i> complex					217	
		<i>C. punctatus</i>					121	
		<i>C. lupicaris</i>					23	
Ankara	Çubuk	<i>C. newsteadi</i>	40.118740	32.944540	956	Cattle, sheep, dog	2	Outdoor
Edirne	Merkez/ Budakdoğanca	<i>C. newsteadi</i>	41.760760	26.340980	116	Cattle, dog, horse	71	Outdoor
		<i>C. obsoletus</i> complex					5	
		<i>C. punctatus</i>					13	
		<i>C. pulicaris</i>					3	
Edirne	Merkez/İskender	<i>C. newsteadi</i>	41.630751	26.669324	93	Cattle, dog, horse	21	Indoor
		<i>C. punctatus</i>					4	
Antalya	Kaş	<i>C. newsteadi</i>	36.339800	29.327030	11	Cattle, dog	4	Outdoor
Erzincan	Merkez/Çatalören	<i>C. newsteadi</i>	39.664770	39.518180	1191	Cattle, dog	6	Outdoor
Trabzon	Sürmene	<i>C. obsoletus</i> complex	40.891111	40.056944	75	Cattle, dog	32	Indoor
Rize	Findıklı	<i>C. obsoletus</i> complex	41.253880	41.156770	30	Cattle, dog	137	Indoor
Mersin	Anamur	<i>C. obsoletus</i> complex	36.133480	32.859470	54	Cattle, sheep, dog	8	Indoor
		<i>C. imicola</i>					84	
Mersin	Silifke	<i>C. schultzei</i> complex	36.335278	34.000833	0,3	Cattle, dog	16	Outdoor
Samsun	Atakum	<i>C. obsoletus</i> complex	41.433580	36.084770	204	Cattle, dog	15	Indoor
		<i>C. lupicaris</i>					1	
Bursa	İzmit	<i>C. obsoletus</i> complex	40.541110	29.834200	850	Cattle, sheep, dog	2	Indoor
Kastamonu	Araç	<i>C. obsoletus</i> complex	41.218780	33.381380	852	Cattle, sheep, dog	23	Indoor
Erzurum	Şenkaya	<i>C. obsoletus</i> complex	40.640000	42.338889	1247	Cattle, dog	2	Indoor
Erzurum	Şenkaya/Aydoğdu	<i>C. punctatus</i>	40.701246	42.471634	1631	Cattle, dog	20	Indoor
Kırklareli	Vize	<i>C. obsoletus</i> complex	41.706903	27.704541	430	Cattle, sheep, dog	2	Indoor
Balıkesir	Karasi	<i>C. obsoletus</i> complex	39.894986	27.843145	280	Cattle, sheep, dog	3	Indoor
Giresun	Merkez	<i>C. obsoletus</i> complex	40.910409	38.313453	35	Cattle, dog	4	Indoor
Burdur	Merkez	<i>C. punctatus</i>	37.633690	30.106700	875	Cattle, dog	4	Outdoor
Hatay	Arsuz	<i>C. imicola</i>	36.406389	35.891111	7	Cattle, dog	49	Outdoor
		<i>C. schultzei</i> complex					101	
Diyarbakır	Sur	<i>C. imicola</i>	37.961110	40.428700	641	Cattle, dog	5	Indoor
Diyarbakır	Bismil	<i>C. schultzei</i> complex	37.826090	40.615240	554	Cattle, dog	52	Outdoor
Gaziantep	Nurdağı	<i>C. schultzei</i> complex	37.190680	36.843299	506	Cattle, sheep, dog	50	Outdoor
		<i>C. imicola</i>					70	
Adana	Kozan	<i>C. schultzei</i> complex	37.363219	35.716955	90	Cattle, sheep, dog	15	Outdoor


Species-specific Multiplex PCR Assay

The primer sets used in this study are given in [Table 2](#). Separate reactions were prepared for each primer set ([Fig. 1](#)). Multiplex-PCR was performed with UNREV1025 in a total volume of 25 µL using 2.5 µL of 10X PCR buffer, 2.5 µL of 2 mM MgCl₂, 1 µL of 10 mM dNTP mix, 1 µL of each primer (10 pmol), 0.125 µL Taq polymerase and 4

µL template DNA. PCR conditions were as follows: initial denaturation at 95°C for 5 min, followed by 40 cycles at 95°C for 30 sEC, 57°C for 1min, 72°C for 1min, and final elongation at 72°C for 5min. The PCR products were loaded onto 1.5% agarose gel stained with SYBR® Safe DNA (Thermo Fisher Scientific, Invitrogen) and visualized under UV light.

Table 2. Primer set used for the identification of blood-meal in *Culicoides abdomens*

Primer Name	Primer Sequence (5'→3')	Length (bp)	Origin
UNREV1025	GGTTGTCTCCAATTCATGTTA		Kent and Norris [29]
Human741	GGCTTACTTCTTTCATCTCTCCT	334	Kent and Norris [29]
Cow121	CATCGGCACAAATTTAGTCG	561	Kent and Norris [29]
Horse	CCCTACATCGGTACTACCC	500	Pitzer et al. [30]
Forward primer UNIV1	GACCAATGATATGAAAAACCATCGTTGT		Garros et al. [21]
Dog, <i>Canis lupus familiaris</i>	CAAGCATACTCCTAGTAAGGATCCG	170	Garros et al. [21]
Forward primer UNIV2	TGAGGACAAATATCATTYTGAGGRGC		Garros et al. [21]
Sheep <i>Ovis aries</i>	GGCGTGAATAGTACTAGTAGCATGAGGATGA	336	Garros et al. [21]
Cow, <i>Bos taurus</i>	TAAGATGTCCTTAATGGTATAGTAG	287	Garros et al. [21]
Forward primer UNIV3	TTTTTTTTTTTCGVTCHATYCCHAAAYAACTAGG		Garros et al. [21]
Horse, <i>Equus caballus</i>	TACGTATGGGTGTTCCACTGGC	208	Garros et al. [21]



RESULTS

In this study, 1225 engorged *Culicoides* females were identified from 24 traps in 20 provinces, which belong to 7 different *Culicoides* species/species complexes: *C. obsoletus* complex (*C. obsoletus* and *C. scoticus*) (n:450), *C. schultzei* complex (n:234), *C. imicola* (n:208), *C. punctatus* (n:162), *C. newsteadi* (n:144), *C. lupicaris* (n:24) and *C. pulicaris* (n:3). Engorged females of *C. obsoletus* complex were collected from 12 different traps, *C. newsteadi* from 5 different traps, *C. punctatus* from 4 different traps, *C. lupicaris* from 2 different traps, *C. pulicaris* from only 1 trap, *C. imicola* from 4 different traps and *C. schultzei* complex from 5 different traps (Table 1). Engorged abdomens were separated and pooled according to the date of collection and species. A total of 69 pools consisting of 1-28 specimens were analysed. DNA from human, sheep, cattle, dog, and horse blood were amplified with species-specific *cytb* primers (Fig. 1). As a result of the molecular analysis of 69 pools, all 7 *Culicoides* species/ species complexes were found to be fed only from cattle (Table 3).

Table 3. Blood-meal identifications by multiplex-PCR of engorged *Culicoides*.

<i>Culicoides</i> Species	Cow (<i>Bos taurus</i>)	Sheep (<i>Ovis aries</i>)	Horse (<i>Equus caballus</i>)	Human (<i>Homo sapiens</i>)	Dog (<i>Canis lupus</i>)
<i>C. newsteadi</i>	+	-	-	-	-
<i>C. obsoletus</i> complex (<i>C. obsoletus</i> / <i>C. scoticus</i>)	+	-	-	-	-
<i>C. punctatus</i>	+	-	-	-	-
<i>C. lupicaris</i>	+	-	-	-	-
<i>C. pulicaris</i>	+	-	-	-	-
<i>C. imicola</i>	+	-	-	-	-
<i>C. schultzei</i> complex	+	-	-	-	-

DISCUSSION

Molecular techniques developed to identify vertebral hosts that blood-feeding arthropods feed on, allow the host to be determined at the species level^[9,17]. In the identification of host preferences of engorged *Culicoides* species, species-specific primers based on the *cytb* that amplify the known host DNA^[20,21] and general primers that amplify the conserved genetic regions of all vertebrate species are used^[10,22,23,25,26]. In this study, we performed a multiplex-PCR test using specific primers based on *cytb* polymorphism and determined the host blood (bovine, sheep, dog, equine and human) in 7 *Culicoides* species/species complexes. Multiplex-PCR test is a fast, cost-effective, and efficient method that can be routinely applied in the laboratory, where large numbers of samples are tested simultaneously. It is widely used to identify the host preference of various blood-feeding arthropods^[17,21,30]. However, it requires preliminary selection to identify primer sets for targeted hosts, which may cause a significant limitation in samples from undisturbed habitats with a wider host range^[17,21]. We did not encounter the stated adverse effects as there were a limited number of target hosts (cattle, sheep, dog, horse, and human) in and around the farms where the light traps were placed. In this study, as a result of multiplex-PCR, 7 *Culicoides* species/species complexes (*C. obsoletus* complex, *C. schultzei* complex, *C. imicola*, *C. punctatus*, *C. newsteadi*, *C. lupicaris*, and *C. pulicaris*) were found to feed on cattle but not on other hosts. This may be due to the small number of other hosts in the farms, the higher CO₂ emission, the larger mass of cattle and the greater number of cattle in the farms^[20]. It has been stated that the host selection of *Culicoides* species may be limited by the presence of hosts in their environment^[15,17]. Lassen et al.^[12] found 74% cattle blood from the collected samples of *Culicoides* species in their study, and they stated that the result was due to placing the traps close to cattle. In our study, all the traps were placed in cattle farms. As a result, cattle appear to be more attractive hosts for *Culicoides* due to their greater number, less mobility than other animals, and limited defensive movements. In a study on the host preference of *Culicoides* between cattle, horse, and sheep, it was found that *Culicoides* preferred cattle more in an area where cattle and sheep coexist, while more species and individuals were collected from horses^[31]. It has been reported that cattle not only attract *Culicoides* but are also important hosts of for these species^[13]. Studies on host preferences are also performed by direct aspiration of species from hosts, sweep nets or drop nets. Comparison of host preference between cattle and sheep with these methods showed that cattle are far more attractive hosts for *Culicoides* species in general^[13,32]. The structure of the farms (indoor or outdoor), seasonal behavioural changes, and the number of hosts are important for host selection of *Culicoides* species^[12].

Culicoides imicola is one of the most important vector species among the species analysed for host identification in this study^[3]. *C. imicola* is an important vector for bluetongue virus (BTV) and is distributed in the south and southeast regions of Türkiye^[33]. There are few number of researches on the host preference of this species. Slama et al.^[34] identified host preferences of *C. imicola* in 96 engorged females and found that they feed on cattle, humans, sheep, goats, and dogs. Martínez-de la Puente et al.^[23] found in their study that *C. imicola* feed on 6 mammal species. In a study carried out in South Africa, it has been detected that *C. imicola* and *C. subschultzei* fed on human, zebra, and kudu blood^[35]. Both species (*C. imicola* and *C. schultzei* complex) were found to feed on blood from cattle in this study.

Newly engorged female specimens are important in identifying the host source with high accuracy. However, it is known that it is quite difficult to collect these newly engorged females. It represents only a small portion of the total individuals captured^[17]. For this reason, it is stated that host preference identification can also be made with *Culicoides* at different stages of digestion^[36]. In our study, 1225 new engorged *Culicoides* specimens were analysed. In future studies, choosing samples from both newly engorged specimens and specimens at different stages of digestion (partially or completely) will expand the number of samples for host identification studies.

Studies carried out in Turkey so far have mainly focused on the *Culicoides* species fauna^[7,8,37,38]. Our study presents the first data on the identification of host preference of *Culicoides* species in Turkey. In this study, we found that 7 *Culicoides* species/species complexes (*C. obsoletus* complex, *C. schultzei* complex, *C. imicola*, *C. punctatus*, *C. newsteadi*, *C. lupicaris* and *C. pulicaris*) did not feed on other hosts but on cattle by multiplex-PCR assays. Although there were other animals on the farms, the fact that they prefer only cattle can be attributed to the low number of other hosts on the farms. It may also be due to cattle being more attractive hosts for *Culicoides* species found in the farms in this study.

Availability of Data and Materials

The data that support the findings of this study are available from the corresponding author (A. Deniz) on reasonable request.

Funding Support

The authors received no specific funding for this work.

Ethical Approval

Not necessary

Competing Interests

The authors declared that there is no conflict of interest

Author Contributions

The study was designed by AD and CK. Fly samples were collected by AD, CK, HB, AIE and OS. *Culicoides* species identification was made by CK, UE and ST. Multiplex-PCR diagnosis was made by AD, HB and ZV. The original draft was prepared by AD. All authors contributed to the review and proofreading of the article.

REFERENCES

- Borkent A:** The Biting Midges, the Ceratopogonidae (Diptera), In, Marquart WC (Ed): *Biology of Disease Vectors*. 2nd ed., 113-126, Elsevier Academic Press, China, 2004.
- Meiswinkel R, Venter GJ, Nevill EM:** Vectors: *Culicoides* spp. In, Goetzer JAW, Tustin RC (Eds): *Infectious Diseases of Livestock*. 2nd ed., 93-136, Oxford University Press, Oxford, 2004.
- Mellor PS, Boorman J, Baylis M:** *Culicoides* biting midges: Their role as arbovirus vectors. *Annu Rev Entomol*, 45 (1): 307-340, 2000. DOI: 10.1146/annurev.ento.45.1.307
- Miranda MA:** Case studies of vector-borne diseases in livestock: Bluetongue virus. In, Garros C, Bouyer J, Takken W, Smallegange RC (Eds): *Pests and Vector-Borne Diseases in the Livestock Industry Ecology and Control of Vector-Borne Diseases*. 221-271, Wageningen Academic Publishers The Netherlands, 2018.
- Roberts H, Nixon E, Matt C:** Case study: Costs of *Culicoides*-borne arboviral diseases. In, Garros C, Bouyer J, Takken W, Smallegange RC (Eds): *Pests and Vector-Borne Diseases in the Livestock Industry Ecology and Control of Vector-Borne Diseases*. 497-508, Wageningen Academic Publishers The Netherlands, 2018.
- Borkent A, Dominiak P:** Catalog of the Biting Midges of the World (Diptera: Ceratopogonidae), *Zootaxa*, 4787 (1): 1-377, 2020. DOI: 10.11646/zootaxa.4787.1.1
- Korkmaz C, Alten B, Erol U, Deniz A:** Updated checklist of *Culicoides* Latreille (Diptera: Ceratopogonidae) of Turkey with ten new records. *J Vector Ecol*, 46, 124-136, 2021. DOI: 10.52707/1081-1710-46.2.124
- Turgut F:** Morphological characteristics of two *Culicoides* Latreille biting midges (Diptera, Ceratopogonidae) with a new record for Turkish fauna. *Trakya Univ J Nat Sci*, 23 (Special Issue): 3-8, 2022. DOI: 10.23902/trkjnat.1082308
- Kent RJ:** Molecular methods for arthropod bloodmeal identification and applications to ecological and vector-borne disease studies. *Mol Ecol Resour*, 9, 4-18, 2009. DOI: 10.1111/j.1755-0998.2008.02469.x
- England ME, Perce-Kelly P, Brugman VA, King S, Gubbins S, Sach F, Sanders CJ, Masters NJ, Denison E, Carpenter S:** *Culicoides* species composition and molecular identification of host blood meals at two zoos in the UK. *Parasit Vectors*, 13:139, 2020. DOI: 10.1186/s13071-020-04018-0
- González MA, Goiri F, Prosser SWJ, Cevidanes A, Hernández-Triana LM, Barandika JF, Hebert PDN, García-Pérez AL:** *Culicoides* species community composition and feeding preferences in two aquatic ecosystems in northern Spain. *Parasit Vectors*, 15 (1):199, 2022. DOI: 10.1186/s13071-022-05297-5
- Lassen SB, Nielsen SA, Skovgård H, Kristensen M:** Molecular identification of bloodmeals from biting midges (Diptera: Ceratopogonidae: *Culicoides* Latreille) in Denmark. *Parasitol Res*, 108 (4): 823-829, 2011. DOI: 10.1007/s00436-010-2123-4
- Ayllón T, Nijhof AM, Weiher W, Bauer B, Allène X, Clausen PH:** Feeding behaviour of *Culicoides* spp. (Diptera: Ceratopogonidae) on cattle and sheep in northeast Germany. *Parasit Vectors*, 7:34, 2014. DOI: 10.1186/1756-3305-7-34
- Bobeva A, Zehindjiev P, Ilieva M, Dimitrov D, Mathis A, Bensch S:** Host preferences of ornithophilic biting midges of the genus *Culicoides* in the Eastern Balkans, *Med Vet Entomol*, 29 (3): 290-296, 2015. DOI: 10.1111/mve.12108
- Videvall E, Bensch S, Ander M, Chirico J, Sigvald R, Ignell R:** Molecular identification of bloodmeals and species composition in *Culicoides* biting midges. *Med Vet Entomol*, 27 (1): 104-112, 2012. DOI: 10.1111/j.1365-2915.2012.01038.x
- Santiago-Alarcon D, Havelka P, Schaefer HM, Segelbacher G:** Bloodmeal analysis reveals avian infections and broad host preferences of *Culicoides* (Diptera: Ceratopogonidae) vectors, *PLoS One*, 7 (2):e31098, 2012. DOI: 10.1371/journal.pone.0031098
- Martínez-de la Puente J, Figuerola J, Soriguer R:** Fur or feather? Feeding preferences of species of *Culicoides* biting midges in Europe. *Trends Parasitol*, 31, 16-22, 2015. DOI: 10.1016/j.pt.2014.11.002
- Blackwell A, Brown M, Mordue W:** The use of an enhanced ELISA method for the identification of *Culicoides* bloodmeals in host-preference studies. *Med Vet Entomol*, 9, 214-218, 1995. DOI: 10.1111/j.1365-2915.1995.tb00183.x
- Nevill EM, Anderson D:** Host preferences of *Culicoides* midges (Diptera: Ceratopogonidae) in South Africa as determined by precipitin tests and light trap catches. *Onderstepoort J Vet Res*, 39 (3): 147-152, 1972.
- Bartsch S, Bauer B, Wiemann A, Clausen PH, Steuber S:** Feeding patterns of biting midges of the *Culicoides* obsoletus and *Culicoides* pulicaris groups on selected farms in Brandenburg, Germany. *Parasitol Res*, 105, 373-380, 2009. DOI: 10.1007/s00436-009-1408-y
- Garros C, Gardes L, Allene X, Rakotoarivony I, Viennet E, Rossi S, Balenghien T:** Adaptation of a species-specific multiplex PCR assay for the identification of blood meal source in *Culicoides* (Ceratopogonidae: Diptera): Applications on Palaearctic biting midge species, vectors of Orbiviruses. *Infect, Genet Evol*, 11 (5): 1103-1110, 2011. DOI: 10.1016/j.meegid.2011.04.002
- Kasičová Z, Schreiberová A, Kimáková A, Kočíšová A:** Blood meal analysis: Host-feeding patterns of biting midges (Diptera, Ceratopogonidae, *Culicoides* Latreille) in Slovakia, *Parasite*, 28:58, 2021. DOI: 10.1051/parasite/2021058
- Martínez-de la Puente J, Navarro J, Ferraguti M, Soriguer R, Figuerola J:** First molecular identification of the vertebrate hosts of *Culicoides* imicola in Europe and a review of its blood-feeding patterns worldwide: implications for the transmission of bluetongue disease and African horse sickness. *Med Vet Entomol*, 31, 333-339, 2017. DOI: 10.1111/mve.12247
- Kampen H, Werner D:** Blood meal analysis in Central European *Culicoides* biting midge species (Diptera: Ceratopogonidae) by reverse line blot hybridization. *Deutsche Gesellschaft für Allgemeine und Angewandte Entomologie*, 22, 225-232, 2020.
- Tomazatos A, Jöst H, Schulze J, Spínu M, Schmidt-Chanasit J, Cadar D, Lühken R:** Blood-meal analysis of *Culicoides* (Diptera: Ceratopogonidae) reveals a broad host range and new species records for Romania. *Parasit Vectors*, 13:79, 2020. DOI: 10.1186/s13071-020-3938-1
- Vasić A, Zdravković N, Anić D, Bojkovski J, Marinov M, Mathis A, Niculaua M, Oşlobanu EL, Pavlović I, Petrić D, Pflüger V, Pudar D, Savuša G, Simeunović P, Veronesi E, Silaghi C, the SCOPES AMSAR training group:** Species diversity, host preference and arbovirus detection of *Culicoides* (Diptera: Ceratopogonidae) in south-eastern Serbia. *Parasit Vectors*, 12:61, 2019. DOI: 10.1186/s13071-019-3292-3
- Mathieu B, Cêtre-Sossah C, Garros C, Chavernac D, Balenghien T, Carpenter S, Setier-Rio ML, Vignes-Lebbe R, Ung V, Candolfi E, Delécolle JC:** Development and validation of IIKC: An interactive identification key for *Culicoides* (Diptera: Ceratopogonidae) females from the Western Palaearctic region. *Parasit Vectors*, 5:137, 2012. DOI: 10.1186/1756-3305-5-137
- Morag N, Saroya Y, Braverman Y, Klement E, Gottlieb Y:** Molecular identification, phylogenetic status, and geographic distribution of *Culicoides oxystoma* (Diptera: Ceratopogonidae) in Israel. *PLoS One*, 7 (3):e33610, 2012. DOI: 10.1371/journal.pone.0033610
- Kent RJ, Norris DE:** Identification of mammalian blood meals in mosquitoes by a multiplexed polymerase chain reaction targeting cytochrome B. *Am J Trop Med Hyg*, 73 (2): 336-342, 2005.
- Pitzer JB, Kaufman PE, Tenbroeck SH, Maruniak JE:** Host blood meal identification by multiplex polymerase chain reaction for dispersal evidence of stable flies (Diptera: Muscidae) between livestock facilities. *J Med Entomol*, 48 (1): 53-60, 2011. DOI: 10.1603/me10123
- Kameke D, Kampen H, Walther D:** Activity of *Culicoides* spp. (Diptera: Ceratopogonidae) inside and outside of livestock stables in late winter and spring, *Parasitol Res*, 116 (3): 881-889, 2017. DOI: 10.1007/s00436-016-5361-2

32. **Kar S, Mondal M, Pal A, Harsha R, Mazumdar A:** Blood meal analysis of *Culicoides* species associated with livestock in West Bengal, India. *Med Vet Entomol*, 36 (4): 503-510, 2022. DOI: 10.1111/mve.12588
33. **Burgu I, Urman HK, Akça Y, Yonguç A, Mellor PS, Hamblin C:** Serologic survey and vector surveillance for bluetongue in southern Turkey. Bluetongue, African Horse Sickness, and Related Orbiviruses. In, Walton TE, Osburn BI (Eds): Proceedings of the Second International Symposium, CRC Press, Boca Raton, 1992.
34. **Slama D, Haouas N, Mezhoud H, Babba H, Chaker E:** Blood meal analysis of *Culicoides* (Diptera: Ceratopogonidae) in central Tunisia. *PLoS One*, 10 (3):e0120528, 2015. DOI: 10.1371/journal.pone.0120528
35. **Snyman J, Snyman LP, Labuschagne K, Venter GJ, Venter M:** The utilisation of CytB and COI barcodes for the identification of bloodmeals and *Culicoides* species (Diptera: Ceratopogonidae) reveals a variety of novel wildlife hosts in South Africa. *Acta Trop*, 219:105913, 2021. DOI: 10.1016/j.actatropica.2021.105913
36. **Riddin MA, Venter GJ, Labuschagne K, Villet MH:** Bloodmeal analysis in *Culicoides* midges collected near horses, donkeys and zebras in the Eastern Cape, South Africa. *Med Vet Entomol*, 33 (4): 467-475, 2019. DOI: 10.1111/mve.12381
37. **Dik B, Kuçlu Ö, Öztürk R:** *Culicoides* Latreille, 1809 (Diptera: Ceratopogonidae) species in the Western Black Sea Region of Turkey, new records for the Turkish fauna. *Turk J Vet Anim Sci*, 41 (2): 228-237, 2017. DOI: 10.3906/vet-1606-41
38. **Korkmaz Ç:** Distribution of *Culicoides* (Diptera: Ceratopogonidae) Species in Turkey and Ecological Approaches. *PhD Thesis*, Hacettepe University, Graduate School of Science and Engineering, 2022.

