

RESEARCH ARTICLE

Evaluating *Linum usitatissimum* Seeds Extract as Potential Alternative Biochemical and Therapeutic Agent Against Induced Coccidiosis in Broiler Chicken

Kashif HUSSAIN ¹  Asghar ABBAS ^{1(*)}  Atif REHMAN ^{1(*)}  Muhammad Umair WAQAS ¹ 
Baseer AHMAD ¹  Muhammad Adnan Sabir MUGHAL ¹  Rao Zahid ABBAS ² 
Muhammad Arfan ZAMAN ³  Junaid Ali KHAN ¹  Muhammad Asif RAZA ¹ 

¹ Department of Pathobiology Pathobiology, Faculty of Veterinary and Animal Sciences, Muhammad Nawaz Sharif University of Agriculture, 66000, Multan, PAKISTAN

² Department of Parasitology, Faculty of Veterinary Sciences, University of Agriculture, 38000, Faisalabad, PAKISTAN

³ College of Veterinary and Animal Sciences, 35200, Jhang, PAKISTAN

**(*) Corresponding authors:**

Atif REHMAN & Asghar ABBAS

Phone: +92 3458176856 (A.R.), +92 3361734029 (A.A.)

E-mail: atif.rehman@mnsuam.edu.pk (A.R.),
asghar.abbas@mnsuam.edu.pk (A.A.)

How to cite this article?

Hussain K, Abbas A, Rehman A, Waqas MU, Ahmad B, Mughal MAS, Abbas RZ, Zaman MA, Khan JA, Raza MA: Evaluating *Linum usitatissimum* seeds extract as potential alternative biochemical and therapeutic agent against induced coccidiosis in broiler chicken.. *Kafkas Univ Vet Fak Derg*, 30 (6): 803-808, 2024.

DOI: 10.9775/kvfd.2024.32618

Article ID: KVFD-2024-32618

Received: 06.07.2024

Accepted: 20.09.2024

Published Online: 03.10.2024

Abstract

Coccidiosis is a significant disease of poultry and is usually treated using various synthetic anticoccidial drugs. However, the effectiveness of this approach has been compromised due to drug resistance. Medicinal plants are being considered as potential alternatives to these drugs. As part of ongoing research, an investigation was conducted to assess the anticoccidial potential of *Linum usitatissimum* seed extract (LUE) and its impact on hematological parameters in broiler chickens with experimental coccidiosis. A total of 108 broiler chicks were divided into six groups. The chicks in Groups I, II, and III were given plant extract at three different concentrations: 4%, 5%, and 6%, respectively, at one week of age. Group IV was the positive control - these chickens were treated with toltrazuril. Group V served as the negative control, meaning they were infected but not given any medication. Group VI was the normal control group. All groups, except Group VI, were orally infected with 60,000 sporulated oocysts when they were 18 days old. After 7 days of giving orally infection, six birds from each group were euthanized so that we could examine their feed conversion ratio (FCR), oocyst score, lesion score, fecal score, serum chemistry and hematology. The results showed that the *L. usitatissimum* extract exhibited anticoccidial activity. It improved the FCR and reduced lesion, oocyst, fecal scores and blood and serum chemistry.

Keywords: Anticoccidial, Chicken, Coccidiosis, *Linum usitatissimum*

INTRODUCTION

Protozoa of the genus *Eimeria*, single-cell parasites from the phylum Apicomplexa, with complicated and several life cycle phases, cause avian coccidiosis ^[1]. *Eimeria* species mainly affect avian intestines, causing symptoms such as bloody diarrhea, poor feed conversion ratio (FCR), and even death. This results in considerable financial losses for chicken farming. *E. tenella* is one of the most harmful *Eimeria* species, causing cecal coccidiosis in hens. Each year, the poultry sector suffers significant financial losses due to coccidiosis, which starts with the intake of

sporulated oocysts ^[2]. Due to quick sporulation process, naturally occurring coccidian oocysts abound and generate millions of oocysts ^[3]. Although Sulfanilamide was first anticoccidial utilized as a treatment against coccidiosis in poultry, a range of anticoccidial feed additives and antibiotics have also been developed and used. Synthetic chemicals and anti-coccidials are usually employed to control coccidiosis. However, reasonable use of anticoccidial medications resistance has evolved ^[4]. Furthermore, time is needed to locate some alternative tools for efficient coccidiosis control. Based on their Therapeutic and immunomodulatory activity,



various plants were documented as anticoccidial and immunomodulatory over the past ten years [5-7]. Using antioxidant-rich and biological active plant extracts has become especially important in view of resistance to synthetic antibiotics, phenols, flavonoids, tannins and saponins are being used as an alternate approach to treat coccidiosis [8,9]. Like other plant with anthelmintic and therapeutic activity, *Linum usitatissimum* often referred as linseed and locally known Alsi contains numerous pharmacological properties against different parasitic and bacterial infections due to its diverse antioxidant chemicals [10]. Therefore, the present study was designed to evaluate the anticoccidial capacity of *L. usitatissimum* seed against induced infection in chickens which was caused by *Eimeria* based on existing literature including antioxidant properties.

MATERIAL AND METHODS

Ethical Approval

The research is conducted by the approval from ethical committee of University of Agriculture Faisalabad under the PSF, Project No. 185 and PARB, Project No. 358 (No. 628/6-08-2013)

Plants Material

After procuring *L. usitatissimum* (Alsi) seeds from the local market and they were crushed using a grinder machine. Afterwards, we prepared an aqueous methanolic extract of *L. usitatissimum* using Soxhlet's apparatus (Velp Italy) following the method described by Abbas et al. [11]. The suspension was evaporated in a rotary evaporator (Heidolph Germany) at a temperature not exceeding 50°C. The prepared *L. usitatissimum* extract was then stored in a refrigerator at 4°C after freeze drying until further use.

Parasite

The parasite material was collected from outbreak cases in Faisalabad as well as the intestines of naturally infected chickens with *Eimeria*. To induce sporulation, the material was immersed in a potassium dichromate solution (2.5%) at 25-29°C and 60-80% humidity. The sample was next inspected under a microscope [12].

The number of sporulated oocysts was calculated using the Modified McMaster procedure. The material was put into the chambers of the McMaster slide and left alone for 2-3 min to let the sporulated oocysts to float and become visible. The slide was next examined under a microscope at low (10x) and high (40x) magnification

Experimental Design

A total of 108 broiler chicks were obtained and divided into six groups, with 18 birds in each group. Groups I, II, and III were given plant extract at three different

concentrations (4%, 5%, and 6%, respectively) at one week of age. Group IV was kept as the positive control (infected plus toltrazuril® from A&K Pharmaceuticals, Faisalabad, provided at a rate of 1 mL/L of water). Group V served as the negative control (infected and non-medicated), and group VI as the normal control (non-infected and non-medicated). By the time the chickens reached eighteen days, all groups except group VI had been orally infected with 60,000 sporulated oocysts. Six birds from each group were sacrificed seven days after the inoculation to collect data on FCR, oocyst score, lesion score, fecal score, serum chemistry, and hematology.

Evaluation of Anticoccidial Activity

The potential of (LUE) as an anticoccidial treatment was investigated using metrics such as (FCR), lesion score [13] and oocyst score [14]. The fecal scores of birds in each group were monitored to assess the severity of illness at day 3 to 7. To evaluate the fecal score in chickens, a standard method is used. The optimal time to assess the emergence of illness is between the third and seventh days after inoculation. The fecal score chart ranges from 1 to 5, representing increasing degrees of disease progression [15].

Hematological Parameters

The collected blood samples were tested for packed cell volume (PCV) by following the microhematocrit method with slight modification. Hemoglobin level (Hb) was determined using Sahli's device. Erythrocyte and leukocyte counts were performed using a hemocytometer using Natt and Herrick solution under compound microscope at 10x.

Serum Chemistry

The plant extract was evaluated for toxic possession, cellular injury, and serum samples using various imported assays (Merck, Germany) to determine the levels of serum enzymes (LAT, LDH, Creatinine) [16].

Statistical Analysis

Statistical analysis was performed using the ANOVA approach and SAS statistical analysis software version 9 [17]. The data was considered statistically significant with a P value <0.05.

RESULTS

All the groups offered with *L. usitatissimum* extract (LUE) revealed improved FCR at classified doses in [Table 1](#). However, the admirable result showed by the group which was administered with higher dose and the results were similar (P>0.05) to standard medicine (Toltrazuril®)

A lower lesion score at graded doses was observed in (LUE) treated group in [Table 2](#). However, the admirable result showed by the group which was administered with

Table 1. Feed conversion ratio (FCR) of plant extract treated groups

Groups	Feed Consumed (g)	Ending Weight (g)	Feed Conversion Ratio (g/g)*
<i>L. usitatissimum</i> 4%	920.1	418.18	2.20
<i>L. usitatissimum</i> 5%	990.2	405.45	2.44
<i>L. usitatissimum</i> 6%	941.5	408.90	2.30
Positive control	910.16	409.86	2.22
Negative control	964.4	368.40	2.61
Normal Control	916.6	422.6	2.16

*Due to feeding in group statistical analysis was not achievable because FCR is simple ratio

Table 2. Lesion score of different plant extract treated groups

Groups	0	+1	+2	+3	+4	Mean
<i>L. usitatissimum</i> 4%	0	1	2	2	1	1.83±0.54 ^b
<i>L. usitatissimum</i> 5%	0	2	2	2	0	1.50±0.51 ^c
<i>L. usitatissimum</i> 6%	0	3	3	0	0	1.66±0.83 ^{bc}
Positive control	1	3	2	0	0	1.33±0.40 ^d
Negative control	0	0	0	3	3	3.33±0.51 ^a
Normal Control	0	0	0	0	0	0

Means with different superscripts are significantly different ($P<0.05$) from each other
+1: No lesions, +2: Very few, +3: Large amount, +4: Blood and Death

Table 3. Oocyst score of treated groups with plant extract

Groups	0	+1	+2	+3	+4	+5	Mean
<i>L. usitatissimum</i> 4%	0	1	3	1	1	0	2.00±0.75 ^b
<i>L. usitatissimum</i> 5%	0	2	2	2	0	0	1.50±0.51 ^c
<i>L. usitatissimum</i> 6%	0	3	3	0	0	0	1.66±0.54 ^{bc}
Positive control	0	2	3	1	0	0	1.66±0.75 ^{bc}
Negative control	0	0	0	3	2	1	3.83±0.40 ^a
Normal Control	-	-	-	-	-	-	-

Means with different superscripts are significantly different ($P<0.05$) from each other
0: No oocyst, +1: 1-10 oocyst/Field, +2: 11-20 oocyst/Field, +3: 21-50 oocyst/Field, +4: 51-100 oocyst/Field, +5: More than 100 oocyst/Field

Table 4. Fecal score of groups treated with Plant Extract

Groups	Day 3 rd	Day 4 th	Day 5 th	Day 6 th	Day 7 th
<i>L. usitatissimum</i> 4%	-	2.04±0.75 ^{ab}	2.15±0.75 ^b	1.12±0.63 ^b	-
<i>L. usitatissimum</i> 5%	-	1.34±0.51 ^b	2.01±0.63 ^b	1.52±0.54 ^b	-
<i>L. usitatissimum</i> 6%	-	1.52±0.54 ^{ab}	1.68±0.51 ^b	1.51±0.54 ^b	-
Positive control	-	1.51±0.83 ^{ab}	1.68±0.51 ^b	1.34±0.51 ^b	-
Negative control	-	2.68±1.40 ^a	2.99±0.42 ^a	2.71±0.79 ^a	-
Normal Control	-	-	-	-	-

Means with different superscripts are significantly different ($P<0.05$) from each other

Table 5. Serum enzymes values of groups treated Plant Extract (Mean±SD)

Groups	ALT	ASAT	LDH	Urea	Creatinine
<i>L. usitatissimum</i> 4%	9.46±0.91 ^b	180.72±10.08 ^b	476.01±16.91 ^b	5.40±0.81 ^b	0.21±0.02 ^b
<i>L. usitatissimum</i> 5%	9.57±0.97 ^b	182.60±12.14 ^b	483.95±21.52 ^b	5.55±0.97 ^b	0.18±0.02 ^b
<i>L. usitatissimum</i> 6%	9.74±1.38 ^b	177.55±14.72 ^b	477.24±22.21 ^b	5.60±1.02 ^b	0.19±0.03 ^b
Positive control	9.69±1.15 ^b	181.85±10.15 ^b	477.42±21.15 ^b	5.20±0.48 ^b	0.19±0.02 ^b
Negative control	24.62±2.31 ^a	288.87±36.21 ^a	891.96±22.16 ^a	20.60±1.12 ^a	0.70±0.03 ^a
Normal Control	8.98±1.78 ^b	195.03±13.46 ^b	471.45±15.78 ^b	5.45±0.59 ^b	0.16±0.02 ^b

Means with different superscripts are significantly different (P<0.05) from each other

Table 6. Hematological values o of groups treated Plant Extract (Mean±SD)

Groups	PCV %	Hb g/dL	RBC 10 ⁶ /μL	WBC 10 ³ /μL
<i>L. usitatissimum</i> 4%	29.14±1.67 ^a	10.22±1.06 ^a	3.99±0.76 ^a	21.67±2.93 ^b
<i>L. usitatissimum</i> 5%	28.20±1.61 ^a	11.43±1.35 ^a	3.64±0.86 ^a	22.74±2.70 ^b
<i>L. usitatissimum</i> 6%	27.01±1.66 ^a	11.81±1.28 ^a	3.41±0.71 ^a	21.66±1.64 ^b
Positive control	24.17±2.14 ^a	11.20±0.64 ^a	3.27±0.71 ^b	21.41±2.78 ^b
Negative control	20.15±1.15 ^b	8.91±0.82 ^b	1.88±0.12 ^c	34.01±5.04 ^a
Normal Control	25.34±1.03 ^a	11.0±1.34 ^a	3.15±0.58 ^{ab}	22.51±3.26 ^b

Means with different superscripts are significantly different (P<0.05) from each other

a higher dose and the results were similar (P>0.05) to standard medicine (Toltrazuril[®])

All groups treated with *L. usitatissimum* extract (LUE) showed minimal oocyst scores at graded doses in Table 3. However, the admirable result showed by the group which was administered with higher dose and the results were similar (P<0.05) to standard medicine (Toltrazuril[®])

L. usitatissimum extract (LUE) administered groups showed minimal fecal score at graded doses in Table 4. However, the admirable result showed by the group which was administered with higher dose and the results were similar (P<0.05) to standard medicine (Toltrazuril[®])

Minimal serum enzyme values in all (LUE) administered groups were observed in Table 5. However, the admirable result showed by the group which was administered with higher dose and the results were similar (P>0.05) to standard medicine (Toltrazuril[®])

Maximum hematological (PCV, Hb, RBCs and WBCs) values were observed in Table 6. However, the admirable result showed by the group which was administered with higher dose and the results were similar (P>0.05) to standard medicine (Toltrazuril[®])

DISCUSSION

Recent reports have explored alternative methods for treating coccidiosis by using plant antioxidant composites such as phenols, flavonoids, tannins, and saponins [1,4,11,18]. This investigation found that *L. usitatissimum* shows

anticoccidial activity and has positive effects on serum chemistry, hematological values, oocyst score, lesion score, and fecal score. The results were identical to those of accomplished with the traditional medication (Toltrazuril[®]). Specifically, *L. usitatissimum*, when administered at doses of 4%, 5%, and 6%, confirmed a dose-dependent potential to combat mixed *Eimeria* infections. The maximum concentration confirmed results corresponding to the Toltrazuril[®] dealt with group (P>0.05). Previous studies inspecting the capacity of various herbal extracts to combat coccidiosis have also mentioned similar findings, with the effectiveness of the extracts varying depending on the dosage administered [19-21].

In a study, the weight gains in broiler chicks infected with *Eimeria* was significantly reduced by an ethanolic extract from *Carica papaya* leaves [22]. Studies showed that orally administered *Ageratum conyzoides* extract to 28-day-old chicks increased the number of red blood cells (RBCs), white blood cells (WBCs), and PCV in the treated birds. This eventually resulted in a decrease in infection levels and excretion of oocysts. The antioxidant chemicals found in *A. conyzoides*, including flavonoids, phenols, conyzorium, methexneblitin, and quercetin, may reduce the oxidative stress caused by coccidiosis. Likewise, *A. conyzoides* has been found to have anticoccidial properties. This means that it can diminish *Eimeria* infection by inquisitive with lipid peroxidation [8,23]. The study found that the groups given the plant extract had lower levels of certain serum enzymes similar to the control groups, indicating that the

extract did not have any harmful effects. Hepatotoxicity can be determined by looking at the levels of AST and ALT, while nephrotoxicity can be determined by studying the levels of serum creatinine and urea.

A recent study showed that pulp of olive fruit when combine with vitamin C gives positive results on hatchability, weight gain, and improved growth rate [24]. Recent study of catechin *Uncaria gambir* extract which is planned on broiler birds to see meat quality, growth rate, serum /Plasma values and also antioxidant activity gives favorable results [25].

There are some other studies which showed the antioxidant, biomedical, immunomodulatory, antibacterial, resistant free effects of plant material, zinc oxide, Sodium Alginate and nanoparticles on animals [26].

The findings mentioned above indicate that plant-derived extracts could effectively help in dealing with chicken coccidiosis and its associated toxicity. These results also highlight the potential of *L. usitatissimum* as a safe and herbal anticoccidial agent, which warrants further research into the plant's active substances and modes of action. By incorporating *L. usitatissimum* into chicken health management, the risk of drug resistance may be reduced, and overall chicken health may be improved. This could present a sustainable and successful alternative to synthetic anticoccidial medicines

DECLARATIONS

Availability of Data and Materials: Research data will be provided by the author (K. Hussain) on request.

Acknowledgements: Support for this study was provided by the Punjab Agricultural Research Board, Project No. 358 (Assessment and Advancement of herbal anticoccidials for the management of coccidiosis in poultry) and the Pakistan Science Foundation, Project No. 185.

Funding Support: The Main Researcher recognizes the financial help for this research from Pakistan Science Foundation, Project No. 185.

Ethical Approval: The research is conducted by the approval from ethical committee of University of Agriculture Faisalabad under the PSF, Project No. 185 and PARB, Project No. 358 (No. 628/6-08-2013)

Competing Interest: There is no conflict of interest.

Declaration of Generative Artificial Intelligence (AI): This whole article is free from any AI tool.

Author Contributions: KH apprehended and planned the study; AA, AR, MUW, BA, MASM did work on methodology and RZA, MAZ, JAK, MAR and investigated the data and help in the writeup of the manuscript

REFERENCES

1. Abbas A, Alkheraije KA: Immunomodulatory effects of *Carica papaya* extract against experimentally induced coccidiosis in broiler chickens. *Pak*

Vet J, 43 (3): 628-632, 2023. DOI: 10.29261/pakvetj/2023.089

2. Chapman HD: Milestones in avian coccidiosis research: A review. *Poult Sci*, 93, 501-511, 2014. DOI: 10.3382/ps.2013-03634

3. Fall AD, Gbati OB, Diatta W, Lapo RA, Diatta-Badji K, Dieng M, Ibra S, Dieng SIM, Bassene E, Pangui LJ: Anticoccidial activity of ethanol roots extract of *Cassia sieberiana* DC in chickens. *European J Med Plants*, 11, 1-7, 2016. DOI: 10.9734/EJMP/2016/21343

4. Hussain K, Abbas A, Alanazi HAH, Alharbi AMA, Alaiiri AA, Rehman A, Waqas MU, Raza MA, Yasin R, Ahmad B, Bano N, Khara HURA: Immunomodulatory effects of *Artemisia brevifolia* extract against experimentally induced coccidiosis in broiler chicken. *Pak Vet J*, 43 (2): 333-338, 2023. DOI: 10.29261/pakvetj/2023.026

5. Khalifa WH, Sallam MG, Kamel NN, Samy A, Yassein SA, El-Mallah GM, Abusinaa GE: Using *in-ovo* injection of olive pulp extract and vitamin C to improve hatchability, post hatch growth performance, carcass traits and some biochemical blood analysis in broiler chickens. *Int J Vet Sci*, 12 (3): 353-359, 2023. DOI: 10.47278/journal.ijvs/2022.200

6. Liaqat I, Noor S, Qureshi AS, Ali S, Liaqat I, Al-Arifa N, Alam S, Ajmal A, Zia T, Munawar M: Biosynthesis and evaluation of *Cinnamomum zeylanicum* nanomaterials for the treatment of polycystic ovary syndrome in mice. *Pak Vet J*, 43 (1): 118-124, 2023. DOI: 10.29261/pakvetj/2023.004

7. Fariha, Jahan N, Rehman KU, Ali S: Effect of herbal mixture as angiotensin-converting enzyme inhibitor in angiotensin-II dependent hypertension. *Pak Vet J*, 39 (1): 25-30, 2019. DOI: 10.29261/pakvetj/2018.103

8. Nweze NE, Obiwulu IS: Anticoccidial effects of *Ageratum conyzoides*. *J Ethnopharmacol*, 122 (1): 6-9, 2009. DOI: 10.1016/j.jep.2008.11.014

9. Velázquez-Antunez J, Olivares-Perez J, Olmedo-Juárez A, Rojas-Hernandez S, Villa-Mancera A, Romero-Rosales T, Zamilpa A, Gonzalez-Cortazar M: Biological activity of the secondary compounds of *Guazuma ulmifolia* leaves to inhibit the hatching of eggs of *Haemonchus contortus*. *Pak Vet J*, 43 (1): 55-60, 2023. DOI: 10.29261/pakvetj/2022.075

10. Al-Saeed FA, Bamarni SSI, Iqbal KJ, Rehman TU, Faruk AZ, Mahmood S, Şahin T, Ölmez M, Riaz R: *In vitro* anthelmintic efficacy of *Haloxylon salicornicum* leaves extract using adult *Haemonchus contortus* worms. *Pak Vet J*, 43 (1): 91-96, 2023. DOI: 10.29261/pakvetj/2022.091

11. Abbas A, Iqbal Z, Abbas RZ, Khan MK, Khan JA, Sindhu ZD, Mahmood MS, Saleemi MK: *In vivo* anticoccidial effects of *Beta vulgaris* (sugar beet) in broiler chickens. *Microb Pathog*, 111, 139-144, 2017. DOI: 10.1016/j.micpath.2017.07.052

12. Ryley JF, Meade R, Burst JH, Robinson TE: Methods in coccidiosis research separation of oocysts from faeces. *J Parasitol*, 73, 311-326, 1976. DOI: 10.1017/S0031182000046990

13. Johnson J, Reid WM: Anticoccidial drugs lesion scoring techniques in battery and floor pen experiments with chickens. *Exp Parasitol*, 28, 30-36, 1970. DOI: 10.1016/0014-4894(70)90063-9

14. Hilbrich P: Krankheiten des geflügelsunterbesondererberücksichtigung der haltung und fütterung. Hermann kuhn kg. Schwenningen am neckar Germany, 1978.

15. Habibi H, Firouzi S, Nili H, Razavi M, Asadi SL, Daneshi S: Anticoccidial effects of herbal extracts on *Eimeria tenella* infection in broiler chickens: *In vitro* and *in vivo* study. *J Parasit Dis*, 2014. DOI: 10.1007/s12639-014-0517-4

16. Abbas RZ, Iqbal Z, Khan MN, Zafar MA, Zia MA: Anticoccidial activity of *Curcuma longa* L. in broiler chickens. *Braz Arch Biol Technol*, 53, 63-67, 2010. DOI: 10.1590/S1516-89132010000100008

17. SAS: SAS Statistical Software Version 9.1. SAS Institute Inc. Cary, NC, USA, 2004.

18. Molan AL, Liu Z, Shampa DE: Effect of pine bark (*Pinus radiata*) extracts on sporulation of coccidian oocysts. *Fol Parasitol*, 56, 1-5, 2009. DOI: 10.14411/fp.2009.001

19. Dkhil MA, Abdel-Baki AS, Wunderlich F, Sies H, Al-Quraishy S: Anticoccidial and anti-inflammatory activity of garlic in murine *Eimeria papillata* infections. *Vet Parasitol*, 175, 66-72, 2011. DOI: 10.1016/j.vetpar.2010.09.009

20. Yang WC, Tien YJ, Chung CY, Chen C, Chiou WH, Hsu SY, Liu HY, Liang CL, Chang CLT: Effect of *Bidens pilosa* on infection and drug

resistance of Eimeria in chickens. *Res Vet Sci*, 98, 74-81, 2015. DOI: 10.1016/j.rvsc.2014.11.002

21. Wang D, Zhou L, Li W, Zhou H, Hou G: Anticoccidial effect of *Piper sarmentosum* extracts in experimental coccidiosis in broiler chickens. *Trop Anim Health Prod*, 5, 1071-1078, 2016. DOI: 10.1007/s11250-016-1034-5

22. Nghonjuyi NW, Tiambo CK, Kimbi HK, Manka'a N, Juliano, F. Lisita RS: Efficacy of ethanolic extract of *Carica papaya* leaves as a substitute of sulphonomide for the control of coccidiosis in KABIR chickens in cameroon. *J Anim Health Prod*, 3 (1): 21-27, 2015. DOI: 10.14737/journal.jahp/2015/3.1.21.27

23. Hussain K, Alsayeqh AF, Abbas A, Abbas RZ, Rehman A, Zaib W, Rehman TU, Mahmood MS: Potential of *Glycyrrhiza glabra* (Licorice) extract an alternative biochemical and therapeutic agent against coccidiosis in broiler chickens. *Kafkas Univ Vet Fak Derg*, 28 (5): 585-591, 2022. DOI:

10.9775/kvfd.2022.27620

24. Ramaiyulis, Mairizal, Salvia, Fati N, Malvin T: Effects of dietary catechin *Uncaria gambir* extract on growth performance, carcass characteristics, plasma lipids, antioxidant activity, and nutrient digestibility in broiler chickens. *Int J Vet Sci*, 12 (2): 169-174, 2023. DOI: 10.47278/journal.ijvs/2022.177

25. Elbehary M, Dowidar YA, Ashour AM, Abd El Fattah EM, Monir A: Effect of *in vitro* maturation medium supplementation with chitosan nanoparticles on the river buffalo cumulus-oocyte complexes. *Int J Vet Sci*, 12 (3): 341-346, 2023. DOI: 10.47278/journal.ijvs/2022.188

26. El-Hamaky AMA, Hassan AA, Wahba AKA, El Mosalamy MMEA: Influence of copper and zinc nanoparticles on genotyping characterizations of multi-drug resistance genes for some calf pathogens. *Int J Vet Sci*, 12 (3): 309-317, 2023. DOI: 10.47278/journal.ijvs/2022.195