

REVIEW ARTICLE

Managing Mycotoxins in Animal/Poultry Feed Through Innovative Control Strategies: A Review

Aisha KHATOON^{1 (*)}  Zain ul ABIDIN²  Ashiq ALI³  Muhammad Kashif SALEEMI¹  Rao Zahid ABBAS⁴  Muhammad Umar IJAZ⁵  Mujeeb ur REHMAN⁶  Bilal MURTAZA⁷ 

¹ Department of Pathology, Faculty of Veterinary Science University of Agriculture Faisalabad 38040 PAKISTAN

² Veterinary Research Institute, Zarrar Shaheed Road Lahore Cantt-13 PAKISTAN

³ Department of Histology and Embryology Shantou Medical University, Shantou, 515041 CHINA

⁴ Department of Parasitology, Faculty of Veterinary Science University of Agriculture Faisalabad, 38040 PAKISTAN

⁵ Department of Zoology, Wildlife and Fisheries, University of Agriculture Faisalabad-38040 PAKISTAN

⁶ Single Cell Bioengineering Group (SCBEG), State Key Laboratory of Marine Resource Utilization in South China Sea, College of Oceanology, Hainan University, Haikou 570228, Hainan, CHINA

⁷ School of Bioengineering, Dalian University of Technology, No. 2 Linggong Road, Dalian, 116024, CHINA



(*) Corresponding author:

Aisha Khatoon

Phone: +92 92-42-9200161

E-mail: aisha.khatoon@uaf.edu.pk;

aishavp@yahoo.com

How to cite this article?

Khatoon A, Abidin ZU, Ali A, Saleemi MK, Abbas RZ, Ijaz MU, Rehman MU, Murtaza B: Managing mycotoxins in animal/poultry feed through innovative control strategies: A review. *Kafkas Univ Vet Fak Derg*, 2025 (Article in Press).

DOI: 10.9775/kvfd.2025.34700

Article ID: KVFD-2025-34700

Received: 30.06.2025

Accepted: 12.09.2025

Published Online: 17.09.2025

Abstract

Mycotoxins are the secondary metabolites of certain toxigenic fungi that have deleterious effects upon the health of humans, animals, and poultry. More than 300 chemically different mycotoxins have been identified to date, among which the most important are aflatoxins, ochratoxins, fumonisins, trichothecenes, and patulins. Approximately 25% of global food crops are significantly affected by mycotoxins every year. Animals become exposed to the adverse effects of mycotoxins when fed mycotoxin-contaminated feed, and animal byproducts containing mycotoxin residues become a constant source of exposure to the human population. Once mycotoxins enter the food chain, their complete removal is inevitable; therefore, different control strategies are being adopted to minimize the adverse effects associated with them. This review encompasses various control strategies adapted to minimize mycotoxicosis.

Keywords: Mycotoxins, Animal feed, Poultry feed, Mycotoxicosis, Control strategies

INTRODUCTION

Certain toxic fungal species are ubiquitous in nature and have strong ecological link with human and animal food supplies. These fungi often produce certain chemical ingredients which are not necessarily required for their growth but play a crucial role in their survival and these chemical compounds are often termed as 'secondary fungal metabolites'. Mycotoxins are a diverse group of chemically different compounds originally produced as secondary metabolites by several toxigenic fungal species. Many fungal genera are predominantly involved in the production of these hazardous chemical compounds, but the most important among them are *Aspergillus*, *Fusarium*, *Penicillium*, *Alternaria* and *Claviceps*^[1]. To date, about 400

chemically diverse mycotoxins have been identified which pose severe toxic effects in different animal species and human population in one way or another. Mycotoxin-associated toxicities and/or adverse effects are directly related to the dose, duration, mycotoxin type, and route of exposure to a specific mycotoxin^[2]. Mycotoxins generally exhibit hepatotoxic, nephrotoxic, carcinogenic, and immunosuppressive effects in certain animal species, ultimately compromising the overall health of animals. When produced within the feed, mycotoxins form 'mycotoxin pockets' within the feed, which are generally rich in their concentration and are considered hotspots for mycotoxins, thereby ensuring an uneven distribution of mycotoxins within the feed.



The route of mycotoxin entry into the animal food/feed chain involves the use of contaminated agricultural byproducts during the formation of feed or production of different mycotoxins within the feed by storage fungi descending either from pre-harvest, harvest, or post-harvest durations of crops^[3]. Contamination by different agricultural products also limits international trade, as certain countries have different regulatory measures regarding the levels of mycotoxins^[4]. Entry of mycotoxins

into the human food chain occurs either through the consumption of mycotoxin-contaminated agricultural products or through animal byproducts derived from animals fed mycotoxin-contaminated rations. The mycotoxin residues exhibit certain anomalies in humans, the details of which have been presented and elaborated in *Table 1* and *Fig. 1*.

Discussing about the classification of mycotoxins, some of the significant mycotoxins from the vast list are aflatoxins,

Table 1. Mycotoxins associated diseases in human population along with details of specific mycotoxins involved in disease production				
Fungi Involved	Source of Contamination	Disease Production	Specific Mycotoxin Involved	Reference
<i>Fusarium verticillioides</i> , <i>Fusarium proliferatum</i>	Corn/maize	Oesophageal tumors	Fumonisin B2	[5]
<i>Fusarium</i> species	Toxic bread (cereal grains)	Alimentary toxic aleukemia	Trichothecene, Fumonisin B1	[6]
<i>Stachybotrys atra</i>	Contaminated dust from ventilation slits, ceilings and walls	Sick-building syndrome	T-2, Diacetoxyscirpenol, Verrucarol	[7]
<i>Fusarium</i> species	Wheat, oats, barley, rice	Akakabio-byo	Fusarium toxins	[8]
<i>Aspergillus flavus</i> and <i>Aspergillus parasiticus</i>	Cereal grains	Kwashiorkor	Aflatoxin M1, AFM2	[9]
<i>Aspergillus ochraceus</i> , <i>Penicillium</i> species	Cereal grains	Balkan endemic nephropathy	Ochratoxin A	[10]
<i>Penicillium</i> and <i>Aspergillus</i> species	Rice	Cardiac beriberi	Aflatoxins	[11]
<i>Fusarium proliferatum</i> , <i>Fusarium verticillioides</i>	Maize/corn	Neural tube defect	Fumonisin B1 and B2	[12]
<i>Claviceps purpurea</i> , <i>Claviceps fusiformis</i>	Cereal grains, rye	Ergotism	Ergotamine-ergocristine Alkaloids	[12]
<i>Aspergillus flavus</i> and <i>Aspergillus parasiticus</i>	Peanuts, cereal grains	Hepatocellular carcinoma	Aflatoxin B1	[13]
<i>Stachybotrys atra</i>	Grain dust	Stachybotryotoxicosis	Satratoxins, Trichothecene	[14]
<i>Aspergillus</i> species	Grain dust	Reye's syndrome	Aflatoxin B1, B2 and M1	[15]
<i>Fusarium</i> species	Wheat, rice, corn	Scabby grain toxicosis	Zearalenone, deoxynivalenol	




Fig 1. Spectrum of mycotoxin associated diseases in human

ochratoxins, zearalenone, trichothecenes, fumonisins, patulin, ergot toxins etc. However, keeping in view the one health perspectives, the most significant among all the mycotoxins are aflatoxins and ochratoxins. This review emphasizes the existence and after-effects of mycotoxins and control strategies adapted for these mycotoxins (in particular), along with the strategies adapted for the control of other mycotoxins in different livestock species.


EXISTENCE OF DIFFERENT MYCOTOXINS IN ANIMAL/POULTRY FEED AND THEIR EFFECTS

Aflatoxins

Contamination of food with aflatoxins has remained a

persistent issue for livestock and poultry feed, along with all processed food products. These chemical compounds were discovered accidentally in 1961 when many turkey poult suddenly died in England due to an unrecognized disorder named as "Turkey-X disease." When investigated, it was revealed that a similar syndrome also appeared in farms where Brazil imported moldy peanut meal during feed formulation. Extraction using chloroform and detailed chemical analysis linked the extracted compound with "*Aspergillus flavus*." Scientists then gave the name "Aflatoxin" to this chemical compound by joining first three letters of both "*Aspergillus*" and "*flavus*." In the same year, this compound was also isolated in crystalline form in Netherlands and further fragmented as Aflatoxin B and G in United Kingdom based on color, they fluoresced under ultraviolet (UV) light. However, further investigations subdivided it into aflatoxins B1, B2, G1, and G2 based on minor differences in their chemical structures [16,17]. Furthermore, many other derivatives are linked to it from time to time, such as aflatoxin M1, aflatoxicol, AFP1, and AFQ1. There are approximately 20 different types of aflatoxins, but aflatoxin B1 (AFB1) (Fig. 2) is considered to be the most important and toxic among all types due to its toxicity [18].

Aflatoxins, belonging to the difuranocoumarin group, are secondary metabolites produced by a variety of toxigenic fungal species belonging to two important genera, *Aspergillus* and *Penicillium* [19] with *Aspergillus flavus* and *Aspergillus parasiticus* are considered major producers of aflatoxins [20,21]. Among all aflatoxins, aflatoxin B1 (AFB1) is the most toxic to animals, humans, and poultry, followed by AFB2, AFG1, and AFG2. Based on its extent of toxicity, the International Agency for Research on Cancer (IARC) has classified it as a group 1 carcinogen in humans and animals. Along with cancer, it causes hepatic disorders, metabolic diseases, vomiting, stunted growth, and diarrhea in human population.



Cereal crops, such as corn, wheat, sorghum, and rice, along with other feed ingredients, are readily contaminated with aflatoxins during their storage period when anaerobic conditions coupled with high humidity develop within the stored ingredients [6,22]. Aflatoxins have a tendency to easily infiltrate body tissues, muscles, and fatty tissues as residues, and when consumed, they become a potent source of contamination in the human population consuming such meat [23]. Table 2 illustrates some studies on mycotoxins reporting the existence of tissue residues in animals. The problem of aflatoxins occasionally occurs in crops prior to harvesting, but they are also produced in the stored ingredients whenever the storage fungi get a favorable environment for their growth, ultimately producing them as their secondary metabolites [24]. There are different legislations for the maximum tolerable levels (MTL) for all foods, including feed/feed ingredients for poultry and large animals (sheep, cattle, and buffalo). The United States Food and Drug Administration (US FDA) recommends 20 µg/kg of feed as a worldwide range for maximum permissible and tolerable levels for the poultry sector, whereas the maximum tolerable level for cattle and buffalo is 100 µg/kg of feed [25,26]. However, many studies are available, particularly in developing countries, which report much higher levels in feed/feed ingredients than in the recommended MTL [23, 27-32].

Table 2. Occurrence of tissue residues of different mycotoxins in various animal species

Mycotoxin	Specie	Level Used	Residues Detected	Reference
Aflatoxins	Broiler	100 µg/kg	0.32 µg/kg 0.08 µg/kg (Muscles)	[33]
	Broiler	1 mg/kg	0.166 µg/kg	[34]
	Broiler	1600-6400 µg/kg	6.97 ng/g 0.49-2.18 ng/g (Muscles)	[35]
	Broiler	Field Study	0.78-10.41 ng/g 0.23-5.67 ng/g (Kidney) 0.01-0.97 ng/g (Muscles)	[36]
	Broiler	600-1800 µg/kg	0.53-2.05 µg/kg	[37]
	Cattle	Field Study	0.36 µg/kg 1.37 µg/kg (Kidney)	[38]
Ochratoxins	Broiler	Field Study	0.58 µg/kg 0.51 µg/kg (Kidney)	[39]
	Broiler	100 µg/kg	1.92 ng/g 3.58 ng/g (Kidney)	[40]
	Broiler	2 mg/kg	1.79 ng/g 4.42 ng/g (Kidney)	[41]
	Beef sausages	Field Study	4.1-7.1 ppm	[42]

Table 3. Health impacts associated with aflatoxins in ruminants and poultry				
Specie	Dose Tested	Duration	Effects	Reference
Cattle	300 µg/kg	133 days	No effects	[48]
Cattle	200-500 µg/kg	14 days	Severe pathological effects	[49]
Cattle	350-455 µg/kg	15-17.5 weeks	No effects	[50]
Cattle	60-300 µg/kg	155 days	No effects	[51]
Beef calves	1400 µg/kg	FS	Neurological signs, ataxia, depression	[52]
Lambs	2 mg/kg	37 days	Decreased body weight and immune responses	[53]
Lambs	350 µg/kg	150 days	Decreased serum parameters	[54]
Lambs	5.9-23.5 µg/kg	91 days	Decreased cellular immunity	[55]
Lactating dairy cows	96 µg/kg	7-12 days	Slight increase in serum proteins	[56]
Broilers	40 µg/kg	42 days	Reduced growth performance	[57]
White Leghorn cockerels	400 µg/kg	60 days	Hematological alterations	[58]
Broiler chicks	100-600 µg/kg	42 days	Immunosuppression	[16]
Broiler chicks	200 µg/kg	42 days	Serum biochemical and immunological alterations	[59]

Ruminants: Ruminants have a complex ecosystem of microflora and microfauna within the rumen [43,44] and nature has provided ruminants with diverse properties for detoxifying mycotoxins into their less toxic products through a diversified range of microflora and microfauna existing within their ruminal fluid [45]. As far as cattle are concerned, mature animals are less prone to the adverse effects of aflatoxins compared to growing, young, and pregnant animals. Aflatoxins are degraded in the rumen and converted to less toxic aflatoxicol [46]. Aflatoxins in feed bind with ruminal contents, and a lower quantity (only 2-5%) of ingested aflatoxin reaches the intestine. Feed levels of approximately 100 µg/kg are toxic to ruminants [47]. The health effects of aflatoxins in different species are listed in *Table 3*.

Aflatoxin B1 (AFB1), which escapes ruminal degradation, enters the liver and is converted into aflatoxin (AFM1), which is released in milk. The maximum tolerable level of AFM1 in milk is 0.5 µg/kg. This metabolite can be detected in milk 6 h after the ingestion of AFB1, whereas its peak level can be noticed 24-48 hours after continuous AFB1 ingestion. Its clearance from milk can be observed 3 days after withdrawal of a controlled diet [60]. It has been reported that AFM1 can cause pronounced aflatoxicosis in weaning calves often characterized by development of histopathological lesions in liver and kidney along with disturbance of hepatic enzymes [61]. About 1-2% of the total ingested AFB1 is released as AFM1 in milk [62]. The average transfer of aflatoxin from feed to milk is 1.7% while the maximum permissible level of aflatoxin for milk is 0.05 µg/liter in Asia [63]. Therefore, to avoid its residues in milk, dietary aflatoxin levels for ruminants must be as low as 25 µg/kg [64].

Poultry: In the poultry industry, chicks of all age groups are prone to adverse effects caused by aflatoxins, particularly AFB1. Apart from adversely affecting all organs of chicks, their residues infiltrate muscles and organs, making them a direct source of contamination for the human population consuming such meat. Aflatoxins in feed result in reduced feed intake and body weight gain, along with suppressed relative organ weights, and birds become dull and less attractive towards feed. Similarly, hematological and serum biochemical parameters are adversely affected, leading to anemic conditions, along with permanent damage to the kidney and liver [65,66]. In addition, the most prominent anomaly expressed by AFB1 in feed is immunosuppression, which makes birds susceptible to secondary bacterial infections [67,68]. Moreover, AFB1 forms DNA adducts, ultimately affecting the overall genome of organisms.


Ochratoxins

Ochratoxins are produced as secondary metabolites of certain species of *Aspergillus* and *Penicillium* with *Aspergillus ochraceus* being the major producer [30]. Based on minor structural differences, it has been further subdivided into ochratoxin A (OTA), ochratoxin B (Mohamed, #123), and OTC, with OTA being the most important and toxic among all. OTA is highly carcinogenic, and the International Agency for Research on Cancer has classified it as Group 2B [69]. *Penicillium verrucosum* is considered a major OTA producer in cold and temperate climatic zones, *Penicillium verrucosum* whereas *Aspergillus carbonarius*, *A. ochraceus*, *A. niger*, and *A. melleus* are considered major OTA producers.

Ochratoxins contaminate different crops, including corn, maize, sorghum, barley, and rice, resulting in a

high chance of contamination in animal/poultry feed. Different regulatory levels are recommended for different raw materials and animal byproducts. According to the European Commission Recommendations (2006, 576), the MTL for complete feedstuff is 100 µg/kg feed [70]. Certain predisposing factors, such as hot and humid environments, poor pre-harvest/harvest conditions, and poor storage environments, enhance the production of OTA in feed.

Ruminants: OTA administered to young ruminants causes severe signs of depression, degeneration of the kidney, and polyuria, ultimately leading to the death of animals [71]. Ruminant microflora causes the degradation of amide bonds between ochratoxin- α and phenylalanine (Fig. 3), ultimately producing less toxic compounds [72]. In the rumen, 50% of the total OTA is degraded within the first 15 min, while 95% of the total OTA is degraded into less toxic ochratoxin- α and phenylalanine moieties by ruminal microflora within 4 h of ingestion. Owing to this phenomenon, OTA does not affect or penetrate vital organs of the body [45].



Ochratoxin A

Fig 3. Structural presentation of ochratoxin A (OTA)

Poultry: Poultry chicks are highly sensitive to the effects of OTA, which affects almost all vital organs of the body. Toxicopathological effects include reduced feed intake, body weight gain, dullness, and reduced attraction to feed. Serum biochemical parameters and hematological indices are also severely affected by exposure to OTA [73,74]. Immunosuppression has been extensively observed in this regard. In this regard, our recent reviews elaborate on experimental ochratoxicosis in poultry [24,69].

Fumonisins

Fumonisins are secondary metabolites of *Fusarium proliferatum* and *Fusarium moniliformis*. Apart from these two species, some *Alternaria*, especially *A. alternata* have been found to produce fumonisins [75]. There are twenty-eight (28) chemically different types of fumonisins (FA1, FA2, FB1, FB2, FB3, and FB4), among which fumonisins

B1 (FB1) is the most important and toxic form of fumonisins [76].

Ruminants: Regarding ruminants, fumonisins are tolerant to ruminal biodegradation; however, due to their low oral bioavailability, acute or chronic intoxication does not occur at the farm level [77]. However, the presence of very high levels in the feed can cause histopathological lesions in the kidneys and liver [78].

Poultry: The mechanism by which fumonisins cause toxicity in animals is thought to be the disruption of sphingolipid metabolism. It has been observed that fumonisins are the specific inhibitors of ceramide synthase enzyme which is needed for ceramide and complex sphingolipids synthesis. Because of this inhibition, a change in the sphingosine [7] to sphinganine [1] ratio occurs, and such an increased ratio has been observed in the tissues of turkeys, ducklings, and broilers exposed to FB1 in feed [79,80]. Turkeys and chicks were relatively resistant to the adverse effects of FB1, but mild to moderate toxicity was observed in turkeys and ducks fed FB1 at 75-400 mg/kg feed for 21 days. The observed changes included reduced body weight gain and hepatic damage, such as hepatic necrosis and biliary hyperplasia [81].

Trichothecenes

Trichothecenes are a diverse group of mycotoxins, the most important of which is deoxynivalenol [20] also known as vomitoxin. Other important members of this group are T-2 toxins, diacetoxyscirpenol (DAS), and scirpenol. They are produced by a variety of *Fusarium* species, while DON is produced primarily by *Fusarium roseum* and T-2 toxins are produced by *F. poae* and *F. sporotrichioides* under storage conditions [82].

Ruminants: Discussing about ruminants, microorganisms present within the ruminal fluid have distinct property of deacetylation and DAS is de-acetylated into monoacettoxyscirpenol [7] and scirpenetriol; and these products are less toxic as compared to their parent compounds [83]. DON occurs in excessive quantities within the concentrates, and it is also readily degraded within the rumen, but in animals suffering from ruminal acidosis, such degradation becomes incomplete, ultimately producing trichothecene-associated adverse effects in the animals [84]. However, in general, ruminants can easily tolerate 8.5 mg/g DON within the feed and is readily degraded within 6-24 h of its ingestion by the ruminants [85]. Furthermore, DON-associated lesions in animals include lesions within the gastrointestinal tract, vomiting, severe dermatitis, hemorrhage, and bloody diarrhea [86].

Poultry: Poultry chicks are also resistant to the adverse effects of DON to some extent, but the associated adverse effects include decreased feed efficiency, reduced body

weight gain, and poor efficiency when fed extremely high doses^[87].

APPROACHES/STRATEGIES FOR DETOXIFICATION/REDUCTION OF MYCOTOXINS

Three main approaches and strategies are generally adopted for the control and prevention of mycotoxins in animal and poultry feedstuff, which include physical, chemical, and biological methods.

Physical Methods

There are different physical methods for the control of mycotoxins, which unfortunately become impractical at a large scale, but such methods can efficiently help detoxify mycotoxins on a small scale. These methods include adsorption, irradiation, heating, solvent extraction, washing and separation.

Sorting and Separation

Grains are the major part of animal/poultry feed formulations^[88] and when these grains are significantly contaminated with mycotoxins, they appear as moldy, broken, discolored and not distributed uniformly in the contaminated cereals rather they cluster together forming 'mycotoxin pockets'^[89,90]. The separation and sorting of mycotoxin-contaminated grains are performed using sieving, aspiration, photoelectric separation, and image separation techniques^[91]. Hand sorting, dehulling, and flotation alone can remove 93%, 63%, and 51% of mycotoxins from white-shelled maize, respectively, whereas using these three methods in combination can remove nearly 98% of mycotoxins^[92]. Similarly, gravity separation and aspiration can reduce mycotoxins by 80%, but this also reduces wheat crop^[93]. Moreover, visual sorting strategies (optical viscosity and near-infrared spectroscopy) have been used to remove mycotoxins from maize and wheat^[90,94-96]. The limitations of these methods are that they are costly and limited to small-scale use.

Washing and Solvent Extraction

Mycotoxins can be easily removed by washing and solvent extraction because of their distinct fat-soluble and water-soluble properties. The floating method has the potential to decrease the concentrations of zearalenone, fumonisins, aflatoxins, and trichothecenes by 61%, 73%, 72%, and 69%, respectively^[92,96,97]. The best results obtained by the floating method can be achieved by adding NaCl and sucrose to water to attain the maximum output^[98]. When these physical techniques are used in combination to control mycotoxins, they provide better results than the individual techniques^[99]. Commonly used solvents for mycotoxin extraction include hexane, methanol, ethanol, and aqueous acetone^[100]. However, the

major disadvantages associated with washing and solvent extraction are that they cause the loss of nutrients and are costly, which limits their application at larger scales.

Heating

This method for the control of mycotoxins has been extensively used for the removal of different mycotoxins; however, AFB1 and FB1 can tolerate heat and require high temperatures (probably more than 150°C) for their decomposition^[101-104] ultimately making their decomposition difficult. Some studies have shown that thermal treatment up to 160°C for 20 min under a pressure of 10 MPa can reduce aflatoxins in rice by 80% from rice^[105] while heating barley at 220°C can destroy 90% of zearalenone (ZEN) and DON^[101]. Similarly, 150-200°C temperature can cause a 70% reduction in the concentration of FB1 in rice^[106]. However, the disadvantage of this method is the production of the Maillard reaction, formation of certain carcinogens such as acrylamide, and reduction of the nutritive value, thereby limiting the use of this method at larger scales^[105].

Decontamination by Irradiation

The irradiation process is usually divided into two forms: non-ionizing and ionizing. Non-ionizing processes involve microwaves, radio waves, visible light waves, and infrared waves, whereas the ionizing form includes ultraviolet rays, X-rays, electron beams, and gamma rays^[107]. Extensive research has been performed by many scientists to evaluate the degradation of different mycotoxins using different irradiation technologies. Table 4 presents different studies reporting the use of irradiation technologies for the degradation of mycotoxins in animal/poultry feed and their ingredients.

Using gamma irradiation: Gamma rays are electromagnetic waves produced as a result of decaying an unstable source such as radioactive isotopes. Gamma rays are preferred in the food industry because of their high reactivity and penetration power. However, certain factors are important in this regard as far as the usage of gamma rays for degradation of mycotoxin is concerned such as dose of radiation used, level of mycotoxin contamination, water content within feed and composition of matrix. Many studies have reported possible degradation of aflatoxins within feed at 5-10 KGy exposure to gamma rays^[112,122,123].

Gamma irradiation is undoubtedly gaining popularity due to its excellent results in finished food products, whereas Di Stefano et al.^[110] reported only up to 21% aflatoxin reduction from finished poultry feed, which suggests that this technique is not suitable for products containing high lipid and vitamin contents^[107].

Using electron beam: Irradiation through electron beams has shown promising results in the degradation of certain mycotoxins because of their short processing time,

Table 4. Degradation of mycotoxins by different irradiation techniques in animal/poultry feed and its ingredients					
Technique used	Feed type	Mycotoxin	Treatment Condition	Degradation Percentage	Reference
Gamma irradiation	Soybean	AFB1	10 kGy	62.20%	[108]
	Wheat	OTA	30.5 kGy	24%	[109]
	Commercial poultry feed	OTA	15 kGy	23.9%	[110]
	Commercial poultry feed	AFB1	15 kGy	18.2%	[110]
	Poultry feed	Aflatoxins	25 kGy	42.7%	[111]
	Maize feed	AFB1	10 kGy	94.5%	[112]
	Broiler feed	AFB1	6 kGy	89.53%	[113]
	Chicken liver	AFB1	10 kGy	25%	[114]
	Chicken liver	OTA	10 kGy	60%	[114]
	Poultry feed	OTA	4 kGy	100%	[115]
	Poultry feed	AFB1	6 kGy	100%	[116]
Electron beam	Corn	Zearalenone	50 kGy	71.1%	[117]
	Barley	Fusarium species	10 kGy	50-98%	[118]
	Wheat	DON	55.8 kGy	78.4%	[119]
Pulsed light	Rice bran	AFB1	0.52 J/cm per pulse for 15 sec	90.3%	[120]
Ultraviolet irradiation	Wheat	254 nm for 160 min	Aflatoxins	65-90%	[121]

dosage control, and low equipment costs. In the case of aflatoxins, this technology breaks down toxins into less toxic products, thereby reducing their toxicity potential of aflatoxins [124,125].

However, this technique has a lower degradation capacity than gamma irradiation. Moreover, Liu et al. [126] reported that this technique was not very efficient in degrading AFB1 from peanut meals, as this technique cannot be declared as a complete solution for degradation/decontamination purposes as far as mycotoxins are concerned.

Using ultraviolet irradiation: Over the past several decades, UV irradiation has been considered an effective technique for the destruction of several mycotoxins, particularly aflatoxins, owing to their photosensitivity. Being a non-thermal technique, UV irradiation depicts the benefits of being practical, cost-efficient and eco-friendly and it does not result in any toxic and/or waste product generation [127].

Ultraviolet (UV) irradiation has a strong penetration capacity through transparent and/or clear liquids, whereas its penetration capacity through solid materials is very limited, leading to its low decontamination ability for compact food products [128]. The disadvantage of this technique is that granular or opaque items should be presented in the form of a thin layer to achieve decontamination of mycotoxins by UV irradiation, thereby limiting its application at the field level for the degradation of mycotoxins in feed.

By photocatalysis: Recent advancements in the field have revealed that UV-visible irradiation, along with semiconducting photocatalysis, can efficiently degrade aflatoxins in a liquid matrix [129,130]. The most commonly used photocatalyst is titanium oxide (TiO_2), which is highly active under UV irradiation. Sun et al. [129] reported that AFB1 in methanol was efficiently degraded up to 95% within 120 min using this technique, while Xu et al. [130] reported up to 60.4% removal of AFB1 within 120 min through UV-vis irradiation. Similarly, DON degradation can also be achieved through photocatalytic techniques [131]. Although this is an efficient technique, less information is available on the safety and stability of photocatalysts.

By pulsed light: Pulsed light is an emerging non-thermal technique for decontaminating mycotoxins from food and feed. Pulsed light is an FDA-approved technique for efficient and rapid decontamination of different food products. In this advanced technique, short- and high-intensity broadband emission light is produced, which includes ultraviolet, visible, and infrared rays [132]. Moreau et al. [133] reported a 92.7% reduction in AFB1 in water using eight flashes of pulsed light, while Wang et al. [120] reported 75% AFB1 and 39.2% AFB2 reductions from raw rice samples using this technique. Pulsed light technology can also be used for the efficient degradation of OTA, ZEN, and DON [133].

Despite the great outcomes of this technology, further studies are needed to investigate the breakdown products

of pulsed light treatments. The design of cost-effective PL equipment is still needed, which can produce high UV output, so that this emerging technique can be effectively used at the industrial level.

By microwave heating: Electromagnetic waves with wavelengths ranging from 1m to 1 mm and frequencies of 300 MHz to 3000 MHz are used in microwave heating. It is a unique volumetric heating technology that efficiently converts electromagnetic field energy into thermal energy via the polarization effect of electromagnetic radiation [134]. Microwave heating is extensively used for heating, drying, extraction, and cooling of certain food products. Various studies have reported the use of this technology for non-thermal degradation of different mycotoxins. Flores et al. [135] studied the effects of microwave heating during alkaline cooling of aflatoxin-contaminated maize grains. They reported 36% AFB1 and 58% AFB2 reduction using microwave heating at 1650 W for 5.5 min.

Microwave manufacturers can customize equipment according to the needs of industrial and food product types. However, the non-uniform distribution of temperature during microwave heating is a challenge that could lead to hot/cold spot formation within food [136]. The presence of mycotoxins within the cold spot cannot be properly detoxified, whereas hot spots may lead to nutritional degradation of the feed. Further studies in this regard are required to ensure the proper distribution of temperature at all spots so that efficient degradation of mycotoxins can be achieved without disturbing the nutritional values of food and feed products.

By cold plasma: Plasma, often referred to as the fourth state of matter, is a highly energetic ionized gas that usually consists of ions, UV irradiation, electrons, and reactive nitrogen and oxygen species (RNS and ROS) [137]. Plasma is further categorized as cold or thermal. Additionally, it can be explained by the type of system generating it, such as dielectric barrier discharge (DBD), corona discharge (CD), and radio frequency plasma (RFP) and many others [138].

This latest technology has been used for the decontamination of different mycotoxins under ambient pressure and temperature conditions [139]. Aflatoxins were degraded using the DBD N₂-plasma technique in hazelnuts, where 70% of AFB1 was detoxified under 1150 W plasma treatment within 12 min [140]. Similarly, cold plasma technology has been reported to efficiently degrade various mycotoxins including OTA [141], DON [142,143], T2 [144], fumonisins [145] and citrinin [140]. Cold plasma technology has been proven to be an efficient technique for the decontamination of mycotoxins. However, this technique is still in the early stages of development, and further advanced research is required to optimize various

food products. Furthermore, the negative impact of plasma treatment on the nutritive value of different food materials needs to be addressed before the commercialization of this technology.

Adsorption

Some adsorbing substances have the potential to bind to mycotoxins and remove them from the gastrointestinal tract [146]. This technique is widely used and well understood, as it has fewer disadvantages than other available methods. The adsorbing agent used for the detoxification of mycotoxins should have some specific properties, including a high adsorption capacity for multiple mycotoxins, low binding efficacy for nutrients, and high safety and palatability [102].

The most commonly used adsorbents for mycotoxin detoxification are aluminosilicate minerals, including montmorillonites and aluminosilicates [146]. The adsorption ability depends on the structures of both the binding agent and the targeted mycotoxin [102]. Some studies have indicated that zeolite and bentonite clay can reduce AFB1 residues in the liver by 87% [147-150]. Similarly, Bentonite clay can decrease the bioavailability of AFB1 in the rumens of lactating animals [151,152]. Many studies have revealed that polar toxins, such as AFB1 and FB1, can be easily adsorbed by many aluminosilicates, which become ineffective against different non-polar mycotoxins [153,154]. Bentonites are considered good agents for the adsorption of mycotoxins because they are bioenvironment-friendly, have high adsorption efficiency, and are generally more economical than other agents [155-157]. Details of some of the adsorbing agents that are effective against mycotoxins are shown in *Table 5*.

Chemical Methods

Different chemical agents can be used to efficiently convert mycotoxins to less toxic or non-toxic compounds by destroying their structural makeup. Certain alkalines and ozone treatments are chemical methods that have proven beneficial in this regard [225,226]. *Fig. 4* shows a schematic flow of different chemical methods used for the control of mycotoxicosis.

Alkaline Treatment

Commonly used alkaline chemicals for the control of mycotoxins in moldy feed include sodium carbonate, potassium hydroxide, sodium hydroxide, and ammonia [226]. Base hydrolysis can open the lactone ring structure of AFB1, and the hydrolyzed product can then be removed by washing with water [227]. Treatment of various cereals with ammonia and hydroxide salts can remove almost 95% of the mycotoxins [228]. Although these treatments can nearly reduce the complete concentration of mycotoxins, the possible transformation of mycotoxins to other

Table 5. Details of different *In vitro* and *in vivo* studies reporting the use of different adsorbents (binders) against mycotoxicosis

Mycotoxin	Agent (Adsorbent)	Type of Study (and Efficacy)	Reference	Mycotoxin	Agent (Adsorbent)	Type of Study (and Efficacy)	Reference
Aflatoxins	Bentonite	<i>In Vitro</i>	[158]	Ochratoxins	Activated charcoal	Leghorn Chicks	[199]
	Activated charcoal	<i>In Vitro</i> Study	[159]		HSCAS	Chickens (Partial)	[200]
	Activated charcoal	Goats	[160]		Activated charcoal	<i>In Vitro</i>	[201]
	Activated charcoal	White Rock Chicks	[161]		Activated Charcoal	Pigs	[201]
	Activated charcoal	Broiler Chicks	[162]		HSCAS, Bento	Pigs	[201]
	HSCAS	Chickens	[163]		Cholestyramine	Pigs	[201]
	HSCAS	Broiler Chicks	[164]		Activated carbon	<i>In Vitro</i>	[202]
	Activated charcoal	Chickens	[165]		Diatomaceous earth	<i>In Vitro</i>	[189]
	HSCAS	Chickens	[165]		Bentonite	<i>In Vitro</i>	[195]
	HSCAS	Pigs	[166]		Bentonite	Broiler (Partial)	[203]
	HSCAS	Pigs	[167]	Fumonisins	Activated charcoal	<i>In Vitro</i>	[204]
	HSCAS	Pigs	[168]		Activated carbon	Rats	[205]
	Zeolite	<i>In Vitro</i>	[169]		Bentonite	<i>In Vitro</i>	[206]
	HSCAS	Pigs	[170]		Divinylbenzene-styrene polymer	Rats	[207]
	Zeolite	Broiler Chicks	[171]	Zearalenone	HSCAS	Minks	[208]
	HSCAS	Pigs	[172]		Cholestyramine	<i>In Vitro</i>	[209]
	HSCAS	<i>In Vitro</i> Study	[173]		Bentonite	Pigs	[210]
	HSCAS	Turkey Poulets	[174]		Maifanite	Pigs	[211]
	Activated charcoal	Rabbits (Partial)	[175]		Montmorillonite	Goat	[151]
	Zeolite	Domestic Fowls	[176]		Activated charcoal	<i>In Vitro</i>	[212]
	Bentonite, HSCAS	Broilers	[177]		Cholestyramine	<i>In Vitro</i>	[212]
	HSCAS	Turkey Poulets	[178]		Montmorillonite, Magnesium trisilicate, cholestyramine	<i>In Vitro</i>	[185]
	Activated charcoal	Minks	[179]		Organophilic montmorillonite	<i>In Vitro</i>	[213]
	HSCAS	Dairy Cows	[180]	DON	Polyvinylpyrrolidone	Pigs	[214]
	Aluminosilicate	Chicks	[181]		Activated carbon	<i>In Vitro</i>	[202]
	HSCAS	Wethers	[179]		Activated carbon	<i>In Vitro</i>	[215]
	HSCAS	Pigs	[182]	Trichothecenes	HSCAS	Turkey Poulets	[174]
	Zeolite	Chickens	[181]		HSCAS	Chicks	[153]
	Zeolite	Broiler Chicks	[183]		HSCAS	Pigs	[216]
	Calcium Bentonite	Pigs	[182]		Super-activated charcoal	Turkey Poulets	[187]
	HSCAS, Bentonite	Pigs	[184]		HSCAS	Broiler Chicks	[217]
	Montmorillonite	<i>In Vitro</i>	[185]		Inorganic clay	Broiler Chicks (Partial)	[188]
	Activated Charcoal	Turkey Poulets	[186]	T-2 toxins	Bentonite	Rats	[218]
	HSCAS	Turkey Poulets	[186]		Divinylbenzene-styrene polymer	Rats	[218]
	Super-activated charcoal	Broiler Chicks	[187]		Super-activated charcoal	Rats	[219]
	Inorganic clay	Broiler Chicks	[188]		Super-activated charcoal	Rats	[220]
	Diatomaceous earth	<i>In Vitro</i>	[189]		Activated charcoal	Swine	[221]
	Clinoptilote	Quail Chicks	[190]	Cyclopiazonic Acid	Acidic clay, neutral clay, clinoptilolite	<i>In Vitro</i>	[222]
	Aluminosilicate	<i>In Vitro</i>	[191]		Acidic clay, neutral clay, clinoptilolite	Broilers	[222]
	HSCAS, bentonite	Rats	[192]		Ergotamine	Montmorillonite	<i>In Vitro</i>
	HSCAS	<i>In Vitro</i>	[193]		Patulin	Activated charcoal	<i>In Vitro</i>
	HSCAS	Broilers	[193]				[224]
	Zeolite	Broiler Chicks	[194]				
	Bentonite	<i>In Vitro</i> Study	[195]				
	Bentonite	Broiler Chicks	[196]				
	Alumino silicate	<i>In Vitro</i>	[197]				
	Sodium bentonite	Poultry Chicks	[147]				
	Bentonite clay	Broilers	[149]				
	Zeolite	Poultry Chicks	[198]				
	Organo-clay composites	Broilers	[152]				

forms, such as masked mycotoxins, along with harmful side effects on the environment and food (changes in nutritional quality, texture, or flavor) are some of the disadvantages that make this method less desirable at larger scales [225].

Ozone Treatment

Oxidizing agents, such as sodium and ozone, play a role in detoxifying mycotoxins by modulating the structures of these secondary metabolites [229,230]. Ozone can degrade the




Fig 4. Flow diagram showing different chemical and biological methods used for the control of mycotoxicosis

FB1, AFs, ZEN and DON in various feeds of animals [231-234]. AFs can be reduced in corn and peanuts through ozone treatment [235-237]. Similarly, DON can be decomposed in corn and wheat using ozone [216,238-240]. Similarly, zearalenone can also be decomposed by treatment with varying concentrations of ozone [241]. Mycotoxins can also be degraded using other oxidizing agents, such as sodium hypochlorite [242,243]. However, the use of oxidizing agents for the detoxification of mycotoxins has some limitations, as these agents can change the physical and chemical composition of the feed, such as protein denaturation and lipid oxidation. Another disadvantage of using oxidizing agents is that they produce harmful chemicals during mycotoxin detoxification, making this method unsuitable for commercial use [230,235,236].

Biological Methods

Many physical and chemical techniques used to reduce the concentration of mycotoxins in feed have various limitations, as discussed above. However, the use of different biological methods for mycotoxin detoxification is necessary, as these methods have shown promising results in various studies [243,244]. *Fig. 4* shows a schematic flow of different biological methods used for the control of mycotoxicosis.

Microorganisms with Detoxification Activities

The use of different probiotics in animals promotes their development and growth and also improves the host's resistance against different diseases and metabolic disorders [245-249]. Certain beneficial microorganisms belonging to the category of probiotics are extensively used for the control of mycotoxicosis. These microorganisms play a significant role in maintaining normal bacterial balance within the GIT and are often used to tackle certain pathological abnormalities, including fungal modifications [250,251]. These beneficial organisms can bind mycotoxins and eliminate them from the gastrointestinal tract [252]. In addition, they also possess the ability to biologically

Table 6. Details of different *In vitro* studies reporting the biological degradation/detoxification of certain mycotoxins

Mycotoxin	Biological Strain/Type	Detoxification Rate [18]	Reference
AFB1	<i>Bacillus subtilis</i>	93.00%	[257]
	<i>Pseudomonas putida</i>	92.00%	[258]
	<i>Bacillus licheniformis</i>	95.70%	[259]
	<i>Bacillus shackletonii</i>	93.10%	[260]
	<i>Bacillus subtilis</i>	66.20%	[13]
	<i>Bacillus velezensis</i>	92.50%	[261]
	<i>Escherichia coli</i>	93.70%	[262]
	<i>S. cerevisiae</i>	68.00%	[244]
	<i>Aspergillus niger</i> RAF105	87.59%	[263]
DON	<i>Stenotrophomonas</i> sDOI:	99.00%	[264]
	<i>Aspergillus niger</i> FS11	97.64%	[265]
	Bacterial strain	100%	[266]
	Bacterial isolates	100.00%	[267]
	<i>Aspergillus</i> (NJA-1)	98.40%	[268]
	<i>Eggerthella</i> sDOI:	100.00%	[269]
	<i>Pseudomonas</i> sDOI: and <i>Lysobacter</i> sDOI:	100.00%	[270]
	<i>Devosia insulae</i>	85.00%	[271]
	Strain E3-39	100.00%	[272]
ZEN	<i>Bacterial consortium</i> C20	73.29%	[273]
	<i>Bacillus subtilis</i>	83.10%	[274]
	<i>Bacillus pumilus</i>	96.70%	[275]
FB1	<i>Bacillus natto/ pumilus</i>	97.70%	[276]
	<i>Bacillus subtilis</i>	88.00%	[272]
	<i>Bacillus</i> spDOI:	22%-50%	[277]
	<i>Saccharomyces cerevisiae</i>	100.00%	[278]
	Bacterial consortium	89.65%	[279]
	Strain NCB/ Bacterial consortium	100.00%	[280]

degrade mycotoxins, ultimately converting them into less toxic metabolites, thereby protecting animals/chicks from mycotoxin-associated lethal damage [253]. Microorganisms, including various species of *Lactobacillus*, *Lactococcus*, *Streptococcus* and *Bifidobacterium* possess antimutagenic, antifungal, and immunomodulatory effects in this regard [254-256]. Many studies have been conducted in this regard, and *Table 6* and *Table 7* summarize some probiotics associated with different types of mycotoxins *in vitro* and *in vivo*.

Biodegradation/Biotransformation by Degradation Enzymes

In addition to the use of different bacteria, fungi, and their byproducts for the degradation of mycotoxins, the use of certain biological enzymes is also gaining popularity.

Table 7. Details of different in vivo studies reporting biological degradation/transformation of certain mycotoxins			
Mycotoxin	Biological Species Used	Animal Model Used	Reference
Aflatoxins	<i>Saccharomyces cerevisiae</i>	Broilers	[281]
	Esterified glucomannan	Cows	[282]
	Manno-oligosaccharides	Wistar rats	[283]
	<i>Nocardia corynebacteroides</i>	Chicks	[284]
	Modified yeast extract	Cows	[285]
	Dried yeast culture	Sheep	[286]
	<i>Lactobacillus casei Shirota</i>	Wistar rats	[287]
	Modified yeast cell wall	Sheep	[288]
	<i>Lactobacillus rhamnosus</i> GAF01	Mice	[289]
	<i>Lactobacillus plantarum</i> MON03	Mice	[290]
	Yeast cell wall	Broilers	[291]
	<i>Pichia kudriavzevii</i>	Broilers	[292]
	<i>Lactobacillus plantarum</i>	Broilers	[293]
Ochratoxins	Esterified glucomannan	Broilers	[293]
	Yeast cell wall preparation (YCW)	Rats	[294]
	<i>Saccharomyces cerevisiae</i>	Ross male broilers	[295]
	<i>Lactobacillus paracasei</i>	Ross Broilers	[296]
	<i>Saccharomyces cerevisiae</i>	Ross Broilers	[296]
	Yeast sludge	Broilers	[297]
	<i>Lactobacillus kunkeei</i>	Male Rats	[298]
T-2	<i>Lactobacillus plantrum</i>	Male Rats	[298]
	<i>Saccharomyces cerevisiae lysate</i> (Masclaux-Daubresse, #1128)	Laying hens	[299]

These enzymes have been isolated from a wide range of microorganisms. These enzymes are also obtained from certain fungal species, but the isolation processes involving crushing of fungal mycelia are quite complicated, and their use on a larger scale is restricted. However, the use of enzymes isolated from different bacteria is promising for mycotoxin biodegradation [260]. **Table 8** shows the details of different degrading enzymes, along with the types of mycotoxins against which their efficacy has been reported.

Mycotoxin	Degrading Enzyme	Source	Reference
AFB ₁	<i>Trametes versicolor</i> Laccase enzyme	<i>Aspergillus niger</i>	[300]
	F42H2-dependent reductase enzyme	<i>Mycobacterium smegmatis</i>	[301]
	Aflatoxin-Oxidase	<i>Aspergillus tabescens</i>	[302]
	Myxobacteria aflatoxin degradation enzyme	<i>Myxococcus fulvus</i>	[303]
	Manganese peroxidase	<i>Phanerochaete sordida</i> YK 624	[304]
	Manganese peroxidase	<i>Pleurotus ostreatus</i>	[305]
	Bacillus aflatoxin-degrading enzyme	<i>Bacillus shackletonii</i> L7	[260]
DON	Cytochrome P450 system	<i>Sphingomonas</i> sDOI: strain KSM1	[306]
	Peroxidase	Extract of rice bran	[307]
	Aldo-keto reductase DepA/DepB	<i>Deeosia mutans</i> 17-2-E-8	[308]
	Quinone-dependent dehydrogenase, NADPH-dependent aldo/keto reductases	<i>Deeosia</i> strain D6-9	[309]
	Manganese peroxidase and Lignin peroxidase	<i>Flammulina velutipes</i>	[310]
ZEN	ZEN-specific lactonohydrolase	<i>Penicillium canescens</i> strain PCA-10	[311]
	Recombinant fusion enzyme (ZHDCP)	Zearalenone hydrolase (ZHD) and carboxypeptidase (CP)	[312]
FB ₁	Fumonisins carboxylesterase FumD	<i>Sphingopyxis</i> sDOI: MTA144	[313]

Nutritional Strategies

Certain nutritional strategies have also been adapted to nullify the adverse effects of mycotoxins in animals. Similarly, the use of certain plants and their extracts has been practiced since ancient times for the rectification of different ailments in both humans and animals [314-317]. These plants contain a variety of flavonoids, alkaloids, essential oils, and tannins, which enhance the body's defense system by mitigating several pathological and management issues in animals and humans [318,319]. The mycotoxin detoxification system can be modulated by nutritional measures. In animals, detoxification systems include ketoreductase, CYP450s and α -glutathione transferase, which can efficiently degrade mycotoxins [239]. Therefore, nutritional regulators may enhance the detoxification potential of the body [320]. Cysteine, glycine, and glutamate synthesize glutathione and help detoxify mycotoxins by forming glutathione. The addition of

Mycotoxins	Nutritional strategy	Animal Model	Reference
AFB1	β -carotene, canthaxanthin, lycopene	Rats	[328]
	Dietary butylated hydroxytoluene	White Turkey poult	[329]
	Vitamin C	<i>Labeo rohita</i>	[330]
	Silymarin-phospholipid complex	Broiler	[331]
	Vitamin C & E	Rabbits	[332]
	Sea buckthorn berries	Broiler	[333]
	Alpha-lipoic acid	Broilers	[334]
	Resveratrol	Broilers	[335]
	Curcumin, quercetin, resveratrol	<i>In vitro</i>	[336]
	Selenium	Cobb male broilers	[280]
DON	Curcumin	Chickens	[337]
	Vitamin E	Leghorn cockerels	[58]
	Vitamin E & C, selenium	Wistar rats	[338]
FB1	L-Arginine	Pig	[339]
	Selenium	<i>In vitro</i>	[276]
	Isoflavones	Rats	[340]
OTA	Vitamin E	rabbits	[341]
	Curcumin/silymarin	<i>In vitro</i>	[342]
	Vitamin C	Mice	[343]
	Vitamin E & C	Broilers	[344]
	Retinol, ascorbic acid, α -tocopherol	Mice	[345]
	Coenzyme Q10, L-carnitine, Zn, Mg	Mice	[346]
	Vitamin C	Nile Tilapia	[347]
	L-carnitine	Leghorn cockerels	[66]
	Silymarin	Leghorn cockerels	[67]
T-2 toxin	Curcumin	<i>In vitro</i>	[342]
	Ducks	Curcumin	[348]
	Vitamin C & E, selenium	Wistar rats	[338]
Zearalenone	Vitamin E & C	Broiler	[344]
	Broilers	Lycopene	[349]
	Retinol, ascorbic acid, α -tocopherol	Mice	[345]
	Vitamin E	<i>In vitro</i>	[350]
	Vitamin C	Weaning piglets	[351]
	Silymarin	Rats	[352]
	Vitamin C	piglets	[353]




Fig 5. Schematic flow showing different nutritional strategies used for the control of mycotoxicosis

specific nutrients and prebiotics to mitigate the damaging effects of pathogens and toxins is another benefit [321-324]. Cytotoxicity occurs due to oxidants produced within the body by mycotoxins, and adding antioxidants can improve the ability of organisms to fight against mycotoxin toxicity. In this regard, selenium and vitamins (C and E) have proven beneficial as they act as superoxide anion scavengers [325-327]. Table 9 describes some nutritional strategies reported in this regard while Fig. 5 shows the schematic flow of use of different nutritional strategies used for the control of mycotoxicosis.

CONCLUSION

The presence of mycotoxins in feed is an unavoidable problem worldwide. This review summarizes a number of strategies to reduce mycotoxicosis, including physical methods (separation, washing, adsorption, heating, and irradiation), chemical strategies (oxidizing agents and basis), biological methods (enzymes and probiotics), and nutritional regulation strategies. Each of these approaches can be used practically, along with its own advantages and disadvantages. However, with the growing awareness of environmental protection, as well as feed and food safety, there is a growing expectation for more green and innovative technologies to control mycotoxin contamination.

Future Perspectives

More advanced techniques should be adapted for the control of mycotoxicosis in poultry. As there is a paradigm shift to the use of natural ingredients for the treatment globally, measures should be taken for the large-scale implementation of biological strategies to avoid physical and chemical strategies-associated disadvantages at farm levels. Widescale commercialization should be done to mitigate mycotoxicosis at farm (both dairy and poultry) levels.

DECLARATIONS

Availability of Data and Materials: All the generated data are included in the manuscript.

Acknowledgements: None

Competing Interests: The authors declared that there is no competing interest.

Declaration of Generative Artificial Intelligence (AI): The article and/or tables and figures were not written/created by AI and AI assisted technologies.

Author Contributions: AK, ZA, and AA: Conceptualization, methodology, and data curation; MKS and RZA: Data extraction and editing. MUI: Editing; MUR and BM: Revised and proof-read. All authors read and approved the final manuscript.

REFERENCES

1. Gurikar C, Shivaprasad D, Sabillón L, Gowda N N, Siliveru K: Impact of mycotoxins and their metabolites associated with food grains. *Grain Oil Sci Technol*, 6 (1): 1-9, 2023. DOI: 10.1016/j.gaost.2022.10.001
2. Furian AF, Fighera MR, Royes LFF, Oliveira MS: Recent advances in assessing the effects of mycotoxins using animal models. *Cur Opin Food Sci*, 47:100874, 2022. DOI: 10.1016/j.cofs.2022.100874
3. Fumagalli F, Ottoboni M, Pinotti L, Cheli F: Integrated mycotoxin management system in the feed supply chain: Innovative approaches. *Toxins (Basel)*, 13 (8): 572, 2021. DOI: 10.3390/toxins13080572
4. Luo S, Du H, Kebede H, Liu Y, Xing F: Contamination status of major mycotoxins in agricultural product and food stuff in Europe. *Food Cont*, 127:108120, 2021. DOI: 10.1016/j.foodcont.2021.108120
5. Marlière CA, Pimenta RCJ, Cunha AC: Fumonisin as a risk factor to esophageal cancer: A review. *Appl Cancer Res*, 29, 102-105, 2009.
6. Benkerroum N: Aflatoxins: Producing-molds, structure, health issues and incidence in Southeast Asian and Sub-Saharan African countries. *Int J Environ Res Public Health*, 17 (4):1215, 2020. DOI: 10.3390/ijerph17041215
7. Wang M, Li L, Hou C, Guo X, Fu H: Building and health: Mapping the knowledge development of sick building syndrome. *Buildings*, 12:287, 2022. DOI: 10.3390/buildings12030287
8. Goudarzi G, Reshadatian N: The study of effective factors in sick building syndrome related to fungi and its control methods. *Res Eng*, 23:102703, 2024. DOI: 10.1016/j.rineng.2024.102703
9. Soriano JM, Rubini A, Morales-Suarez-Varela M, Merino-Torres JE, Silvestre D: Aflatoxins in organs and biological samples from children affected by kwashiorkor, marasmus and marasmic-kwashiorkor: A scoping review. *Toxicon*, 185, 174-183, 2020. DOI: 10.1016/j.toxicon.2020.07.010
10. Stoyanov GS, Kobakova I, Petkova L, Dzhenkov DL, Popov H: Balkan endemic nephropathy: An autopsy case report. *Cureus*, 13 (1):e12415, 2021. DOI: 10.7759/cureus.12415
11. Sadashivanavar V, Madalageri M, Pai KSR, Sharma H, Halagali P, Seenivasan R, Tippavajhala VK, Somanna P, Noman AI: Cardiovascular impacts of foodborne toxins. In, *Physiological Perspectives on Food Safety: Exploring the Intersection of Health and Nutrition*. 351-375, Cham: Springer Nature Switzerland, 2025.
12. Lumsangkul C, Tso KH, Fan YK, Chiang HI, Ju JC: Mycotoxin fumonisin B1 interferes sphingolipid metabolism and neural tube closure during early embryogenesis in brown tsaiya ducks. *Toxins*, 13 (11):743, 2021. DOI: 10.3390/toxins13110743
13. Cai P, Zheng H, She J, Feng N, Zou H, Gu J, Yuan Y, Liu X, Liu Z, Bian J: Molecular mechanism of aflatoxin-induced hepatocellular carcinoma derived from a bioinformatics analysis. *Toxins*, 12:203, 2020. DOI: 10.3390/toxins12030203
14. Jagels A, Stephan F, Ernst S, Lindemann V, Cramer B, Hübner F, Humpf HU: Artificial vs natural *Stachybotrys* infestation - Comparison of mycotoxin production on various building materials. *Indoor Air*, 30 (6): 1268-1282, 2020. DOI: 10.1111/ina.12705
15. Leslie JF, Moretti A, Mesterházy Á, Ameye M, Audenaert K, Singh PK, Richard-Forget F, Chulze SN, Ponte EMD, Chala A, Battilani P: Key global actions for mycotoxin management in wheat and other small grains. *Toxins*, 13 (10):725, 2021. DOI: 10.3390/toxins13100725
16. Khatoon A, Khan MZ, Abidin Z, Saleemi MK, Oguz H, Gul S T, Abbas RZ, Ali A, Bhatti SA: Aflatoxin B1-associated oxidative stress along with toxicopathological and immunological alterations is efficiently counteracted by dietary supplementation of distillery yeast sludge in broilers. *Mycotox Res*, 40 (4): 615-629, 2024. DOI: 10.1007/s12550-024-00549-y
17. Sarwar MK, Ijaz S, Javed N, Akbar N: Molecular and analytical approaches based characterization of aflatoxins producing *Aspergillus* species affecting groundnut. *Pak J Agric Sci*, 61 (3): 931-940, 2024.
18. Shabeer S, Asad S, Jamal A, Ali A: Aflatoxin contamination, its impact and management strategies: An updated review. *Toxins*, 14 (5):307, 2022. DOI: 10.3390/toxins14050307
19. Abidin Z, Khatoon A, Numan MJ: Mycotoxins in broilers: Pathological alterations induced by aflatoxins and ochratoxins, diagnosis and determination, treatment and control of mycotoxicosis. *World Poult Sci J*, 67 (3): 485-496, 2011. DOI: 10.1017/S0043933911000535
20. Cui R, Pan A, Wang T, Liang Y, Yu HF: Aflatoxin B1 in animals: metabolism and immunotoxicity. *Pak Vet J*, 2025 (Early Online). DOI: 10.29261/pakvetj/2025.207
21. El-Hamaky AM, Hassan AA, Wahba AK, El-Mosalamy MM: Influence of copper and zinc nanoparticles on genotyping characterizations of multi-drug resistance genes for some calf pathogens. *Int J Vet Sci*, 12: 309-317, 2023. DOI: 10.47278/journal.ijvs/2022.195
22. Khashan SA, Khashan BA, Thalij KM, Konca Y: The effect of nanochitosan in reducing the toxicity of aflatoxin B1 and fumonisin B1 in broilers. *Pak Vet J*, 45 (1): 268-276, 2025. DOI: 10.29261/pakvetj/2025.116
23. Khan MA, Khan IA, Tahir AH, Shahid MA, Nazish N, Zafar MA, Bhatti SA, Pasha RH, Abbas Y, Sadiq S, Jamil B: Isolation and identification of deleterious fungi associated with stored grains and cattle feedstuff of Potohar region of Pakistan. *Pak Vet J*, 44 (3): 861-867, 2024. DOI: 10.29261/pakvetj/2024.189
24. Khatoon A, Abidin Z: Mycotoxicosis - diagnosis, prevention and control: Past practices and future perspectives. *Toxin Rev*, 39, 99-114, 2020. DOI: 10.1080/15569543.2018.1485701
25. Mengesha G, Bekele T, Ashagrie H, Woldegiorgis AZ: Level of aflatoxins in dairy feeds, poultry feeds, and feed ingredients produced by feed factories in Addis Ababa, Ethiopia. *Mycotox Res*, 40: 309-318, 2024. DOI: 10.1007/s12550-024-00531-8
26. Aboagye-Nuamah F, Kwoseh CK, Maier DE: Toxigenic mycoflora, aflatoxin and fumonisin contamination of poultry feeds in Ghana. *Toxicon*, 198, 164-170, 2021. DOI: 10.1016/j.toxicon.2021.05.006
27. Meijer N, Kleter G, de Nijs M, Rau ML, Derkx R, van der Fels-Klerx HJ: The aflatoxin situation in Africa: Systematic literature review. *Comp Rev Food Sci Food Saf*, 20, 2286-2304, 2021. DOI: 10.1111/1541-4337.12731
28. Priya MS, Jagadeeswaran A, Natarajan A: Detection of aflatoxin B1 (AFB1) in common ingredients of poultry and broiler feed under different seasons. *J Livest Sci*, 14, 163-168, 2023. DOI: 10.33259/JLivestSci.2023.163-168
29. Negash D: Animal feed safety: Cases and approaches to identify the contaminants and toxins. *Safety*, 4, 1-8, 2020.
30. Abidin Z, Khatoon A, Numan M: Mycotoxins in broilers: Pathological alterations induced by aflatoxins and ochratoxins, diagnosis and determination, treatment and control of mycotoxicosis. *World Poult Sci J*, 67 (3): 485-496, 2011. DOI: 10.1017/S0043933911000535
31. Olatoye O, Aiyedun J, Oludairo O: Incidence of aflatoxin B1 in commercial poultry feed and tissues of broiler chickens in ibadan, Nigeria. *Sahel J Vet Sci*, 17 (2): 13-18, 2020
32. Kassaw TS, Megerssa YC, Woldemariyam FT: Occurrence of aflatoxins in poultry feed in selected chicken rearing villages of bishoftu Ethiopia. *Vet Med (Auckl)*, 13: 277-286, 2022. DOI: 10.2147/VMRR.S384148
33. Bintvihok A, Kositcharoenkul S: Effect of dietary calcium propionate on performance, hepatic enzyme activities and aflatoxin residues in broilers fed a diet containing low levels of aflatoxin B1. *Toxicon*, 47 (1): 41-6, 2006. DOI: 10.1016/j.toxicon.2005.09.009
34. Denli M, Blandon JC, Guynot ME, Salado S, Perez JF: Effects of dietary

- AflaDetox on performance, serum biochemistry, histopathological changes, and aflatoxin residues in broilers exposed to aflatoxin B (1). *Poult Sci.* 88 (7): 1444-1451, 2009. DOI: 10.3382/ps.2008-00341
- 35. Hussain Z, Khan MZ, Khan A, Javed I, Saleemi MK, Mahmood S, Asi MR:** Residues of aflatoxin B1 in broiler meat: effect of age and dietary aflatoxin B1 levels. *Food Chem Toxicol.* 48 (12): 3304-3307, 2010. DOI: 10.1016/j.fct.2010.08.016
- 36. Khan MZ, Hameed MR, Hussain T, Khan A, Javed I, Ahmad I, Hussain A, Saleemi MK, Islam NU:** Aflatoxin residues in tissues of healthy and sick broiler birds at market age in Pakistan: A one year study. *Pak Vet J.* 33 (4): 423-427, 2013.
- 37. Fowler J, Li W, Bailey C:** Effects of a calcium bentonite clay in diets containing aflatoxin when measuring liver residues of aflatoxin B1 in starter broiler chicks. *Toxins*, 7 (9): 3455-3464, 2015. DOI: 10.3390/toxins7093455
- 38. Aljazzar A, El-Ghareeb WR, Darwish WS, Abdel-Raheem SM, Ibrahim AM:** Content of total aflatoxin, lead, and cadmium in the bovine meat and edible offal: Study of their human dietary intake, health risk assessment, and molecular biomarkers. *Environ Sci Poll Res.* 28 (43): 61225-61234, 2021. DOI: 10.1007/s11356-021-12641-2
- 39. Milicevic D, Jovanovic M, Matekalo-Sverak V, Radicevic T, Petrovic MM, Lilic S:** A survey of spontaneous occurrence of ochratoxin A residues in chicken tissues and concurrence with histopathological changes in liver and kidneys. *J Environ Sci Health C Environ Carcinog Ecotoxicol Rev.* 29 (2): 159-175, 2011. DOI: 10.1080/10590501.2011.577687
- 40. Pozzo L, Cavallarin L, Antoniazzi S, Guerre P, Biasibetti E, Capucchio MT, Schiavone A:** Feeding a diet contaminated with ochratoxin A for broiler chickens at the maximum level recommended by the EU for poultry feeds (0.1 mg/kg). 2. Effects on meat quality, oxidative stress, residues and histological traits. *J Anim Physiol Anim Nutr.* 97, 23-31, 2013. DOI: 10.1111/jpn.12051
- 41. Qu D, Huang X, Han J, Man N:** Efficacy of mixed adsorbent in ameliorating ochratoxicosis in broilers fed ochratoxin A contaminated diets. *It J Anim Sci.* 16 (4): 573-579, 2017. DOI: 10.1080/1828051X.2017.1302822
- 42. Hussein MA, Gherbawy Y:** Genotypic identification of ochratoxigenic Aspergilli that contaminated beef luncheon and their protease activity. *Rendiconti Lincei. Scienze Fisiche e Naturali.* 30 (4): 767-773, 2019. DOI: 10.1007/s12210-019-00845-1
- 43. Yue S, Li X, Qian J, Du J, Liu X, Xu H, Liu H, Chen X:** Impact of enzymatic hydrolyzed protein feeding on rumen microbial population, blood metabolites and performance parameters of lactating dairy cows. *Pak Vet J.* 43 (4): 804-808, 2023. DOI: 10.29261/pakvetj/2023.081
- 44. Agustin F, Pazla R, Jamarun N, Suryadi H:** Exploring the impact of processed cassava peel on microbial dynamics and *in vitro* nutrient digestibility in ruminant diets. *Int J Vet Sci.* 13 (4): 463-470, 2024. DOI: 10.47278/journal.ijvs/2023.119
- 45. Abidin Z, Khatoon A:** Ruminal microflora, mycotoxin inactivation by ruminal microflora and conditions favouring mycotoxicosis in ruminants: A review. *Int J Vet Sci.* 1, 36-43, 2012.
- 46. Min L, Fink-Gremmels J, Li D, Tong X, Tang J, Nan X, Yu Z, Chen W, Wang G:** An overview of aflatoxin B1 biotransformation and aflatoxin M1 secretion in lactating dairy cows. *Anim Nutr.* 7 (1): 42-48, 2021. DOI: 10.1016/j.aninu.2020.11.002
- 47. Jiang, Y, Ogunade IM, Vyas D, Adesogan AT:** Aflatoxin in dairy cows: toxicity, occurrence in feedstuffs and milk and dietary mitigation strategies. *Toxins*, 13:283, 2021
- 48. Keyl AC, Booth AN:** Aflatoxin effects in livestock. *J Am Oil Chem Soc.* 48 (10): 599-604, 1971. DOI: 10.1007/BF02544571
- 49. Pier A, Richard J, Thurston J:** Effects of mycotoxins on immunity and resistance of animals. *Nat Toxins*, 691-699, 1980. DOI: 10.1016/B978-0-08-024952-0.50087-9
- 50. Richard JL, Pier AC, Stubblefield RD, Shotwell OL, Lyon RL, Cutlip RC:** Effect of feeding corn naturally contaminated with aflatoxin on feed efficiency, on physiologic, immunologic, and pathologic changes, and on tissue residues in steers. *Am J Vet Res.* 44 (7): 1294-1299, 1983.
- 51. Helferich WG, Baldwin RL, Hsieh DP:** [14C]-aflatoxin B1 metabolism in lactating goats and rats. *J Anim Sci.* 62 (3): 697-705, 1986. DOI: 10.2527/jas1986.623697x
- 52. D'Angelo A, Bellino C, Alborali GL, Biancardi A, Borrelli A, Capucchio MT, Catalano D, Dellaferreira G, Maurella C, Cagnasso A:** Neurological signs associated with aflatoxicosis in Piedmontese calves. *Vet Rec.* 160 (20): 698-700, 2007. DOI: 10.1136/vr.160.20.698
- 53. Fernandez A, Hernandez M, Verde MT, Sanz M:** Effect of aflatoxin on performance, hematology, and clinical immunology in lambs. *Can J Vet Res.* 64 (1): 53-58, 2000.
- 54. Gowda N, Suganthi R, Malathi V, Raghavendra A:** Efficacy of heat treatment and sun drying of aflatoxin-contaminated feed for reducing the harmful biological effects in sheep. *Anim Feed Sci Technol.* 133 (1-2): 167-175, 2007. DOI: 10.1016/j.anifeedsci.2006.08.009
- 55. Tripathi M, Mondal D, Karim S:** Growth, haematology, blood constituents and immunological status of lambs fed graded levels of animal feed grade damaged wheat as substitute of maize. *J Anim Physiol Anim Nutr.* 92 (1): 75-85, 2008. DOI: 10.1111/j.1439-0396.2007.00712.x
- 56. Masoero F, Moschini M, Gallo A, Diaz D:** *In vivo* release of aflatoxin B1 bound to different sequestering agents in dairy cows. *It J Anim Sci.* 6 (Suppl.-1): 315-317, 2007. DOI: 10.4081/ijas.2007.1s.315
- 57. Guo H, Wang P, Liu C, Chang J, Yin Q, Wang L, Jin S, Zhu Q, Lu F:** Compound mycotoxin detoxifier alleviating aflatoxin B(1) toxic effects on broiler growth performance, organ damage and gut microbiota. *Poult Sci.* 102 (3):102434, 2023. DOI: 10.1016/j.psj.2022.102434
- 58. Saleemi MK, Raza A, Khatoon A, Zubair M, Murtaza B, Jamil M, Imran M, Muhammad F, Zubair K, Bhatti SA:** Toxic effects of aflatoxin B1 on hematobiochemical and histopathological parameters of juvenile white Leghorn male birds and their amelioration with vitamin E and *Moringa oleifera*. *Pak Vet J.* 43 (3): 405-411, 2023. DOI: 10.29261/pakvetj/2023.053
- 59. Khatoon A, Amin A, Majeed S, Gul ST, Arshad MI, Saleemi MK, Ali A, Abbas RZ, Bhatti SA:** Dietary Chlorella vulgaris mitigates aflatoxin B1 toxicity in broiler chicken: Toxicopathological, hematobiochemical and immunological perspectives. *Toxicon.* 251:108127, 2024. DOI: 10.1016/j.toxicon.2024.108127
- 60. Zentai A, Jozwiak A, Süth M, Farkas Z:** Carry-over of aflatoxin B1 from feed to cow milk—A review. *Toxins*, 15:195, 2023. DOI: 10.3390/toxins15030195
- 61. Alnuimy A:** Toxopathological and histopathological effects of aflatoxins. *Mag Al-Kufa Univ Biol.* 16, 25-43, 2024. DOI: 10.36320/ajb/v16.i1.13241
- 62. Muaz K, Riaz M, Oliveira CAFD, Akhtar S, Ali SW, Nadeem H, Park S, Balasubramanian B:** Aflatoxin M1 in milk and dairy products: Global occurrence and potential decontamination strategies. *Toxin Rev.* 41, 588-605, 2022. DOI: 10.1080/15569543.2021.1873387
- 63. Abedullah A, Kouser S, Badar H, Ibrahim MNM:** Consumer demand for aflatoxin-free raw milk in Pakistan. *J Anim Plant Sci.* 33, 125-134, 2023. DOI: 10.36899/JAPS.2023.1.0602
- 64. Bervis N, Loran S, Juan T, Carramiñana JJ, Herrera A, Arino A, Herrera M:** Field monitoring of aflatoxins in feed and milk of high-yielding dairy cows under two feeding systems. *Toxins*, 13:201, 2021. DOI: 10.3390/toxins13030201
- 65. Abidin Z, Khan MZ, Khatoon A, Saleemi MK, Khan A:** Protective effects of L-carnitine upon toxicopathological alterations induced by ochratoxin A in white Leghorn cockerels. *Toxin Rev.* 35 (3-4): 157-164, 2016. DOI: 10.1080/15569543.2016.1219374
- 66. Abidin Z, Khan MZ, Khatoon A, Saleemi MK, Khan A, Javed I:** Ameliorative effects of L-carnitine and vitamin E (alpha-tocopherol) on haematological and serum biochemical parameters in white Leghorn cockerels given ochratoxin A contaminated feed. *Br Poult Sci.* 54 (4): 471-477, 2013. DOI: 10.1080/00071668.2013.796509
- 67. Khatoon A, Khan MZ, Khan A, Saleemi MK, Javed I:** Amelioration of ochratoxin A-induced immunotoxic effects by silymarin and vitamin E in white Leghorn cockerels. *J Immunotoxicol.* 10 (1): 25-31, 2013. DOI: 10.3109/1547691X.2012.686533
- 68. Khatoon A, Khan MZ, Abidin Z, Khan A, Saleemi M:** Mitigation potential of distillery sludge against ochratoxin A induced immunological alterations in broiler chicks. *World Mycotox J.* 10 (3): 255-262, 2017.

- 69.** Khatoon A, Abidin Z: An extensive review of experimental ochratoxosis in poultry: I. Growth and production parameters along with histopathological alterations. *World Poult Sci J*, 74 (4): 627-646, 2018. DOI: 10.1017/S0043933918000685
- 70.** Abidin Z, Khatoon A, Arooj N, Hussain S, Ali S, Manzoor AW, Saleemi MK: Estimation of ochratoxin A in poultry feed and its ingredients with special reference to temperature conditions. *Br Poult Sci*, 58 (3): 251-255, 2017. DOI: 10.1080/00071668.2017.1293797
- 71.** Longobardi C, Ferrara G, Andretta E, Montagnaro S, Damiano S, Ciarcia R: Ochratoxin A and kidney oxidative stress: The role of nutraceuticals in veterinary medicine - A review. *Toxins*, 14:398, 2022. DOI: 10.3390/toxins14060398
- 72.** Xu H, Wang L, Sun J, Wang L, Guo H, Ye Y, Sun X: Microbial detoxification of mycotoxins in food and feed. *Crit Rev Food Sci Nutr*, 62, 4951-4969, 2022. DOI: 10.1080/10408398.2021.1879730
- 73.** Khatoon A, Khan MZ, Khan A, Javed I: Toxicopathological and serum biochemical alterations induced by ochratoxin A in broiler chicks and their amelioration by locally available bentonite clay. *Pak J Agric Sci*, 53 (4): 977-984, 2016.
- 74.** Khatoon A, Nawaz MY, Mehboob G, Saleemi MK, Gul ST, Abbas RZ, Ijaz MU, Murtaza B, Bhatti SA, Abidin Z: Unraveling the combined deleterious effects of ochratoxin a and atrazine upon broiler's health: Toxicopathological, serum biochemical and immunological perspectives. *Toxicon*, 236:107327, 2023. DOI: 10.1016/j.toxicon.2023.107327
- 75.** Meng J, Li R, Huang Q, Guo D, Fan K, Zhang J, Zhu X, Wang M, Chen X, Nie D, Cao C: Survey and toxicogenic abilities of *Aspergillus*, *Fusarium*, and *Alternaria* fungi from wheat and paddy grains in Shanghai, China. *Front Plant Sci*, 14:1202738, 2023. DOI: 10.3389/fpls.2023.1202738
- 76.** Chen J, Wen J, Tang Y, Shi J, Mu G, Yan R, Cai J, Long M: Research progress on fumonisins B1 contamination and toxicity: A review. *Molecules*, 26 (17):5238, 2021. DOI: 10.3390/molecules26175238
- 77.** Mostrom MS, Jacobsen BJ: Ruminant mycotoxicosis: An update. *Vet Clin North Am Food Anim Pract*, 36 (3): 745-774, 2020. DOI: 10.1016/j.cvfa.2020.08.011
- 78.** Loh ZH, Ouwerkerk D, Klieve AV, Hungerford NL, Fletcher MT: Toxin degradation by rumen microorganisms: A Review. *Toxins (Basel)*, 12 (10):664, 2020. DOI: 10.3390/toxins12100664
- 79.** Guerre P, Gilleron C, Matard-Mann M, Nyvall Collen P: Targeted sphingolipid analysis in heart, gizzard, and breast muscle in chickens reveals possible new target organs of fumonisins. *Toxins*, 14:828, 2022. DOI: 10.3390/toxins14120828
- 80.** Guerre P, Travel A, Tardieu D: Targeted analysis of sphingolipids in turkeys fed fusariotoxins: First evidence of key changes that could help explain their relative resistance to Fumonisin toxicity. *Int J Mol Sci*, 23 (5):2512, 2022. DOI: 10.3390/ijms23052512
- 81.** Galli GM, Griss LG, Fortuoso BF, Silva AD, Fracasso M, Lopes TF, Schetinger MRS, Gundel S, Ourique AF, Carneiro C, Mendes RE: Feed contaminated by fumonisins (*Fusarium* spp.) in chicks has a negative influence on oxidative stress and performance, and the inclusion of curcumin-loaded nanocapsules minimizes these effects. *Microb Pathog*, 148:104496, 2020. DOI: 10.1016/j.micpath.2020.104496
- 82.** Wang J, Zhang M, Yang J, Yang X, Zhang J, Zhao Z: Type A trichothecene metabolic profile differentiation, mechanisms, biosynthetic pathways, and evolution in *Fusarium* species - A mini review. *Toxins*, 15:446, 2023. DOI: 10.3390/toxins15070446
- 83.** Kibugu J, Munga L, Mburu D, Maloba F, Auma JE, Grace D, Lindahl JF: Dietary mycotoxins: an overview on toxicokinetics, toxicodynamics, toxicity, epidemiology, detection, and their mitigation with special emphasis on aflatoxicosis in humans and animals. *Toxins*, 16:483, 2024. DOI: 10.3390/toxins16110483
- 84.** Dong JN, Zhao ZK, Wang ZQ, Li SZ, Zhang YP, Sun Z, Qin GX, Zhang XF, Zhao W, Aschalew ND, Wang T: Impact of deoxynivalenol on rumen function, production, and health of dairy cows: Insights from metabolomics and microbiota analysis. *J Hazard Mater*, 465:133376, 2024. DOI: 10.1016/j.jhazmat.2023.133376
- 85.** Zhang F, Wu Q, Wang W, Guo S, Li W, Lv L, Chen H, Xiong F, Liu Y, Chen Y, Li S: Inhibitory effect mediated by deoxynivalenol on rumen fermentation under high-forage substrate. *Ferment*, 8:369, 2022. DOI: 10.3390/fermentation8080369
- 86.** Cope RB: Trichothecenes. In, Veterinary Toxicology. Academic Press, pp. 1083-1093, 2025.
- 87.** Riahi I, Marquis V, Ramos AJ, Brufau J, Esteve-Garcia E, Perez-Vendrell AM: Effects of deoxynivalenol-contaminated diets on productive, morphological, and physiological indicators in broiler chickens. *Animals*, 10:1795, 2020. DOI: 10.3390/ani10101795
- 88.** Anwar U, Yousaf M, Mirza MA, Aziz-ur-Rahman M: Impact of stored wheat-based feed on gut morphology, digesta viscosity and blood metabolites of broiler chickens. *Pak Vet J*, 43 (1): 179-183, 2023.
- 89.** Yu J, Pedroso IR: Mycotoxins in cereal-based products and their impacts on the health of humans, livestock animals and pets. *Toxins*, 15:480, 2023. DOI: 10.3390/toxins15080480
- 90.** Wenndt AJ, Sudini HK, Mehta R, Pingali P, Nelson R: Spatiotemporal assessment of post-harvest mycotoxin contamination in rural North Indian food systems. *Food Cont*, 126:108071, 2021. DOI: 10.1016/j.foodcont.2021.108071
- 91.** Wasti S, Sah N, Singh AK, Lee CN, Jha R, Mishra B: Dietary supplementation of dried plum: A novel strategy to mitigate heat stress in broiler chickens. *J Anim Sci Biotechnol*, 12:58, 2021. DOI: 10.1186/s40104-021-00571-5
- 92.** Liu M, Zhao L, Gong G, Zhang L, Shi L, Dai J, Han Y, Wu Y, Khalil MM, Sun L: Remediation strategies for mycotoxin control in feed. *J Anim Sci Biotechnol*, 13:19, 2022. DOI: 10.1186/s40104-021-00661-4
- 93.** Ismail AM, Raza MH, Zahra N, Ahmad R, Sajjad Y, Khan SA: Aflatoxins in wheat grains: detection and detoxification through chemical, physical, and biological means. *Life*, 14:535, 2024. DOI: 10.3390/life14040535
- 94.** Pascale M, Logrieco AF, Lippolis V, De Girolamo A, Cervellieri S, Lattanzio VM, Ciasca B, Vega A, Reichel M, Graeber M, Slettengren K: Industrial-scale cleaning solutions for the reduction of *Fusarium* toxins in maize. *Toxins*, 14:728, 2022. DOI: 10.3390/toxins14110728
- 95.** Cujescu, D, Nenciu F, Persu C, Găgeanu I, Gabriel G, Vlăduț NV, Matache M, Voicea I, Pruteanu A, Bularda M, Paraschiv G: Evaluation of an optical sorter effectiveness in separating maize seeds intended for sowing. *Appl Sci*, 13:8892, 2023. DOI: 10.3390/app13158892
- 96.** Bailly S, Orlando B, Brustel J, Bailly JD, Levasseur-Garcia C: Rapid detection of aflatoxins in ground maize using near infrared spectroscopy. *Toxins*, 16:385, 2024. DOI: 10.3390/toxins16090385
- 97.** Conte G, Fontanelli M, Galli F, Cotrozzi L, Pagni L, Pellegrini E: Mycotoxins in feed and food and the role of ozone in their detoxification and degradation: An update. *Toxins*, 12:486, 2020. DOI: 10.3390/toxins12080486
- 98.** Shetty PH, Bhat RV: A physical method for segregation of fumonisins-contaminated maize. *Food Chem*, 66 (3): 371-374, 1999. DOI: 10.1016/S0308-8146(99)00052-7
- 99.** Van der Westhuizen L, Shephard GS, Rheeder J, Burger HM, Gelderblom W, Wild C, Gong Y: Optimising sorting and washing of home-grown maize to reduce fumonisin contamination under laboratory-controlled conditions. *Food Cont*, 22 (3-4): 396-400, 2011. DOI: 10.1016/j.foodcont.2010.09.009
- 100.** Bian Y, Zhang Y, Zhou Y, Wei B, Feng X: Recent insights into sample pretreatment methods for mycotoxins in different food matrices: A critical review on novel materials. *Toxins*, 15:215, 2023. DOI: 10.3390/toxins15030215
- 101.** Yumbe-Guevara BE, Imoto T, Yoshizawa T: Effects of heating procedures on deoxynivalenol, nivalenol and zearalenone levels in naturally contaminated barley and wheat. *Food Addit Contam*, 20 (12): 1132-1140, 2003. DOI: 10.1080/02652030310001620432
- 102.** Kabak B, Dobson AD, Var I: Strategies to prevent mycotoxin contamination of food and animal feed: A review. *Crit Rev Food Sci Nutr*, 46 (8): 593-619, 2006. DOI: 10.1080/10408390500436185
- 103.** Ryu D, Hanna MA, Eskridge KM, Bullerman LB: Heat stability of zearalenone in an aqueous buffered model system. *J Agric Food Chem*, 51 (6): 1746-1748, 2003. DOI: 10.1021/jf0210021

- 104.** Colovic R, Puvaca N, Cheli F, Avantaggiato G, Greco D, Duragic O, Kos J, Pinotti L: Decontamination of mycotoxin-contaminated feedstuffs and compound feed. *Toxins (Basel)*, 11 (11):617, 2019. DOI: 10.3390/toxins11110617
- 105.** Park JW, Kim YB: Effect of pressure cooking on aflatoxin B1 in rice. *J Agric Food Chem*, 54 (6): 2431-2435, 2006. DOI: 10.1021/jf053007e
- 106.** Becker-Algeri TA, Heidtmann-Bemvenuti R, dos Santos Hackbart HC, Badiale-Furlong E: Thermal treatments and their effects on the fumonisin B1 level in rice. *Food Cont*, 34 (2): 488-493, 2013. DOI: 10.1016/j.foodcont.2013.05.016
- 107.** Guo Y, Zhao L, Ma Q, Ji C: Novel strategies for degradation of aflatoxins in food and feed: A review. *Food Res Int*, 140:109878, 2021. DOI: 10.1016/j.foodres.2020.109878
- 108.** Zhang X, Chen Z, Gu W, Ji W, Wang Y, Hao C, He Y, Huang L, Wang M, Shao X, Yan Y: Viral and bacterial co-infection in hospitalised children with refractory *Mycoplasma pneumoniae* pneumonia. *Epidemiol Infect*, 146 (11): 1384-1388, 2018. DOI: 10.1017/S0950268818000778
- 109.** Calado T, Venâncio A, Abrunhosa L: Irradiation for mold and mycotoxin control: A review. *Comp Rev Food Sci Food Saf*, 13 (5): 1049-1061, 2014. DOI: 10.1111/1541-4337.12095
- 110.** Di Stefano V, Pitonzo R, Cicero N, D’Oca MC: Mycotoxin contamination of animal feedingstuff: detoxification by gamma-irradiation and reduction of aflatoxins and ochratoxin A concentrations. *Food Addit Contam*, 31 (12): 2034-2039, 2014. DOI: 10.1080/19440049.2014.968882
- 111.** Herzallah S, Alshawabkeh K, Fataftah AA: Aflatoxin decontamination of artificially contaminated feeds by sunlight, γ -radiation, and microwave heating. *J Appl Poult Res*, 17 (4): 515-521, 2008. DOI: 10.3382/japr.2007-00107
- 112.** Markov K, Mihaljević B, Domijan AM, Pleadin J, Delaš F, Frece J: Inactivation of aflatoxigenic fungi and the reduction of aflatoxin B1 *in vitro* and *in situ* using gamma irradiation. *Food Cont*, 54, 79-85, 2015. DOI: 10.1016/j.foodcont.2015.01.036
- 113.** Serra MS, Pulles MB, Mayanquer FT, Vallejo MC, Rosero MI, Ortega JM, Naranjo LN: Evaluation of the use of gamma radiation for reduction of aflatoxin B1 in corn (*Zea mays*) used in the production of feed for broiler chickens. *J Agric Chem Environ*, 7 (1): 21-33, 2018. DOI: 10.4236/jacen.2018.71003
- 114.** Abd El-Tawaab AA, El-Hofy FI, Mahmoud AH, Rashed DM: Mycotoxin residues in different chicken products by HPLC and their inactivation using Gamma radiation. *Int J Pharma Res All Sci*, 8 (4-2019): 71-81, 2019.
- 115.** Refai M, Aziz N, El-Far F, Hassan A: Detection of ochratoxin produced by *A. ochraceus* in feedstuffs and its control by γ radiation. *Appl Rad Iso*, 47 (7): 617-621, 1996. DOI: 10.1016/0969-8043(96)00022-X
- 116.** El-Far F, Aziz N, Hegazy S: Inhibition by gamma-irradiation and antimicrobial food additives of aflatoxin B1 production by *Aspergillus flavus* in poultry diet. *Food/Nahrung*, 36 (2): 143-149, 1992. DOI: 10.1002/food.19920360207
- 117.** Luo X, Zhai Y, Qi L, Pan L, Wang J, Xing J, Wang R, Wang L, Zhang Q, Yang K, Chen Z: Influences of electron beam irradiation on the physical and chemical properties of zearalenone- and ochratoxin A-contaminated corn and *in vivo* toxicity assessment. *Foods*, 9 (3):376, 2020. DOI: 10.3390/foods9030376
- 118.** Kottapalli B, Wolf-Hall CE, Schwarz P: Effect of electron-beam irradiation on the safety and quality of Fusarium-infected malting barley. *Int J Food Microbiol*, 110 (3): 224-31, 2006. DOI: 10.1016/j.ijfoodmicro.2006.04.007
- 119.** Stepanik T, Kost D, Nowicki T, Gaba D: Effects of electron beam irradiation on deoxynivalenol levels in distillers dried grain and solubles and in production intermediates. *Food Addit Contam*, 24 (9): 1001-1006, 2007. DOI: 10.1080/02652030701329629
- 120.** Wang B, Mahoney NE, Pan Z, Khir R, Wu B, Ma H, Zhao L: Effectiveness of pulsed light treatment for degradation and detoxification of aflatoxin B1 and B2 in rough rice and rice bran. *Food Cont*, 59, 461-467, 2016. DOI: 10.1016/j.foodcont.2015.06.030
- 121.** Ghangro AB, Channa MJ, Sheikh SA, Nizamani SM, Ghangro IH: Assessment of aflatoxin level in stored wheat of godowns of Hyderabad division and decontamination by uv radiation. *Int J Biosci*, 8 (1): 8-16, 2016. DOI: 10.12692/ijb/8.1.8-16
- 122.** Alkadi H, Altal J: Effect of microwave oven processing treatments on reduction of aflatoxin B1 and ochratoxin A in maize flour. *Europ J Chem*, 10 (3): 224-227, 2019. DOI: 10.5155/eurjchem.10.3.224-227.1840
- 123.** Mohamed NF, El-Dine RS, Kotb MAM, Saber A: Assessing the possible effect of gamma irradiation on the reduction of aflatoxin B1, and on the moisture content in some cereal grains. *Am J Biomed Sci*, 7 (1): 33-39, 2015. DOI: 10.5099/ajb150100033
- 124.** Wang SQ, Huang GQ, Li YP, Xiao JX, Zhang Y, Jiang WL: Degradation of aflatoxin B1 by low-temperature radio frequency plasma and degradation product elucidation. *Europ Food Res Technol*, 241 (1): 103-113, 2015. DOI: 10.1007/s00217-015-2439-5
- 125.** Liu R, Wang R, Lu J, Chang M, Jin Q, Du Z, Wang S, Li Q, Wang X: Degradation of AFB1 in aqueous medium by electron beam irradiation: Kinetics, pathway and toxicology. *Food Cont*, 66, 151-157, 2016. DOI: 10.1016/j.foodcont.2016.02.002
- 126.** Liu R, Lu M, Wang R, Wang S, Chang M, Jin Q, Wang X: Degradation of aflatoxin B1 in peanut meal by electron beam irradiation. *Int J Food Prop*, 21 (1): 892-901, 2018. DOI: 10.1080/10942912.2018.1466321
- 127.** Gayán E, Condón S, Álvarez I: Biological aspects in food preservation by ultraviolet light: A review. *Food Bioproc Technol*, 7 (1): 1-20, 2014. DOI: 10.1007/s11947-013-1168-7
- 128.** Fan X, Huang R, Chen H: Application of ultraviolet C technology for surface decontamination of fresh produce. *Trends Food Sci Technol*, 70, 9-19, 2017. DOI: 10.1016/j.tifs.2017.10.004
- 129.** Sun S, Zhao R, Xie Y, Liu Y: Photocatalytic degradation of aflatoxin B1 by activated carbon supported TiO₂ catalyst. *Food Cont*, 100, 183-188, 2019. DOI: 10.1016/j.foodcont.2019.01.014
- 130.** Xu C, Ye S, Cui X, Song X, Xie X: Modelling photocatalytic detoxification of aflatoxin B1 in peanut oil on TiO₂ layer in a closed-loop reactor. *Biosys Eng*, 180, 87-95, 2019. DOI: 10.1016/j.biosystemseng.2019.01.018
- 131.** Wu S, Wang F, Li Q, Zhou Y, He C, Duan N: Detoxification of DON by photocatalytic degradation and quality evaluation of wheat. *RSC Adv*, 9 (59): 34351-34358, 2019. DOI: 10.1039/c9ra04316k
- 132.** Oms-Oliu G, Martín-Belloso O, Soliva-Fortuny R: Pulsed light treatments for food preservation. A review. *Food Bioprocess Technol*, 3 (1): 13-23, 2010.
- 133.** Moreau M, Lescure G, Agoulon A, Svinareff P, Orange N, Feuilloley M: Application of the pulsed light technology to mycotoxin degradation and inactivation. *J Appl Toxicol*, 33 (5): 357-63, 2013. DOI: 10.1002/jat.1749
- 134.** Soni A, Smith J, Thompson A, Brightwell G: Microwave-induced thermal sterilization-A review on history, technical progress, advantages and challenges as compared to the conventional methods. *Tr Food Sci Technol*, 97, 433-442, 2020. DOI: 10.1016/j.tifs.2020.01.030
- 135.** Perez-Flores GC, Moreno-Martinez E, Mendez-Albores A: Effect of microwave heating during alkaline-cooking of aflatoxin contaminated maize. *J Food Sci*, 76 (2): T48-T52, 2011. DOI: 10.1111/j.1750-3841.2010.01980.x
- 136.** Menon A, Stojceska V, Tassou SA: A systematic review on the recent advances of the energy efficiency improvements in non-conventional food drying technologies. *Tr Food Sci Technol*, 100, 67-76, 2020. DOI: 10.1016/j.tifs.2020.03.014
- 137.** Kogelschatz U: Atmospheric-pressure plasma technology. *Plasma Physics and Controlled Fusion*, 46 (12B):B63, 2004. DOI: 10.1088/0741-3335/46/12B/006
- 138.** Fridman A, Chirokov A, Gutsol A: Non-thermal atmospheric pressure discharges. *J Phys*, 38 (2):R1, 2005. DOI: 10.1088/0022-3727/38/2/R01
- 139.** Misra NN, Yadav B, Roopesh MS, Jo C: Cold plasma for effective fungal and mycotoxin control in foods: Mechanisms, inactivation effects, and applications. *Compr Rev Food Sci Food Saf*, 18 (1): 106-120, 2019. DOI: 10.1111/1541-4337.12398
- 140.** Siciliano I, Spadaro D, Prella A, Vallauri D, Cavallero MC, Garibaldi A, Gullino ML: Use of cold atmospheric plasma to detoxify hazelnuts from

- aflatoxins. *Toxins (Basel)*, 8 (5):125, 2016. DOI: 10.3390/toxins8050125
- 141. Ouf SA, Mohamed AAH, El-Sayed WS:** Fungal decontamination of fleshy fruit water washes by double atmospheric pressure cold plasma. *CLEAN-Soil Air Water*, 44 (2): 134-142, 2016. DOI: 10.1002/clen.201400575
- 142. Park BJ, Takatori K, Sugita-Konishi Y, Kim IH, Lee MH, Han DW, Chung KH, Hyun SO, Park JC:** Degradation of mycotoxins using microwave-induced argon plasma at atmospheric pressure. *Surf Coat Technol*, 201 (9-11): 5733-5737, 2007. DOI: 10.1016/j.surfcoat.2006.07.092
- 143. Kriz P, Petr B, Zbynek H, Jaromir K, Pavel O, Petr S, Miroslav D:** Influence of plasma treatment in open air on mycotoxin content and grain nutriments. *Plasma Med*, 5 (2-4): 145-148, 2015. DOI: 10.1615/PlasmaMed.2016015752
- 144. Iqdam BM, Feizollahi E, Arif MF, Jeganathan B, Vasanthan T, Thilakarathna MS, Roopesh MS:** Reduction of T-2 and HT-2 mycotoxins by atmospheric cold plasma and its impact on quality changes and germination of wheat grains. *J Food Sci*, 86 (4): 1354-1371, 2021. DOI: 10.1111/1750-3841.15658
- 145. Ten Bosch L, Pfohl K, Avramidis G, Wieneke S, Viol W, Karlovsky P:** Plasma-based degradation of mycotoxins produced by Fusarium, Aspergillus and Alternaria species. *Toxins (Basel)*, 9 (3):97, 2017. DOI: 10.3390/toxins9030097
- 146. Adamović M, Stojanović M, Grubišić M, Ileš D, Milojković J:** Importance of aluminosilicate minerals in safe food production. *Mac J Anim Sci*, 1, 175-180, 2011.
- 147. Magnoli AP, Texeira M, Rosa CA, Mazzola RD, Cavagliari LR, Magnoli CE, Dalcerio AM, Chiacchiera SM:** Sodium bentonite and monensin under chronic aflatoxicosis in broiler chickens. *Poult Sci*, 90 (2): 352-357, 2011. DOI: 10.3382/ps.2010-00834
- 148. Chen Y, Kong Q, Chi C, Shan S, Guan B:** Biotransformation of aflatoxin B1 and aflatoxin G1 in peanut meal by anaerobic solid fermentation of *Streptococcus thermophilus* and *Lactobacillus delbrueckii* subsp. *bulgaricus*. *Int J Food Microbiol*, 211: 1-5, 2015. DOI: 10.1016/j.ijfoodmicro.2015.06.021
- 149. Bhatti SA, Khan MZ, Saleemi MK, Saqib M, Khan A, Hassan Z:** Protective role of bentonite against aflatoxin B1-and ochratoxin A-induced immunotoxicity in broilers. *J Immunotox*, 14 (1): 66-76, 2017. DOI: 10.1080/1547691X.2016.1264503
- 150. Sumantri I, Murti T, Van der Poel A, Boehm J, Agus A:** Carry-over of aflatoxin B1-feed into aflatoxin M1-milk in dairy cows treated with natural sources of aflatoxin and bentonite. *J Indo Trop Anim Agric*, 37 (4): 271-277, 2012. DOI: 10.14710/jitaa.37.4.271-277
- 151. Gouda G, Khattab H, Abdel-Wahhab M, El-Nor SA, El-Sayed H, Khalif S:** Clay minerals as sorbents for mycotoxins in lactating goat's diets: Intake, digestibility, blood chemistry, ruminal fermentation, milk yield and composition, and milk aflatoxin M1 content. *Small Rum Res*, 175, 15-22, 2019. DOI: 10.1016/j.smallrumres.2019.04.003
- 152. Tzou YM, Chan YT, Chen SE, Wang CC, Chiang PN, Teah HY, Hung JT, Wu JJ, Liu YT:** Use 3-D tomography to reveal structural modification of bentonite-enriched clay by nonionic surfactants: Application of organo-clay composites to detoxify aflatoxin B1 in chickens. *J Hazard Mat*, 375, 312-319, 2019. DOI: 10.1016/j.jhazmat.2019.04.084
- 153. Kubena LF, Harvey RB, Huff WE, Elissalde MH, Yersin A G, Phillips TD, Rottinghaus GE:** Efficacy of a hydrated sodium calcium aluminosilicate to reduce the toxicity of aflatoxin and diacetoxyscirpenol. *Poult Sci*, 72 (1): 51-59, 1993. DOI: 10.3382/ps.0720051
- 154. Mussadeq Y, Begum I, Akhter S:** Activity of aflatoxin adsorbents in poultry feed. *Pak J Biol Sci*, 10, 1697-1699, 2000.
- 155. Robinson A, Johnson NM, Strey A, Taylor JE, Marroquin-Cardona A, Mitchell NJ, Afriyie-Gyawu E, Ankrah NA, Williams JH, Wang JS, Jolly PE, Nachman RJ, Phillips TD:** Calcium montmorillonite clay reduces urinary biomarkers of fumonisins B(1) exposure in rats and humans. *Food Addit Contam*, 29 (5): 809-818, 2012. DOI: 10.1080/19440049.2011.651628
- 156. Li Y, Tian G, Dong G, Bai S, Han X, Liang J, Meng J, Zhang H:** Research progress on the raw and modified montmorillonites as adsorbents for mycotoxins: A review. *Appl Clay Sci*, 163, 299-311, 2018. DOI: 10.1016/j.clay.2018.07.032
- 157. Phillips TD, Wang M, Elmore SE, Hearon S, Wang JS:** NovaSil clay for the protection of humans and animals from aflatoxins and other contaminants. *Clays Clay Miner*, 67 (1): 99-110, 2019. DOI: 10.1007/s42860-019-0008-x
- 158. Masimango N, Remacle J, Ramaut J:** The role of adsorption in the elimination of aflatoxin B1 from contaminated media. *Europ J Appl Micro Biotechnol*, 6, 101-105, 1978.
- 159. Decker WJ, Corby DG:** Activated charcoal adsorbs aflatoxin B1. *Vet Hum Toxicol*, 22 (6): 388-389, 1980.
- 160. Hatch RC, Clark JD, Jain AV, Weiss R:** Induced acute aflatoxicosis in goats: treatment with activated charcoal or dual combinations of oxytetracycline, stanozolol, and activated charcoal. *Am J Vet Res*, 43 (4): 644-648, 1982. DOI: 10.2460/ajvr.1982.43.04.644
- 161. Ademoyero AA, Dalvi RR:** Efficacy of activated charcoal and other agents in the reduction of hepatotoxic effects of a single dose of aflatoxin B1 in chickens. *Toxicol Lett*, 16 (1-2): 153-157, 1983. DOI: 10.1016/0378-4274(83)90024-3
- 162. Dalvi R, Ademoyero A:** Toxic effects of aflatoxin B1 in chickens given feed contaminated with Aspergillus flavus and reduction of the toxicity by activated charcoal and some chemical agents. *Avian Dis*, 61-69, 1984. DOI: 10.2307/1590128
- 163. Davidson J, Babish J, Delaney K, Taylor D, Phillips T:** Hydrated sodium calcium aluminosilicate decreases the bioavailability of aflatoxin in the chicken. *Poult Sci*, 6 (1): (Abstract) 1987.
- 164. Phillips TD, Kubena LF, Harvey RB, Taylor DR, Heidelbaugh ND:** Hydrated sodium calcium aluminosilicate: A high affinity sorbent for aflatoxin. *Poult Sci*, 67 (2): 243-247, 1988. DOI: 10.3382/ps.0670243
- 165. Kubena L, Harvey R, Phillips T, Huff W:** Modulation of aflatoxicosis in growing chickens by dietary addition of a hydrated sodium calcium aluminosilicate. *Poult Sci*, 67:106, 1988.
- 166. Doerr J:** Effect of aluminosilicate on broiler chickens during aflatoxicosis. *Poult Sci*, 68 (1):45, 1989.
- 167. Harvey R, Kubena L, Phillips T, WEH, Corrier D:** Prevention of clinical signs of aflatoxicosis with hydrated sodium calcium aluminosilicate added to diets. *Proc 20th Annu Meet Am Assoc Swine Pract*, 99-102, 1989.
- 168. Gelven RE:** Approaches for alleviating toxic effects of aflatoxin in lactating dairy cows and weanling pigs. *MSc Thesis*, University of Missouri-Columbia, 2010.
- 169. Dvorak M:** Ability of bentonite and natural zeolite to adsorb aflatoxin from liquid media. *Vet Med*, 34 (5): 307-316, 1989.
- 170. Lindemann M, Blodgett D, Schurig G, Kornegay E:** Evaluation of potential ameliorators of aflatoxicosis in weanling/growing swine. *J Anim Sci*, 67:36, 1989.
- 171. Fukal L, Slamova A, Novak L, Sova Z:** The effect of a high aflatoxin B1 concentration in feed on the weights of organs and strength of bones in grown-up chickens. *Biolog Chem Zivoc Vyrobny-Veterinaria (Czechoslovakia)*, 26 (2), 1990.
- 172. Beaver RW, Wilson DM, James MA, Haydon KD, Colvin BM, Sangster LT, Piluk AH, Groopman JD:** Distribution of aflatoxins in tissues of growing pigs fed an aflatoxin-contaminated diet amended with a high affinity aluminosilicate sorbent. *Vet Hum Toxicol*, 32 (1): 16-18, 1990.
- 173. Phillips TD, Clement BA, Kubena LF, Harvey RB:** Detection and detoxification of aflatoxins: prevention of aflatoxicosis and aflatoxin residues with hydrated sodium calcium aluminosilicate. *Vet Hum Toxicol*, 32, 15-19, 1990.
- 174. Kubena LF, Harvey RB, Huff WE, Corrier DE, Phillips TD, Rottinghaus GE:** Efficacy of a hydrated sodium calcium aluminosilicate to reduce the toxicity of aflatoxin and T-2 toxin. *Poult Sci*, 69 (7): 1078-1086, 1990. DOI: 10.3382/ps.0691078
- 175. Abdelhamid AM, el-Shawaf, el-Ayoty SA, Ali MM, Gamil T:** Effect of low level of dietary aflatoxins on baladi rabbits. *Arch Tierernahr*, 40 (5-6): 517-537, 1990. DOI: 10.1080/17450399009421084
- 176. Sova Z, Pohunková H, Reisnerová H, Slámová A, Haisl K:** Hematological and histological response to the diet containing aflatoxin B 1 and zeolite in broilers of domestic fowl. *Acta Vet Brno*, 60 (1): 31-40, 1991.
- 177. Araba M, Wyatt R:** Effects of sodium bentonite, hydrated sodium

- calcium aluminosilicate NovaSil™, and ethacal on aflatoxicosis in broiler chickens. *Poult Sci*, 70 (6), 1991.
- 178. Kubena LF, Huff WE, Harvey RB, Yersin AG, Elissalde MH, Witzel DA, Giroir LE, Phillips TD, Petersen HD:** Effects of a hydrated sodium calcium aluminosilicate on growing turkey pouls during aflatoxicosis. *Poult Sci*, 70 (8): 1823-1830, 1991. DOI: 10.3382/ps.0701823
- 179. Bonna RJ, Aulerich RJ, Bursian SJ, Poppenga RH, Braselton WE, Watson GL:** Efficacy of hydrated sodium calcium aluminosilicate and activated charcoal in reducing the toxicity of dietary aflatoxin to mink. *Arch Environ Contam Toxicol*, 20 (3): 441-447, 1991. DOI: 10.1007/BF01064418
- 180. Harvey RB, Kubena LF, Phillips TD, Corrier DE, Elissalde MH, Huff WE:** Diminution of aflatoxin toxicity to growing lambs by dietary supplementation with hydrated sodium calcium aluminosilicate. *Am J Vet Res*, 52 (1): 152-156, 1991. DOI: 10.2460/ajvr.1991.52.01.152
- 181. Scheideler SE:** Effects of various types of aluminosilicates and aflatoxin B1 on aflatoxin toxicity, chick performance, and mineral status. *Poult Sci*, 72 (2): 282-288, 1993. DOI: 10.3382/ps.0720282
- 182. Schell T, Lindemann M, Kornegay E, Blodgett D:** Effects of feeding aflatoxin-contaminated diets with and without clay to weanling and growing pigs on performance, liver function, and mineral metabolism. *J Anim Sci*, 71 (5): 1209-1218, 1993. DOI: 10.2527/1993.71151209x
- 183. Harvey RB, Kubena LF, Elissalde MH, Phillips TD:** Efficacy of zeolitic ore compounds on the toxicity of aflatoxin to growing broiler chickens. *Avian Dis*, 37 (1): 67-73, 1993. DOI: 10.2307/1591459
- 184. Lindemann MD, Blodgett DJ, Kornegay ET, Schurig GG:** Potential ameliorators of aflatoxicosis in weanling/growing swine. *J Anim Sci*, 71 (1): 171-178, 1993. DOI: 10.2527/1993.711171x
- 185. Ramos AJ, Fink-Gremmels J, Hernandez E:** Prevention of Toxic Effects of Mycotoxins by Means of Nonnutritive Adsorbent Compounds. *J Food Prot*, 59 (6): 631-641, 1996. DOI: 10.4315/0362-028X-59.6.631
- 186. Edrington TS, Sarr AB, Kubena LF, Harvey RB, Phillips TD:** Hydrated sodium calcium aluminosilicate (HSCAS), acidic HSCAS, and activated charcoal reduce urinary excretion of aflatoxin M1 in turkey pouls. Lack of effect by activated charcoal on aflatoxicosis. *Toxicol Lett*, 89 (2): 115-122, 1996. DOI: 10.1016/s0378-4274(96)03795-2
- 187. Edrington TS, Kubena LF, Harvey RB, Rottinghaus GE:** Influence of a superactivated charcoal on the toxic effects of aflatoxin or T-2 toxin in growing broilers. *Poult Sci*, 76 (9): 1205-1211, 1997. DOI: 10.1093/ps/76.9.1205
- 188. Bailey R, Kubena L, Harvey R, Buckley S, Rottinghaus G:** Efficacy of various inorganic sorbents to reduce the toxicity of aflatoxin and T-2 toxin in broiler chickens. *Poult Sci*, 77 (11): 1623-1630, 1998. DOI: 10.1093/ps/77.11.1623
- 189. Natour R, Yousef S:** Adsorption efficiency of diatomaceous earth for mycotoxin. *Arab Gulf J Sci Res*, 16, 113-127, 1998.
- 190. Parlat SS, Yildiz AO, Oguz H:** Effect of clinoptilolite on performance of Japanese quail (*Coturnix coturnix japonica*) during experimental aflatoxicosis. *Br Poult Sci*, 40 (4): 495-500, 1999. DOI: 10.1080/00071669987269
- 191. Flores CM, Dominguez JM, Diaz-De-Leon J:** Modeling and experimental comparison of the differential adsorption of B1 and G1 aflatoxins on mineral aluminosilicate surfaces. *J Environ Pathol Toxicol Oncol*, 18 (3): 213-220, 1999.
- 192. Abdel-Wahhab MA, Nada SA, Amra HA:** Effect of aluminosilicates and bentonite on aflatoxin-induced developmental toxicity in rat. *J Appl Toxicol*, 19 (3): 199-204, 1999. DOI: 10.1002/(sici)1099-1263(199905/06)19:3<199::aid-jat558>3.0.co;2-d
- 193. Ledoux DR, Rottinghaus GE, Bermudez AJ, Alonso-Debolt M:** Efficacy of a hydrated sodium calcium aluminosilicate to ameliorate the toxic effects of aflatoxin in broiler chicks. *Poult Sci*, 78 (2): 204-210, 1999. DOI: 10.1093/ps/78.2.204
- 194. Miazzo R, Rosa CA, De Queiroz Carvalho EC, Magnoli C, Chiacchiera SM, Palacio G, Saenz M, Kikot A, Basaldella E, Dalcero A:** Efficacy of synthetic zeolite to reduce the toxicity of aflatoxin in broiler chicks. *Poult Sci*, 79 (1): 1-6, 2000. DOI: 10.1093/ps/79.1.1
- 195. Soufiani GN, Razmara M, Kermanshahi H, Velazquez AB, Daneshmand A:** Assessment of aflatoxin B1 adsorption efficacy of natural and processed bentonites: In vitro and in vivo assays. *Appl Clay Sci*, 123: 129-133, 2016. DOI: 10.1016/j.clay.2016.01.019
- 196. Rosa CA, Miazzo R, Magnoli C, Salvano M, Chiacchiera SM, Ferrero S, Saenz M, Carvalho EC, Dalcero A:** Evaluation of the efficacy of bentonite from the south of Argentina to ameliorate the toxic effects of aflatoxin in broilers. *Poult Sci*, 80 (2): 139-144, 2001. DOI: 10.1093/ps/80.2.139
- 197. Aly SE, Abdel-Galil MM, Abdel-Wahhab MA:** Application of adsorbent agents technology in the removal of aflatoxin B(1) and fumonisin B(1) from malt extract. *Food Chem Toxicol*, 42 (11): 1825-1831, 2004. DOI: 10.1016/j.fct.2004.06.014
- 198. Sumantri I, Herliani H, Yuliani M, Nuryono N:** Effects of zeolite in aflatoxin B1 contaminated diet on aflatoxin residues and liver histopathology of laying duck. In, *IOP Conf Ser: Earth Environ Sci*, 2018
- 199. Rotter RG, Frohlich AA, Marquardt RR:** Influence of dietary charcoal on ochratoxin A toxicity in Leghorn chicks. *Can J Vet Res*, 53 (4): 449-453, 1989
- 200. Huff WE, Kubena LF, Harvey RB, Phillips TD:** Efficacy of hydrated sodium calcium aluminosilicate to reduce the individual and combined toxicity of aflatoxin and ochratoxin A. *Poult Sci*, 71 (1): 64-69, 1992. DOI: 10.3382/ps.0710064
- 201. Bauer J:** Methods for detoxification of mycotoxins in feedstuffs. *Monatshefte fuer Veterinaermedizin (Germany)*, 49 (4): 175-181, 1994.
- 202. Galvano F, Pietri A, Bertuzzi T, Piva A, Chies L, Galvano M:** Activated carbons: *In vitro* affinity for ochratoxin A and deoxynivalenol and relation of adsorption ability to physicochemical parameters. *J Food Prot*, 61 (4): 469-475, 1998. DOI: 10.4315/0362-028x-61.4.469
- 203. Khatoon A, Khan M Z, Abidin Z, Bhatti SA:** Effects of feeding bentonite clay upon ochratoxin A-induced immunosuppression in broiler chicks. *Food Add Contam*, 35 (3): 538-545, 2018. DOI: 10.1080/19440049.2017.1411612
- 204. Galvano F, Pietri A, Bertuzzi T, Bognanno M, Chies L, Galvano M:** Activated carbons: *In vitro* affinity for fumonisin B(1) and relation of adsorption ability to physicochemical parameters. *J Food Prot*, 60 (8): 985-991, 1997. DOI: 10.4315/0362-028X-60.8.985
- 205. Solfrizzo M, Carratu MR, Avantaggiato G, Galvano F, Pietri A, Visconti A:** Ineffectiveness of activated carbon in reducing the alteration of sphingolipid metabolism in rats exposed to fumonisin-contaminated diets. *Food Chem Toxicol*, 39 (5): 507-511, 2001. DOI: 10.1016/s0278-6915(00)00160-5
- 206. Vila-Donat P, Marín S, Sanchis V, Ramos AJ:** Tri-octahedral bentonites as potential technological feed additive for Fusarium mycotoxin reduction. *Food Add Contam*, 37 (8): 1374-1387, 2020. DOI: 10.1080/19440049.2020.1766702
- 207. Smith TK:** Dietary influences on excretory pathways and tissue residues of zearalenone and zearalenols in the rat. *Can J Physiol Pharmacol*, 60 (12): 1444-1449, 1982. DOI: 10.1139/y82-214
- 208. Bursian SJ, Aulerich RJ, Cameron JK, Ames NK, Steficek BA:** Efficacy of hydrated sodium calcium aluminosilicate in reducing the toxicity of dietary zearalenone to mink. *J Appl Toxicol*, 12 (2): 85-90, 1992. DOI: 10.1002/jat.2550120204
- 209. Ramos A, Hernandez E:** Resin colestiramina: Un adsorbente de zearalenona de alta afinidad. In, *Procc. XIV Congreso Nacional de Microbiologia*. Zaragoza, Spain. 17-19 September, 1993.
- 210. Williams K, Blaney B, Peters R:** Pigs fed Fusarium-infected maize containing zearalenone and nivalenol with sweeteners and bentonite. *Livest Prod Sci*, 39 (3): 275-281, 1994. DOI: 10.1016/0301-6226(94)90207-0
- 211. Chen Q, Lu Z, Hou W, Shi B, Shan A:** Effects of modified maifanite on zearalenone toxicity in female weaner pigs. *It J Anim Sci*, 14 (2): 3597, 2015. DOI: 10.4081/ijas.2015.3597
- 212. Avantaggiato G, Havenga R, Visconti A:** Assessing the zearalenone-binding activity of adsorbent materials during passage through a dynamic *in vitro* gastrointestinal model. *Food Chem Toxicol*, 41 (10): 1283-1290, 2003. DOI: 10.1016/S0278-6915(03)00113-3

- 213.** Lemke SL, Grant PG, Phillips TD: Adsorption of zearalenone by organophilic montmorillonite clay. *J Agric Food Chem*, 46 (9): 3789-3796, 1998. DOI: 10.1021/jf9709461
- 214.** Friend D, Trenholm H, Martin K, Young J, Thompson B: Effect of adding potential vomitoxin (deoxynivalenol) detoxicants or aF graminearum inoculated corn supplement to wheat diets fed to pigs. *Can J Anim Sci*, 64 (3): 733-741, 1984. DOI: 10.4141/cjas84-081
- 215.** Avantaggiato G, Havenaar R, Visconti A: Evaluation of the intestinal absorption of deoxynivalenol and nivalenol by an *in vitro* gastrointestinal model, and the binding efficacy of activated carbon and other adsorbent materials. *Food Chem Toxicol*, 42 (5): 817-824, 2004. DOI: 10.1016/j.fct.2004.01.004
- 216.** Young JC, Subryan LM, Potts D, McLaren ME, Cobran FH: Reduction in levels of deoxynivalenol in contaminated wheat by chemical and physical treatment. *J Agric Food Chem*, 34 (3): 461-465, 1986. DOI: 10.1021/jf00069a021
- 217.** Kubena LF, Harvey RB, Bailey RH, Buckley SA, Rottinghaus GE: Effects of a hydrated sodium calcium aluminosilicate (T-Bind) on mycotoxicosis in young broiler chickens. *Poult Sci*, 77 (10): 1502-1509, 1998. DOI: 10.1093/ps/77.10.1502
- 218.** Carson MS, Smith TK: Role of bentonite in prevention of T-2 toxicosis in rats. *J Anim Sci*, 57 (6): 1498-506, 1983. DOI: 10.2527/jas1983.5761498x
- 219.** Buck W, Bratich P: Activated charcoal: Preventing unnecessary death by poisoning. *Vet Med*, 81, 73-77, 1986.
- 220.** Galey FD, Lambert RJ, Busse M, Buck WB: Therapeutic efficacy of superactive charcoal in rats exposed to oral lethal doses of T-2 toxin. *Toxicol*, 25 (5): 493-499, 1987. DOI: 10.1016/0041-0101(87)90285-6
- 221.** Poppenga RH, Lundein GR, Beasley VR, Buck WB: Assessment of a general therapeutic protocol for the treatment of acute T-2 toxicosis in swine. *Vet Hum Toxicol*, 29 (3): 237-239, 1987.
- 222.** Dwyer MR, Kubena LF, Harvey RB, Mayura K, Sarr AB, Buckley S, Bailey RH, Phillips TD: Effects of inorganic adsorbents and cyclopiazonic acid in broiler chickens. *Poult Sci*, 76 (8): 1141-1149, 1997. DOI: 10.1093/ps/76.8.1141
- 223.** Huebner HJ, Lemke SL, Ottinger SE, Mayura K, Phillips TD: Molecular characterization of high affinity, high capacity clays for the equilibrium sorption of ergotamine. *Food Addit Contam*, 16 (4): 159-171, 1999. DOI: 10.1080/026520399284118
- 224.** Sands DC, McIntyre JL, Walton GS: Use of activated charcoal for the removal of patulin from cider. *Appl Environ Microbiol*, 32 (3): 388-391, 1976. DOI: 10.1128/aem.32.3.388-391.1976
- 225.** Dvorak Z: Mechanical engineering problems in preserving biological objects by temperature lowering. *Sb Ved Pr Lek Fak Karlovy Univ Hradci Kralove*, 33 (2): 115-125, 1990.
- 226.** Jalili M, Jinap S, Son R: The effect of chemical treatment on reduction of aflatoxins and ochratoxin A in black and white pepper during washing. *Food Addit Contam*, 28 (4): 485-493, 2011. DOI: 10.1080/19440049.2010.551300
- 227.** Javanmardi F, Khodaei D, Sheidaei Z, Bashiry M, Nayebzadeh K, Vasseghian Y, Mousavi Khanegah A: Decontamination of aflatoxins in edible oils: A comprehensive review. *Food Rev Int*, 38 (7): 1410-1426, 2022. DOI: 10.1080/87551920.2020.1812635
- 228.** Park DL: Perspectives on mycotoxin decontamination procedures. *Food Addit Contam*, 10 (1): 49-60, 1993. DOI: 10.1080/02652039309374129
- 229.** Natarajan KR: Chemical inactivation of aflatoxins in peanut protein ingredients. *J Environ Pathol Toxicol Oncol*, 11 (4): 217-227, 1992.
- 230.** McKenzie KS, Sarr AB, Mayura K, Bailey RH, Miller DR, Rogers TD, Norred WP, Voss KA, Plattner RD, Kubena LF, Phillips TD: Oxidative degradation and detoxification of mycotoxins using a novel source of ozone. *Food Chem Toxicol*, 35 (8): 807-820, 1997. DOI: 10.1016/s0278-6915(97)00052-5
- 231.** de Alencar ER, Faroni LRDA, Soares N FF, da Silva WA, da Silva Carvalho MC: Efficacy of ozone as a fungicidal and detoxifying agent of aflatoxins in peanuts. *J Sci Food Agric*, 92 (4): 899-905, 2012. DOI: 10.1002/jsfa.4668
- 232.** Agriopoulou S, Koliadima A, Karauskakis G, Kapolos J: Kinetic study of aflatoxins' degradation in the presence of ozone. *Food Cont*, 61, 221-226, 2016. DOI: 10.1016/j.foodcont.2015.09.013
- 233.** Piemontese L, Messia MC, Marconi E, Falasca L, Zivoli R, Gambacorta L, Perrone G, Solfrizzo M: Effect of gaseous ozone treatments on DON, microbial contaminants and technological parameters of wheat and semolina. *Food Addit Contam*, 35 (4): 760-771, 2018. DOI: 10.1080/19440049.2017.1419285
- 234.** Santos Alexandre AP, Vela-Paredes RS, Santos AS, Costa NS, Canniatti-Brazaca SG, Calori-Domingues MA, Augusto PED: Ozone treatment to reduce deoxynivalenol (DON) and zearalenone (ZEN) contamination in wheat bran and its impact on nutritional quality. *Food Addit Contam*, 35 (6): 1189-1199, 2018. DOI: 10.1080/19440049.2018.1432899
- 235.** Jr AP, King J: Efficacy and safety evaluation of ozonation to degrade aflatoxin in corn. *J Food Sci*, 67 (8): 2866-2872, 2002. DOI: 10.1111/j.1365-2621.2002.tb08830.x
- 236.** Trombete F, Porto Y, Freitas-Silva O, Pereira R, Direito G, Saldanha T, Fraga M: Efficacy of ozone treatment on mycotoxins and fungal reduction in artificially contaminated soft wheat grains. *J Food Process Preserv*, 41 (3):e12927, 2017. DOI: 10.1111/jfpp.12927
- 237.** McKenzie K, Kubena L, Denvir A, Rogers T, Hitchens G, Bailey R, Harvey R, Buckley S, Phillips T: Aflatoxicosis in turkey poult is prevented by treatment of naturally contaminated corn with ozone generated by electrolysis. *Poult Sci*, 77 (8): 1094-1102, 1998. DOI: 10.1093/ps/77.8.1094
- 238.** Li MM, Guan EQ, Bian K: Effect of ozone treatment on deoxynivalenol and quality evaluation of ozonised wheat. *Food Addit Contam*, 32 (4): 544-553, 2015. DOI: 10.1080/19440049.2014.976596
- 239.** Sun C, Ji J, Wu S, Sun C, Pi F, Zhang Y, Tang L, Sun X: Saturated aqueous ozone degradation of deoxynivalenol and its application in contaminated grains. *Food Cont*, 69, 185-190, 2016. DOI: 10.1016/j.foodcont.2016.04.041
- 240.** Wang L, Luo Y, Luo X, Wang R, Li Y, Li Y, Shao H, Chen Z: Effect of deoxynivalenol detoxification by ozone treatment in wheat grains. *Food Cont*, 66, 137-144, 2016. DOI: 10.1016/j.foodcont.2016.01.038
- 241.** Qi L, Li Y, Luo X, Wang R, Zheng R, Wang L, Li Y, Yang D, Fang W, Chen Z: Detoxification of zearalenone and ochratoxin A by ozone and quality evaluation of ozonised corn. *Food Addit Contam*, 33 (11): 1700-1710, 2016. DOI: 10.1080/19440049.2016.1232863
- 242.** Altug T, Yousef AE, Marth EH: Degradation of aflatoxin B1 in dried figs by sodium bisulfite with or without heat, ultraviolet energy or hydrogen peroxide. *J Food Prot*, 53 (7): 581-583, 1990. DOI: 10.4315/0362-028X-53.7.581
- 243.** Abd Alla ES: Zearalenone: Incidence, toxigenic fungi and chemical decontamination in Egyptian cereals. *Nahrung*, 41 (6): 362-365, 1997. DOI: 10.1002/food.19970410610
- 244.** Chlebic A, Slizewska K: *In vitro* detoxification of aflatoxin B(1), deoxynivalenol, fumonisins, T-2 toxin and zearalenone by probiotic bacteria from genus *Lactobacillus* and *Saccharomyces cerevisiae* yeast. *Prob Antimicrob Proteins*, 12 (1): 289-301, 2020. DOI: 10.1007/s12602-018-9512-x
- 245.** Bidura IGNG, Siti NW, Wibawa AAP, Puspani E, Candrawati DPMA: Improving the quality of tofu waste by mixing it with carrots and probiotics as a feed source of probiotics and β -carotene. *Int J Vet Sci*, 12, 407-413, 2023. DOI: 10.47278/jurnal.ijvs/2022.213
- 246.** Akhtar T, Shahid S, Asghar A, Naeem MI, Aziz S, Ameer T: Utilisation of herbal bullets against Newcastle disease in poultry sector of Asia and Africa (2012-2022). *Int J Agri Biosci*, 12, 56-65, 2023. DOI: 10.47278/journal.ijab/2023.044
- 247.** Coniglio MV, Luna MJ, Provencal P, Magnoli AP: Use of the probiotic *Saccharomyces cerevisiae* var. boulardii RC009 in the rearing stage of calves. *Int J Agri Biosci*, 12, 188-192, 2023. DOI: 10.47278/journal.ijab/2023.063
- 248.** Kalita R, Pegu A, Baruah C: Prospects of probiotics and fish growth promoting bacteria in aquaculture: a review. *Int J Agri Biosci*, 12 (4): 234-244, 2023. DOI: 10.47278/journal.ijab/2023.070
- 249.** Rashid S, Alsayeqh AF, Akhtar T, Abbas R Z, Ashraf R: Probiotics: alternative to antibiotics in poultry production. *Int J Vet Sci*, 12, 45-53, 2023. DOI: 10.47278/journal.ijvs/2022.175

- 250.** Shah NP: Probiotic bacteria: Selective enumeration and survival in dairy foods. *J Dairy Sci.* 83 (4): 894-907, 2000. DOI: 10.3168/jds.S0022-0302(00)74953-8
- 251.** Mehmood A, Nawaz M, Rabbani M, Mushtaq MH: Probiotic effect of Limosilactobacillus fermentum on growth performance and competitive exclusion of *Salmonella gallinarum* in poultry. *Pak Vet J.* 43 (4): 659-664, 2023. DOI: 10.29261/pakvetj/2023.103
- 252.** Madrigal-Santillan E, Madrigal-Bujaidar E, Marquez-Marquez R, Reyes A: Antigenotoxic effect of *Saccharomyces cerevisiae* on the damage produced in mice fed with aflatoxin B(1) contaminated corn. *Food Chem Toxicol.* 44 (12): 2058-2063, 2006. DOI: 10.1016/j.fct.2006.07.006
- 253.** Teniola OD, Addo PA, Brost IM, Farber P, Jany KD, Alberts JF, van Zyl WH, Steyn PS, Holzapfel WH: Degradation of aflatoxin B(1) by cell-free extracts of *Rhodococcus erythropolis* and *Mycobacterium fluoranthenivorans* sp. nov. DSM44556(T). *Int J Food Microbiol.* 105 (2): 111-117, 2005. DOI: 10.1016/j.ijfoodmicro.2005.05.004
- 254.** Gul ST, Alsayeqh AF: Probiotics improve physiological parameters and meat production in broiler chicks. *Int J Vet Sci.* 12, 182-191, 2023. DOI: 10.47278/journal.ijvs/2022.191
- 255.** Raza A, Abbas RZ, Karadağlı Ö, Raheem A, Khan AMA, Khalil MZ, Maheen N, Quddus A, Hussain A, Kanchev KP: Role of probiotics in increasing meat and egg production in poultry. *Kafkas Univ Vet Fak Derg.* 30, 753-760, 2024. DOI: 10.9775/kvfd.2024.32861
- 256.** Dabool AS, Atwah B, Alghamdi S, Momenah MA, Saleh O, Alhazmi N, Mostafa YS, Alamri SA, Alyoubi WA, Alshammari NM: Could *Paenibacillus xylanexedens* MS58 be an ecofriendly antibiotic in poultry production? Impacts on performance, blood biochemistry, gut microbiota and meat quality. *Pak Vet J.* 44, 352-360, 2024. DOI: 10.29261/pakvetj/2024.180
- 257.** Farzaneh M, Shi ZQ, Ghassemour A, Sedaghat N, Ahmadzadeh M, Mirabolfathy M, Javan-Nikkhah M: Aflatoxin B1 degradation by *Bacillus subtilis* UTBSP1 isolated from pistachio nuts of Iran. *Food Cont.* 23 (1): 100-106, 2012. DOI: 10.1016/j.foodcont.2011.06.018
- 258.** Samuel MS, Sivararamakrishna A, Mehta A: Degradation and detoxification of aflatoxin B1 by *Pseudomonas putida*. *Int Biodeterior Biodegrad.* 86, 202-209, 2014. DOI: 10.1016/j.ibiod.2013.08.026
- 259.** Rao KR, Vipin A, Hariprasad P, Appaiah KA, Venkateswaran G: Biological detoxification of Aflatoxin B1 by *Bacillus licheniformis* CFR1. *Food Cont.* 71, 234-241, 2017. DOI: 10.1016/j.foodcont.2016.06.040
- 260.** Xu L, Eisa Ahmed MF, Sangare L, Zhao Y, Selvaraj JN, Xing F, Wang Y, Yang H, Liu Y: Novel aflatoxin-degrading enzyme from *Bacillus shackletonii* L7. *Toxins (Basel).* 9 (1):36, 2017. DOI: 10.3390/toxins9010036
- 261.** Shu X, Wang Y, Zhou Q, Li M, Hu H, Ma Y, Chen X, Ni J, Zhao W, Huang S, Wu L: Biological degradation of aflatoxin B(1) by cell-free extracts of *Bacillus velezensis* DY3108 with broad pH stability and excellent thermostability. *Toxins (Basel).* 10 (8):330, 2018. DOI: 10.3390/toxins10080330
- 262.** Wang L, Wu J, Liu Z, Shi Y, Liu J, Xu X, Hao S, Mu P, Deng F, Deng Y: Aflatoxin B(1) degradation and detoxification by *Escherichia coli* CG1061 isolated from chicken cecum. *Front Pharmacol.* 9:1548, 2018. DOI: 10.3389/fphar.2018.01548
- 263.** Fang Q, Du M, Chen J, Liu T, Zheng Y, Liao Z, Zhong Q, Wang L, Fang X, Wang J: Degradation and detoxification of aflatoxin B1 by tea-derived *Aspergillus niger* RAF106. *Toxins (Basel).* 12 (12):777, 2020. DOI: 10.3390/toxins12120777
- 264.** Cai M, Qian Y, Chen N, Ling T, Wang J, Jiang H, Wang X, Qi K, Zhou Y: Detoxification of aflatoxin B1 by *Stenotrophomonas* sp. CW117 and characterization the thermophilic degradation process. *Environ Pollut.* 261:114178, 2020. DOI: 10.1016/j.envpol.2020.114178
- 265.** Qiu T, Wang H, Yang Y, Yu J, Ji J, Sun J, Zhang S, Sun X: Exploration of biodegradation mechanism by AFB1-degrading strain *Aspergillus niger* FS10 and its metabolic feedback. *Food Cont.* 121:107609, 2021. DOI: 10.1016/j.foodcont.2020.107609
- 266.** Fuchs E, Binder E M, Heidler D, Krska R: Structural characterization of metabolites after the microbial degradation of type A trichothecenes by the bacterial strain BBSH 797. *Food Addit Contam.* 19 (4): 379-386, 2002. DOI: 10.1080/02652030110091154
- 267.** Young JC, Zhou T, Yu H, Zhu H, Gong J: Degradation of trichothecene mycotoxins by chicken intestinal microbes. *Food Chem Toxicol.* 45 (1): 136-143, 2007. DOI: 10.1016/j.fct.2006.07.028
- 268.** He C, Fan Y, Liu G, Zhang H: Isolation and identification of a strain of *Aspergillus tubingensis* with deoxynivalenol biotransformation capability. *Int J Mol Sci.* 9 (12): 2366-2375, 2008. DOI: 10.3390/ijms9122366
- 269.** Gao X, Mu P, Wen J, Sun Y, Chen Q, Deng Y: Detoxification of trichothecene mycotoxins by a novel bacterium, *Eggerthella* sp. DII-9. *Food Chem Toxicol.* 112, 310-319, 2018. DOI: 10.1016/j.fct.2017.12.066
- 270.** Zhai Y, Zhong L, Gao H, Lu Z, Bie X, Zhao H, Zhang C, Lu F: Detoxification of deoxynivalenol by a mixed culture of soil bacteria with 3-epi-deoxynivalenol as the main intermediate. *Front Microbiol.* 10:2172, 2019. DOI: 10.3389/fmicb.2019.02172
- 271.** Wang G, Wang Y, Ji F, Xu L, Yu M, Shi J, Xu J: Biodegradation of deoxynivalenol and its derivatives by *Devosia insulae* A16. *Food Chem.* 276, 436-442, 2019. DOI: 10.1016/j.foodchem.2018.10.011
- 272.** Ju J, Tinyiro S E, Yao W, Yu H, Guo Y, Qian H, Xie Y: The ability of *Bacillus subtilis* and *Bacillus natto* to degrade zearalenone and its application in food. *J Food Process Preserv.* 43 (10):e14122, 2019. DOI: 10.1111/jfpp.14122
- 273.** Wang Y, Wang G, Dai Y, Wang Y, Lee Y W, Shi J, Xu J: Biodegradation of deoxynivalenol by a novel microbial consortium. *Front Microbiol.* 10:2964, 2019. DOI: 10.3389/fmicb.2019.02964
- 274.** Jia R, Cao L, Liu W, Shen Z: Detoxification of deoxynivalenol by *Bacillus subtilis* ASAG 216 and characterization the degradation process. *Europ Food Res Technol.* 247 (1): 67-76, 2021. DOI: 10.1007/s00217-020-03607-8
- 275.** Xu J, Wang H, Zhu Z, Ji F, Yin X, Hong Q, Shi J: Isolation and characterization of *Bacillus amyloliquefaciens* ZDS-1: Exploring the degradation of zearalenone by *Bacillus* spp. *Food Cont.* 68, 244-250, 2016. DOI: 10.1016/j.foodcont.2016.03.030
- 276.** Wang G, Yu M, Dong F, Shi J, Xu J: Esterase activity inspired selection and characterization of zearalenone degrading bacteria *Bacillus pumilus* ES-21. *Food Cont.* 77, 57-64, 2017. DOI: 10.1016/j.foodcont.2017.01.021
- 277.** Štyriak I, Conková E, Kmec V, Böhm J, Razzazi E: The use of yeast for microbial degradation of some selected mycotoxins. *Mycotox Res.* 17 (Suppl 1): 24-27, 2001.
- 278.** Benedetti R, Nazzi F, Locci R, Firrao G: Degradation of fumonisin B1 by a bacterial strain isolated from soil. *Biodegrad.* 17 (1): 31-38, 2006. DOI: 10.1007/s10532-005-2797-y
- 279.** Lei Y, Zhao L, Ma Q, Zhang J, Zhou T, Gao C, Ji C: Degradation of zearalenone in swine feed and feed ingredients by *Bacillus subtilis* ANSB01G. *World Mycotox J.* 7 (2): 143-151, 2014. DOI: 10.3920/WMJ2013.1623
- 280.** Zhao Z, Zhang Y, Gong A, Liu N, Chen S, Zhao X, Li X, Chen L, Zhou C, Wang J: Biodegradation of mycotoxin fumonisin B1 by a novel bacterial consortium SAAS79. *Appl Microbiol Biotechnol.* 103 (17): 7129-7140, 2019. DOI: 10.1007/s00253-019-09979-6
- 281.** Stanley VG, Ojo R, Woldesenbet S, Hutchinson DH, Kubena LF: The use of *Saccharomyces cerevisiae* to suppress the effects of aflatoxicosis in broiler chicks. *Poult Sci.* 72 (10): 1867-1872, 1993. DOI: 10.3382/ps.0721867
- 282.** Diaz DE, Hagler WM, Blackwelder JT, Eve JA, Hopkins BA, Anderson KL, Jones FT, Whitlow LW: Aflatoxin binders II: Reduction of aflatoxin M1 in milk by sequestering agents of cows consuming aflatoxin in feed. *Mycopathol.* 157 (2): 233-241, 2004. DOI: 10.1023/b:mco.0000020587.93872.59
- 283.** Sampaio Baptista A, Horii J, Antonia Calori-Domingues M, Micotti da Glória E, Mastrodi Salgado J, Roberto Vizioli M: The capacity of manno-oligosaccharides, thermolysed yeast and active yeast to attenuate aflatoxicosis. *World J Microbiol Biotechnol.* 20 (5): 475-481, 2004. DOI: 10.1023/B:WIBI.0000040397.48873.3b
- 284.** Tejada-Castañeda ZI, Avila-Gonzalez E, Casaubon-Huguenin MT, Cervantes-Olivares RA, Vasquez-Pelaez C, Hernandez-Baumgarten EM, Moreno-Martinez E: Biodegradation of aflatoxin-contaminated chick feed. *Poult Sci.* 87 (8): 1569-1576, 2008. DOI: 10.3382/ps.2007-00304

- 285. Kutz RE, Sampson JD, Pompeu LB, Ledoux DR, Spain JN, Vazquez-Anon M, Rottinghaus GE:** Efficacy of Solis, NovasilPlus, and MTB-100 to reduce aflatoxin M1 levels in milk of early to mid lactation dairy cows fed aflatoxin B1. *J Dairy Sci*, 92 (8): 3959-3963, 2009. DOI: 10.3168/jds.2009-2031
- 286. Battacone G, Nudda A, Palomba M, Mazzette A, Pulina G:** The transfer of aflatoxin M1 in milk of ewes fed diet naturally contaminated by aflatoxins and effect of inclusion of dried yeast culture in the diet. *J Dairy Sci*, 92 (10): 4997-5004, 2009. DOI: 10.3168/jds.2008-1684
- 287. Hernandez-Mendoza A, Guzman-De-Peña D, González-Córdova AF, Vallejo-Córdoba B, Garcia HS:** *In vivo* assessment of the potential protective effect of Lactobacillus casei Shirota against aflatoxin B1. *Dairy Sci Technol*, 90 (6): 729-740, 2010. DOI: 10.1051/dst/2010030
- 288. Firmin S, Morgavi DP, Yiannikouris A, Boudra H:** Effectiveness of modified yeast cell wall extracts to reduce aflatoxin B1 absorption in dairy ewes. *J Dairy Sci*, 94 (11): 5611-5619, 2011. DOI: 10.3168/jds.2011-4446
- 289. Abbes S, Salah-Abbes JB, Sharafi H, Jebali R, Noghabi KA, Oueslati R:** Ability of Lactobacillus rhamnosus GAF01 to remove AFM1 *in vitro* and to counteract AFM1 immunotoxicity *in vivo*. *J Immunotoxicol*, 10 (3): 279-286, 2013. DOI: 10.3109/1547691X.2012.718810
- 290. Jebali R, Abbes S, Salah-Abbes JB, Younes RB, Haous Z, Oueslati R:** Ability of Lactobacillus plantarum MON03 to mitigate aflatoxins (B1 and M1) immunotoxicities in mice. *J Immunotoxicol*, 12 (3): 290-299, 2015. DOI: 10.3109/1547691X.2014.973622
- 291. Liu N, Wang JQ, Jia SC, Chen YK, Wang JP:** Effect of yeast cell wall on the growth performance and gut health of broilers challenged with aflatoxin B1 and necrotic enteritis. *Poult Sci*, 97 (2): 477-484, 2018. DOI: 10.3382/pss/pey342
- 292. Ali A, Khatoon A, Saleemi MK, Abbas RZ:** Aflatoxins associated oxidative stress and immunological alterations are mitigated by dietary supplementation of Pichia kudriavzevii in broiler chicks. *Microb Pathog*, 161 (Pt A):105279, 2021. DOI: 10.1016/j.micpath.2021.105279
- 293. Ali A, Khatoon A, Almohaimeed HM, Al-Sarraj F, Albiheyri R, Alotibi I, Abidin Z:** Mitigative potential of novel *Lactobacillus plantarum* TISTR 2076 against the aflatoxins-associated oxidative stress and histopathological alterations in liver and kidney of broiler chicks during the entire growth period. *Toxins (Basel)*, 14 (10):689, 2022. DOI: 10.3390/toxins14100689
- 294. Firmin S, Gandia P, Morgavi DP, Houin G, Jouany JP, Bertin G, Boudra H:** Modification of aflatoxin B1 and ochratoxin A toxicokinetics in rats administered a yeast cell wall preparation. *Food Addit Contam*, 27 (8): 1153-1160, 2010. DOI: 10.1080/19440041003801174
- 295. El Barkouky E, Mohamed F, Atta A, Abu Taleb A, El Menawey M, Hatab M:** Effect of Saccharomyces cerevisiae and vitamin C supplementation on performance of broilers subjected to ochratoxin a contamination. *Egypt Poult Sci*, 30 (1): 89-113, 2010.
- 296. Slizewska K, NowakA, Smulikowska S:** Probiotic preparation reduces faecal water genotoxicity and cytotoxicity in chickens fed ochratoxin A contaminated feed (*in vivo* study). *Acta Biochim Pol*, 63 (2): 281-286, 2016.
- 297. Mujahid H, Hashmi AS, Khan MZ, Tayyab M, Shezad W:** Protective effect of yeast sludge and whey powder against ochratoxicosis in broiler chicks. *Pak Vet J*, 39, 588-592, 2019. DOI: 10.29261/pakvetj/2019.077
- 298. Hmood KA, Habeeb AH, Al-Mhnna KI:** Antioxidant role of *Lactobacillus* sp isolated from honey bee against histological effects of ochratoxin *in vivo*. *Al-Kufa Univ J Biol*, 11 (2): 67-80, 2019.
- 299. Dazuk V, Boiago MM, Rolim G, Paravisi A, Copetti PM, Bissacotti BF, Morsch VM, Vedovatto M, Gazoni FL, Matte F, Gloria EM, Da Silva AS:** Laying hens fed mycotoxin-contaminated feed produced by Fusarium fungi (T-2 toxin and fumonisin B1) and *Saccharomyces cerevisiae* lysate: Impacts on poultry health, productive efficiency, and egg quality. *Microb Pathog*, 149:104517, 2020. DOI: 10.1016/j.micpath.2020.104517
- 300. Alberts JE, Gelderblom WC, Botha A, van Zyl WH:** Degradation of aflatoxin B(1) by fungal laccase enzymes. *Int J Food Microbiol*, 135 (1): 47-52, 2009. DOI: 10.1016/j.ijfoodmicro.2009.07.022
- 301. Taylor MC, Jackson CJ, Tattersall DB, French N, Peat TS, Newman J, Briggs LJ, Lopalikar GV, Campbell PM, Scott C:** Identification and characterization of two families of F420H2-dependent reductases from mycobacteria that catalyse aflatoxin degradation. *Mol Microbiol*, 78 (3): 561-575, 2010. DOI: 10.1111/j.1365-2958.2010.07356.x
- 302. Cao H, Liu D, Mo X, Xie C, Yao D:** A fungal enzyme with the ability of aflatoxin B(1) conversion: purification and ESI-MS/MS identification. *Microbiol Res*, 166 (6): 475-483, 2011. DOI: 10.1016/j.micres.2010.09.002
- 303. Zhao LH, Guan S, Gao X, Ma QG, Lei YP, Bai XM, Ji C:** Preparation, purification and characteristics of an aflatoxin degradation enzyme from *Myxococcus fulvus* ANSM068. *J Appl Microbiol*, 110 (1): 147-155, 2011. DOI: 10.1111/j.1365-2672.2010.04867.x
- 304. Wang J, Ogata M, Hirai H, Kawagishi H:** Detoxification of aflatoxin B1 by manganese peroxidase from the white-rot fungus *Phanerochaete sordida* YK-624. *FEMS Microbiol Lett*, 314 (2): 164-169, 2011. DOI: 10.1111/j.1574-6968.2010.02158.x
- 305. Yehia RS:** Aflatoxin detoxification by manganese peroxidase purified from *Pleurotus ostreatus*. *Braz J Microbiol*, 45 (1): 127-133, 2014. DOI: 10.1590/S1517-83822014005000026
- 306. Ito M, Sato I, Ishizaka M, Yoshida S, Koitabashi M, Yoshida S, Tsushima S:** Bacterial cytochrome P450 system catabolizing the Fusarium toxin deoxynivalenol. *Appl Environ Microbiol*, 79 (5): 1619-1628, 2013. DOI: 10.1128/AEM.03227-12
- 307. Feltrin ACP, Garcia SO, Caldas SS, Primel EG, Badiale-Furlong E, Garda-Buffon J:** Characterization and application of the enzyme peroxidase to the degradation of the mycotoxin DON. *J Environ Sci Health*, 52 (10): 777-783, 2017. DOI: 10.1080/03601234.2017.1356672
- 308. Carere J, Hassan YI, Lepp D, Zhou T:** The enzymatic detoxification of the mycotoxin deoxynivalenol: identification of DepA from the DON epimerization pathway. *Microb Biotechnol*, 11 (6): 1106-1111, 2018. DOI: 10.1111/1751-7915.12874
- 309. He WJ, Shi MM, Yang P, Huang T, Zhao Y, Wu AB, Dong WB, Li HP, Zhang JB, Liao YC:** A quinone-dependent dehydrogenase and two NADPH-dependent aldo/keto reductases detoxify deoxynivalenol in wheat via epimerization in a Devosia strain. *Food Chem*, 321:126703, 2020. DOI: 10.1016/j.foodchem.2020.126703
- 310. Tso KH, Lumsangkul C, Ju JC, Fan YK, Chiang HI:** The potential of peroxidases extracted from the spent mushroom (*Flammulina velutipes*) substrate significantly degrade mycotoxin deoxynivalenol. *Toxins*, 13 (1):72, 2021. DOI: 10.3390/toxins13010072
- 311. Shcherbakova L, Rozhkova A, Osipov D, Zorov I, Mikityuk O, Statsyuk N, Sinitysyna O, Dzhavakhya V, Sinitsyn A:** Effective zearalenone degradation in model solutions and infected wheat grain using a novel heterologous lactonohydrolase secreted by recombinant *Penicillium canescens*. *Toxins (Basel)*, 12 (8):475, 2020. DOI: 10.3390/toxins12080475
- 312. Azam MS, Yu D, Liu N, Wu A:** Degrading ochratoxin A and zearalenone mycotoxins using a multifunctional recombinant enzyme. *Toxins (Basel)*, 11 (5): 301, 2019. DOI: 10.3390/toxins11050301
- 313. Masching S, Naehrer K, Schwartz-Zimmermann HE, Sarandan M, Schaumberger S, Dohnal I, Nagl V, Schatzmayr D:** Gastrointestinal degradation of fumonisins B(1) by carboxylesterase FumD prevents fumonisins induced alteration of sphingolipid metabolism in turkey and swine. *Toxins (Basel)*, 8 (3):84, 2016. DOI: 10.3390/toxins8030084
- 314. Ahmad S, Humak F, Ahmad M, Altaf H, Qamar W, Hussain A, Ashraf U, Abbas RZ, Siddique A, Ashraf T:** Phytochemicals as alternative anthelmintics against poultry parasites: A review. *Agrobiol Rec*, 12, 34-45, 2023. DOI: 10.47278/journal.abr/2023.015
- 315. Hegazy SA, Abd ES, Khorshed M, Salem F:** Productive and immunological performance of small ruminants offered some medicinal plants as feed additives. *Int J Vet Sci*, 12, 120-125, 2023. DOI: 10.47278/journal.ivs/2022.163
- 316. Krishnaveni P, Thangapandian M, Raja P, Rao G:** Pathological and molecular studies on antitumor effect of curcumin and curcumin solid lipid nanoparticles. *Pak Vet J*, 43, 315-320, 2023. DOI: 10.29261/pakvetj/2023.022
- 317. Ratajac R, Pavlićević A, Petrović J, Stojanov I, Orčić D, Štrbac F, Simin N:** *In vitro* evaluation of acaricidal efficacy of selected essential oils against *Dermanyssus gallinae*. *Pak Vet J*, 44, 93-98, 2023. DOI: 10.29261/pakvetj/2023.123

- 318. Mohammad LM, Kamil AM, Tawfeeq RK, Jamal Ahmed S:** Ameliorating effects of herbal mixture for dexamethasone induced histological changes in mice. *Int J Vet Sci*, 12, 126-131, 2023. DOI: 10.47278/journal.ijvs/2022.170
- 319. Rasheed M, Aljohani AS:** Evaluation of anthelmintic effects of essential oil of Star Anise against Ascaridia galli of poultry. *Pak Vet J*, 44, 1223-1228, 2024. DOI: 10.29261/pakvetj/2024.294
- 320. Saeed Z, Abbas RZ, Khan MK, Saleemi MK:** Anticoccidial activities of essential oil of Amomum subulatum in broiler chicks. *Pak J Agric Sci*, 60, 377-384, 2023.
- 321. Shi D, Zhou J, Zhao L, Rong X, Fan Y, Hamid H, Li W, Ji C, Ma Q:** Alleviation of mycotoxin biodegradation agent on zearalenone and deoxynivalenol toxicosis in immature gilts. *J Anim Sci Biotechnol*, 9 (1):42, 2018. DOI: 10.1186/s40104-018-0255-z
- 322. Abbas R Z, Saeed Z, Bosco A, Qamar W, Subhani Z, Sorin CM, Kasli MAF, Munir F:** Botanical control of coccidiosis in ruminants. *Pak J Agric Sci*, 60, 473-485, 2023.
- 323. Liu L, Xie M, Wei D:** Biological detoxification of mycotoxins: Current status and future advances. *Int J Mol Sci*, 23 (3): 1064, 2022. DOI: 10.3390/ijms23031064
- 324. Ahmad S, Yousaf MS, Tahir SK, Rashid MA, Majeed KA, Naseem M, Raza M, Hayat Z, Khalid A, Zaneb H:** Effects of co-supplementation of β -galacto-oligosaccharides and methionine on breast meat quality, meat oxidative stability and selected meat quality genes in broilers. *Pak Vet J*, 43, 428-434, 2023. DOI: 10.29261/pakvetj/2023.043
- 325. Zhao Y, Wang Q, Huang J, Chen Z, Liu S, Wang X, Wang F:** Mycotoxin contamination and presence of mycobiota in rice sold for human consumption in China. *Food Cont*, 98, 19-23, 2019. DOI: 10.1016/j.foodcont.2018.11.014
- 326. Aboubakr M, Elmahdy AM, Taima S, Emam MA, Farag A, Alkafafy M, Said AM, Soliman A:** Protective effects of N acetylcysteine and vitamin E against acrylamide-induced neurotoxicity in rats. *Pak Vet J*, 43, 262-268, 2023.
- 327. El-Sheikh ESA, Hamed IA, Alduwish MA, Momenah MA, Melebary SJ, Alsolmy SA, Alghamdi MS, Alharbi AA, Sherif RM, Shalaby AA:** The ameliorative effect of vitamin C against sub-chronic thiamethoxam toxicity in male rats. *Pak Vet J*, 44, 803-811, 2024. DOI: 10.29261/pakvetj/2024.206
- 328. Gradelet S, Le Bon AM, Berges R, Suschetet M, Astorg P:** Dietary carotenoids inhibit aflatoxin B1-induced liver preneoplastic foci and DNA damage in the rat: role of the modulation of aflatoxin B1 metabolism. *Carcinog*, 19 (3): 403-411, 1998. DOI: 10.1093/carcin/19.3.403
- 329. Klein PJ, Van Vleet TR, Hall JO, Coulombe RA, Jr:** Dietary butylated hydroxytoluene protects against aflatoxicosis in Turkeys. *Toxicol Appl Pharmacol*, 182 (1): 11-19, 2002. DOI: 10.1006/taap.2002.9433
- 330. Sahoo PK, Mukherjee SC:** Immunomodulation by dietary vitamin C in healthy and aflatoxin B1-induced immunocompromised rohu (*Labeo rohita*). *Comp Immunol Microbiol Infect Dis*, 26 (1): 65-76, 2003. DOI: 10.1016/s0147-9571(01)00038-8
- 331. Tedesco D, Steidler S, Galletti S, Tameni M, Sonzogni O, Ravarotto L:** Efficacy of silymarin-phospholipid complex in reducing the toxicity of aflatoxin B1 in broiler chicks. *Poult Sci*, 83 (11): 1839-1843, 2004. DOI: 10.1093/ps/83.11.1839
- 332. Karakilcik AZ, Zerin M, Arslan O, Nazligul Y, Vural H:** Effects of vitamin C and E on liver enzymes and biochemical parameters of rabbits exposed to aflatoxin B1. *Vet Hum Toxicol*, 46 (4): 190-192, 2004.
- 333. Solcan C, Gogu M, Floristean V, Oprisan B, Solcan G:** The hepatoprotective effect of sea buckthorn (*Hippophae rhamnoides*) berries on induced aflatoxin B1 poisoning in chickens 1. *Poult Sci*, 92 (4): 966-974, 2013. DOI: 10.3382/ps.2012-02572
- 334. Li Y, Ma QG, Zhao LH, Guo YQ, Duan GX, Zhang JY, Ji C:** Protective efficacy of alpha-lipoic acid against aflatoxin B1-induced oxidative damage in the liver. *Asian-Aust J Anim Sci*, 27 (6): 907-915, 2014. DOI: 10.5713/ajas.2013.13588
- 335. Sridhar M, Suganthi RU, Thammiah V:** Effect of dietary resveratrol in ameliorating aflatoxin B1-induced changes in broiler birds. *J Anim Physiol Anim Nutr (Berl)*, 99 (6): 1094-1104, 2015. DOI: 10.1111/jpn.12260
- 336. Ghadiri S, Spalenza V, Dellafiora L, Badino P, Barbarossa A, Dall'Asta C, Nebbia C, Girolami F:** Modulation of aflatoxin B1 cytotoxicity and aflatoxin M1 synthesis by natural antioxidants in a bovine mammary epithelial cell line. *Toxicol In Vitro*, 57, 174-183, 2019. DOI: 10.1016/j.tiv.2019.03.002
- 337. Cheng P, Ishfaq M, Yu H, Yang Y, Li S, Li X, Fazlani SA, Guo W, Zhang X:** Curcumin ameliorates duodenal toxicity of AFB1 in chicken through inducing P-glycoprotein and downregulating cytochrome P450 enzymes. *Poult Sci*, 99 (12): 7035-7045, 2020. DOI: 10.1016/j.psj.2020.09.055
- 338. Atroschi F, Rizzo A, Biese I, Salonen M, Lindberg LA, Saloniemi H:** Effects of feeding T-2 toxin and deoxynivalenol on DNA and GSH contents of brain and spleen of rats supplemented with vitamin E and C and selenium combination. *J Anim Physiol Anim Nutr*, 74 (1-5): 157-164, 1995. DOI: 10.1111/j.1439-0396.1995.tb00447.x
- 339. Wu L, Liao P, He L, Feng Z, Ren W, Yin J, Duan J, Li T, Yin Y:** Dietary L-arginine supplementation protects weanling pigs from deoxynivalenol-induced toxicity. *Toxins (Basel)*, 7 (4): 1341-1354, 2015. DOI: 10.3390/toxins7041341
- 340. Lu Z:** Dose-dependent fumonisin B (1) hepatotoxicity and hepatocarcinogenicity, detoxification of fumonisin B (1), and suppression by isoflavones of fumonisin B (1)-promoted hepatocarcinogenesis in rats. *Iowa State University*. 1997.
- 341. Gbore FA, Adu OA:** Ameliorative potential of vitamin E on the impact of dietary fumonisins B1 on reproductive performance of female rabbits. *J Agric Rural Develop Trop Subtrop*, 118 (2): 161-169, 2017.
- 342. Ledur PC, Santurio JM:** Cytoprotective effects of curcumin and silymarin on PK-15 cells exposed to ochratoxin A, fumonisin B(1) and deoxynivalenol. *Toxicon*, 185, 97-103, 2020. DOI: 10.1016/j.toxicon.2020.06.025
- 343. Bose S, Sinha SP:** Modulation of ochratoxin-produced genotoxicity in mice by vitamin C. *Food Chem Toxicol*, 32 (6): 533-537, 1994. DOI: 10.1016/0278-6915(94)90110-4
- 344. Hoehler D, Marquardt RR:** Influence of vitamins E and C on the toxic effects of ochratoxin A and T-2 toxin in chicks. *Poult Sci*, 75 (12): 1508-1515, 1996. DOI: 10.3382/ps.0751508
- 345. Grosse Y, Chekir-Ghedira L, Huc A, Obrecht-Pflumio S, Dirheimer G, Bacha H, Pfohl-Leszkoicz A:** Retinol, ascorbic acid and alpha-tocopherol prevent DNA adduct formation in mice treated with the mycotoxins ochratoxin A and zearalenone. *Cancer Lett*, 114 (1-2): 225-229, 1997. DOI: 10.1016/s0304-3835(97)04669-7
- 346. Atroschi F, Biese I, Saloniemi H, Ali-Vehmas T, Saari S, Rizzo A, Veijalainen P:** Significance of apoptosis and its relationship to antioxidants after ochratoxin A administration in mice. *J Pharm Pharm Sci*, 3 (3): 281-91, 2000.
- 347. Shalaby AM:** The opposing effect of ascorbic acid (vitamin C) on ochratoxin toxicity in Nile tilapia (*Oreochromis niloticus*). 2004.
- 348. Zhai S, Ruan D, Zhu Y, Li M, Ye H, Wang W, Yang L:** Protective effect of curcumin on ochratoxin A-induced liver oxidative injury in duck is mediated by modulating lipid metabolism and the intestinal microbiota. *Poult Sci*, 99 (2): 1124-1134, 2020. DOI: 10.1016/j.psj.2019.10.041
- 349. Leal M, Shimada A, Ruiz F, Gonzalez de Mejia E:** Effect of lycopene on lipid peroxidation and glutathione-dependent enzymes induced by T-2 toxin *in vivo*. *Toxicol Lett*, 109 (1-2): 1-10, 1999. DOI: 10.1016/s0378-4274(99)00062-4
- 350. Ghedira-Chekir L, Maaroufi K, Zakhama A, Ellouz F, Dhouib S, Creppy EE, Bacha H:** Induction of a SOS repair system in lysogenic bacteria by zearalenone and its prevention by vitamin E. *Chem Biol Interact*, 113 (1): 15-25, 1998. DOI: 10.1016/s0009-2797(98)00013-1
- 351. Shi B, Su Y, Chang S, Sun Y, Meng X, Shan A:** Vitamin C protects piglet liver against zearalenone-induced oxidative stress by modulating expression of nuclear receptors PXR and CAR and their target genes. *Food Funct*, 8 (10): 3675-3687, 2017. DOI: 10.1039/C7FO01301A
- 352. Gao X, Xiao ZH, Liu M, Zhang NY, Khalil MM, Gu CQ, Qi DS, Sun LH:** Dietary silymarin supplementation alleviates zearalenone-induced hepatotoxicity and reproductive toxicity in rats. *J Nutr*, 148 (8): 1209-1216, 2018. DOI: 10.1093/jn/nxy114
- 353. Su Y, Sun Y, Ju D, Chang S, Shi B, Shan A:** The detoxification effect of vitamin C on zearalenone toxicity in piglets. *Ecotoxicol Environ Saf*, 158, 284-292, 2018. DOI: 10.1016/j.ecoenv.2018.04.046