

REVIEW ARTICLE

Potential Application of *Rhododendron* Flavonoids' Anti-inflammatory and Antioxidant Activities in Animal Health Support and Protection

Ziqin ZHANG¹ , Sheng KANG¹ , Meixue NI¹ , Xiaoqing GUO¹ , Hailong DONG¹ ,
Qingxia WU^{1(*)} , Zhenyu CHANG^{1(*)} 

¹ Xizang Agricultural and Animal Husbandry University, School of Animal Science, Tibetan Key Laboratory of Clinical Veterinary Medicine, Linzhi, Tibet, P. R. CHINA



(*) **Corresponding author:**

Qingxia Wu & Zhenyu Chang
Phone: +86-17689548312 (ZC),
+86-18089945473 (QW)
E-mail: zychang2014@sina.com (ZC);
441594343@qq.com (QW)

How to cite this article?

Zhang Z, Kang S, Ni M, Guo X, Dong H, Wu Q, Chang Z: Potential Application of *Rhododendron* Flavonoids' Anti-inflammatory and Antioxidant Activities in Animal Health Support and Protection. *Kafkas Univ Vet Fak Derg*, 2026 (Article in Press).
DOI: 10.9775/kvfd.2025.35766

Article ID: KVFD-2025-35766

Received: 25.11.2025

Accepted: 18.02.2026

Published Online: 29.03.2026

Abstract

Rhododendron plants are abundant in flavonoid compounds, whose prominent antioxidant and anti-inflammatory physiological activities have attracted considerable attention. However, the application of related research in animal health maintenance and bodily protection, along with their underlying mechanisms, remains unclear. This ambiguity has constrained their translation into practical applications as physiological protective factors within animal husbandry. This study systematically reviewed relevant literature, identifying 17 *Rhododendron* flavonoids exhibiting significant antioxidant and anti-inflammatory activity. The antioxidant effects of these flavonoids are primarily achieved through multiple pathways: direct scavenging of free radicals, inhibition of oxidative enzymes (such as NADPH oxidase and xanthine oxidase), chelation of metal ions, supply of hydrogen ions, enhancement of the antioxidant system, and regulation of endogenous antioxidant capacity via the Keap1-Nrf2/ARE and MAPKs signaling pathways. Meanwhile, their anti-inflammatory effects are mediated through four main mechanisms: inhibition of inflammatory mediator production, modulation of cytokines and their receptors, scavenging of reactive oxygen species, and regulation of cell signaling pathways (including NF- κ B and AP-1). To date, flavonoids have demonstrated significant efficacy in the treatment of various human diseases, including hypertension, atherosclerosis, cardiomyopathy, myocarditis, diabetes mellitus, and cancer. However, their potential in veterinary medicine has not been fully explored. This review summarizes recent advances in the antioxidant and anti-inflammatory physiological activities of *Rhododendron* flavonoids and their underlying mechanisms. It aims to provide scientific rationale for developing flavonoid-based physiological protective agents suitable for animals, thereby advancing their scientific application in animal health management and livestock production. This aligns with current priority areas in veterinary research.

Keywords: Antioxidant, Anti-inflammatory, Animal health support and protection, Flavonoids, *Rhododendron* plants

INTRODUCTION

Rhododendron belongs to the *Rhododendron* family, which is a large genus of *Rhododendron*, with common types such as Ying shan hong, Haunted Goat Flower, Zhao shan White, Goat Tramp, and so on, about 967 species, which are widely distributed in Asia, Europe, and North America, and are mainly native to East Asia and Southeast Asia^[1]. About 700 species in China, except Xinjiang, Ningxia, everywhere, but in the southwest provinces and districts, mostly^[2]. The flowers, leaves and stems of *Rhododendron* plants are rich in flavonoids, which are high quality herbal medicines. Flavonoids have excellent antioxidant and anti-inflammatory activities and therefore also have many pharmacological effects

such as anti-cancer, treatment of cardiovascular diseases and diabetes mellitus^[3].

The development of many diseases is closely related to oxidative damage in the body, and reactive oxygen species (ROS) and reactive nitrogen species (RNS) free radicals are naturally produced during normal metabolism, and they play an important role in physiological activities. However, when the concentration of ROS/RNS is too high, they produce oxidative stress, a state that leads to tissue damage and has been linked to the development of a variety of diseases, including cancer, cardiovascular disease, neurodegenerative diseases, diabetes, respiratory infections, and others^[4]. The high cost of treating related diseases caused by oxidative damage places a huge economic burden on the global community. Flavonoids



possess diverse physiological activities and low toxicity. Therefore, investigating their antioxidant and anti-inflammatory mechanisms to elucidate their protective effects in alleviating oxidative stress and mitigating cellular oxidative damage holds significant value for safeguarding animal organisms from injury, maintaining health homeostasis. Clarifying the antioxidant and anti-inflammatory mechanisms of these flavonoids may also facilitate the development of physiologically protective formulations, providing theoretical support for standardised animal health management^[5].

Regarding the antioxidant and anti-inflammatory activities of flavonoid compounds, existing research has predominantly focused on their pharmacological effects in humans, while studies on their physiological support applications, pharmacokinetics, and safety in animals -particularly livestock and companion animals- remain fragmented and lack systematic integration. To address this research gap, this paper first provides a comprehensive review of studies concerning the antioxidant and anti-inflammatory physiological properties of flavonoids. It systematically outlines their mechanisms of action and applications in maintaining animal health and providing physiological protection, thereby establishing a foundational reference for subsequent research into the application of *Rhododendron* flavonoids in animal health management and livestock production. Subsequently, the focus shifts to the physiological protective value of *Rhododendron* flavonoids in veterinary health management. This involves systematically reviewing relevant domestic and international research findings, thoroughly elucidating their mechanisms of action, and examining their current applications in maintaining animal health and protecting against damage. This aims to establish a robust foundation for advancing research in this field, construct a standardised reference framework, and facilitate the translational application of flavonoids (particularly represented by *Rhododendron* flavonoids) in veterinary health management and livestock production.

FLAVONOID COMPOSITION

In medicinal chemistry, a compound's chemical formula not only determines its physicochemical properties, but more crucially, the presence and positioning of specific functional groups within its structure directly dictate how it interacts with biological targets. This, in turn, influences its pharmacological activity and therapeutic applications. Consequently, understanding a compound's chemical formula forms the foundation for investigating its mode of action. The research on flavonoids of *Rhododendron* plants in China was first started in 1974, when the total flavonoid glycosides isolated from *Rhododendron* plants were acid-hydrolysed by the teaching and research group

of pharmacology at Lanzhou Medical College, from which 3,5,7,8,3',4'-hexahydroxyflavonoids were isolated^[6]. Nowadays, flavonoids are mainly detected and analysed by ultra-high performance liquid chromatography-quadrupole time-of-flight mass spectrometry (UHPLC-TOFMS) and various chromatographic methods^[7]. To date, 17 flavonoid compounds with antioxidant or anti-inflammatory activity have been studied (*Table 1*).

ANTIOXIDANT MECHANISM OF ACTION

Antioxidant and oxidative stress are inextricably linked to the occurrence of oxidative stress, which is a state in which excessive oxidative molecules (e.g. ROS) are produced or accumulated within an organism, exceeding the antioxidant capacity of the cells or tissues, resulting in oxidative damage to biomolecules (e.g. proteins, lipids, DNA, etc.)^[20], and antioxidant refers to the process of scavenging oxygen free radicals, thereby protecting cells from oxidative stress. Therefore, as illustrated in *Fig. 1*, an understanding of the factors that induce oxidative stress in cells and the subsequent application of antioxidants to mitigate these triggers enables the attainment of effective antioxidant protection. Flavonoids, as an excellent antioxidant, can address oxidative stress through several pathways, such as stabilisation or reduction of reactive oxygen species (ROS), scavenging of free radicals, and modulation of endogenous antioxidant capacity are the three main pathways.

Stabilises or Reduces ROS, Scavenges Free Radicals

The two pathways, stabilisation or reduction of ROS and scavenging of free radicals, usually occur simultaneously in the process of flavonoids exerting antioxidant effects^[21]. Thus, the two pathways are described together. The antioxidant modalities of both pathways are direct scavenging of free radicals, inhibition of oxidative enzymes (NADPH oxidase, xanthine oxidase), chelation of metal ions, provision of hydrogen ions and enhancement of the antioxidant system.

Direct Scavenging of Free Radicals

Free radicals are atoms that exist in free form without pairs or free electrons, which can be produced by normal cellular metabolism^[22]. Common types of free radicals include hydroxyl radicals (HO·), superoxide anion radicals (O₂^{·-}) and hydrogen peroxide radicals (HO₂·). Hydroxyl radicals (HO·) are the most active free radicals in the cell, which can react with lipids, proteins, and DNA, triggering the oxidation of unsaturated fatty acids and the formation of lipid peroxides (LPO), which can disrupt the membrane structure and lead to cell damage^[23]. As illustrated in *Fig. 2-a*, quercetin, isoquercetin, rutin and other azalea plant flavonoids were able to directly scavenge HO· at concentrations up to 260 μM^[24]. When


















Table 1. Flavonoid constituents of <i>Rhododendron</i> spp. with antioxidant and anti-inflammatory effects				
Serial Number	Phyto-Chemicals	Source Plant	Chemical Structure	Ref.
1	Quercetin	Almost all <i>Rhododendron</i> plants		[8]
2	Rutin	<i>Rhododendron arboreum</i>		[9]
3	Catechin	<i>Rhododendron nivale</i> Hook <i>Rhododendron dauricum</i> <i>Rhododendron wiltonii</i> Hemsl		[10,11]
4	Hyperin	Almost all <i>Rhododendron</i> plants		[8]
5	Farrerol	<i>Rhododendron primuliflorum</i> Bureau <i>Rhododendron dauricum</i>		[12,13]
6	Morin-3-O-β-L-lyxoside	<i>Rhododendron nivale</i> Hook		[10]
7	Epicatechin	<i>Rhododendron nivale</i> Hook		[10]
8	dihydroquercetin	<i>Rhododendron nivale</i> Hook		[10]

Table 1. Continue				
Serial Number	Phyto-Chemicals	Source Plant	Chemical Structure	Ref.
9	Naringenin	<i>Rhododendron pachypodum</i> Balf		[14]
10	Kaempferol	<i>Rhododendron dauricum</i> <i>Rhododendron wiltonii</i> Hemsl		[11,13]
11	Myricetin	<i>Rhododendron dauricum</i> <i>Rhododendron anthopogon</i> D		[15]
12	Luteolin	<i>Rhododendron anthopogonoides</i> Maxim		[16,17]
13	Apigenin	<i>Rhododendron anthopogonoides</i> Maxim <i>Rhododendron amesiae</i> Rehder		[18]
14	4-O-Methylcedrin	<i>Rhododendron pachypodum</i> Balf		[14]
15	(2Z)-2,6 dimethyl-2,7-octadiene-1,6-diol	<i>Rhododendron pachypodum</i> Balf		[14]
16	6,8-di-C-methyl-dihydrokaempferol	<i>Rhododendron pachypodum</i> Balf		[14]
17	Isoquercitrin	<i>Rhododendron amesiae</i> Rehder		[19]

flavonoids bind to free radicals, the phenolic hydroxyl group of the flavonoid can provide an electron to the free radical, thus neutralising and deactivating it. When free

radicals are scavenged, lipids, proteins, DNA and other biomolecules in the cell can exert their effects normally, reducing oxidative damage [25].

Inhibition of Oxidative Enzymes (NADPH Oxidase, Xanthine Oxidase)

Both NADPH oxidase and xanthine oxidase catalyse free radical generation. NADPH oxidase is a transmembrane enzyme complex consisting of several components such as NOX2, p22phox, p47phox, p67phox, etc., whose main function is the production of superoxide anion radicals ($O_2^{\cdot-}$). In phagocytes, such as neutrophils, NADPH oxidase is activated upon detection of pathogens, which, through the signalling pathway, leads to the production of cytokines (mainly p47phox) phosphorylation, prompting their transfer to membrane-bound NOX2 components and initiating the catalytic production of superoxide anion, the catalytic mechanism of which involves the transfer of electrons from NADPH to flavin adenine dinucleotide (FAD), then to proximal haemoglobin, rapidly to distal haemoglobin, and ultimately to molecular oxygen, to form superoxide anion [26]. Xanthine oxidase (XO), in catalysing the conversion of hypoxanthine to xanthine and further to uric acid, uses molecular oxygen as an electron acceptor to produce large amounts of uric acid and H_2O_2 , of which H_2O_2 can be further decomposed to produce hydroxyl radicals ($\cdot OH$), increasing the production of reactive oxygen species [27].

Both quercetin and lignans from the flavonoids of the genus *Rhododendron* showed inhibition of NADPH oxidase and xanthine oxidase. Quercetin inhibits NADPH oxidase

activity and reduces ROS production, while it also inhibits xanthine oxidase activity, lowers serum uric acid levels and reduces H_2O_2 production [28]. Lignans also inhibit NADPH oxidase, reduce oxidative stress, and can reverse the elevation of xanthine oxidase activity induced by potassium oxonate, reduce serum and liver levels of uric acid and blood urea nitrogen, and reduce H_2O_2 production [29]. As illustrated in Fig. 2-b, populin acts on NADPH oxidase, and in silico predictions reveal a high docking score between populin and this enzyme, suggesting its potential to inhibit NADPH oxidase [30]. As illustrated in Fig 2-c, rutin and chrysin exert their primary inhibitory effects on xanthine oxidase; notably, rutin possesses diverse biological activities including antioxidant and anticancer properties, while chrysin inhibits xanthine oxidase activity via hydrogen bonding and van der Waals forces with the enzyme's catalytic active site, acting in a mixed competitive manner [31]. These flavonoids exert their antioxidant and protective effects by inhibiting the activity of oxidative enzymes and reducing oxidative stress and uric acid production.

Chelating Metal Ions

The mechanism of action of flavonoids chelating metal ions to achieve antioxidant action mainly includes the following aspects. The first is that flavonoids form stable complexes through the formation of ligand bonds with metal ions (Fe, Cu, etc.) through specific functional groups such as hydroxyl ($-OH$) and carbonyl ($C=O$)




Fig 1. Several ways in which cells generate oxidative stress. (a) The reduction of oxygen to water produces ROS substances such as $O_2^{\cdot-}$ and NO, and the production of ROS causes oxidative damage to cells, (b) An imbalance in the antioxidant defence system increases ROS and swamps AOS, (c) Activation of NADPH oxidase leads to excessive ROS production, causing oxidative stress, (d) XOD catalyses the production of uric acid and H_2O_2 under aerobic conditions, and H_2O_2 is broken down to $HO\cdot$, loss of cells, (e) Metal ions catalyse the production of ROS by oxidative enzymes, (f) Metal ions such as iron and copper catalyse oxidative reactions that produce $HO\cdot$, causing cell damage




Fig 2. Azalea flavonoids as antioxidants through ROS stabilisation and free radical scavenging. (a) Quercetin, isoquercetin, and rutin directly scavenge HO \cdot , (b) Quercetin, lignocerotin, and populin inhibit NADPH oxidase activity and reduce free radical production, (c) Quercetin, lignans, rutin, and chrysin inhibit xanthine oxidase activity and reduce free radical production, (d) Quercetin, lignans, rutin, apigenin and kaempferol chelate metal ions, reduce HO \cdot -production and inhibit oxidase activity, (e) Quercetin, lignans, apigenin, and kaempferol provide hydrogen ions to scavenge HO \cdot , (f) Flavonoids enhance the antioxidant capacity of the antioxidant system and stabilise ROS

groups in their molecular structure, and such complexes have stronger free radical scavenging ability than free flavonoids, in which metal ions can facilitate the transfer of electrons to enhance the capture of free radicals; finally, there may be synergistic antioxidant activity between flavonoids and metal ions in particular cases [32,33]. For example, complexes formed by flavonoids with copper ions show synergistic effects in scavenging superoxide anion (O $_2^{\cdot-}$) and hydroxyl radical (OH \cdot) [33].

As illustrated in Fig. 2-d, quercetin, rutin, apigenin, lignans, and kaempferol in the flavonoids of the genus *Rhododendron* can form complexes with metal ions and exhibit antioxidant activity [34].

Hydrogen Atom

Flavonoids provide hydrogen ions to scavenge free radicals in two main ways. The first is that the antioxidant activity of flavonoids interrupts the chain reaction mainly through the reaction of their phenolic hydroxyl groups with oxygen radicals to form resonance-stabilised semiquinone radicals. In this process, hydrogen atoms on the phenolic hydroxyl groups of flavonoids can combine with peroxy radicals to form flavonoid radicals, which in turn react with other radicals, thus terminating the free radical chain reaction. The second is that intramolecular hydrogen bonding in flavonoids reduces the antioxidant activity of hydroxyl groups that act as hydrogen bond donors (e.g., 5-OH, 3-OH, and 3'-OH) while enhancing

the antioxidant activity of hydroxyl groups that act as hydrogen bond acceptors (e.g., 4'-OH) [35].

As illustrated in Fig. 2-e, quercetin, populin and kaempferol among the flavonoids of the genus *Azalea* exhibit antioxidant activity by providing hydrogen atoms to scavenge free radicals [24], and the antioxidant activities of populin and kaempferol were also positively correlated with the number of phenolic hydroxyl groups on the B-ring [36].

Antioxidant Defence System

The antioxidant system is mainly composed of a number of antioxidant enzymes, non-enzymatic antioxidants, and low molecular weight antioxidants [37]. Antioxidant enzymes are the first-line defence mechanisms of the antioxidant defence system, including superoxide dismutase (SOD), catalase (CAT), glutathione peroxidase (GPx), etc. These enzymes are able to scavenge reactive oxygen species, such as superoxide radicals and hydrogen peroxide, to prevent them from causing cellular damage [37]. Non-enzymatic antioxidants, including reduced glutathione (GSH), coenzyme Q (CoQ), lipoic acid, uric acid, and various proteins (e.g. ferritin, transferrin, cuprocyenin, and albumin), which are able to directly neutralise free radicals and protect cells from oxidative damage [38]. Low molecular weight antioxidants such as vitamin C, vitamin E, polyphenols, coenzyme Q or the metabolic compound urate, which scavenge free radicals and reduce oxidative stress [37].

As illustrated in *Fig. 2-f*, flavonoids reduce oxidative stress by modulating the antioxidant enzyme system, enhancing the activity of antioxidant enzymes and the antioxidant capacity of the AsA-GSH cycle, and are also able to stimulate the ascorbic acid biosynthetic pathway and its upstream glycolytic metabolic pathway, leading to an increased accumulation of flavonoids, which act as one of the antioxidants in the non-enzymatic system, providing hydroxyl radicals to scavenge excess reactive oxygen species^[39]. It can also act synergistically with vitamin E, partially regenerating in the presence of combined antioxidants, driving the regeneration process by the scavenging of semiquinone radicals in a synergistic reaction, and also by interactions with glutathione and ascorbic acid, in which oxidised flavonoids can be re-reduced by glutathione and ascorbic acid to continue to exert their antioxidant effects^[40].

Modulates Endogenous Antioxidant Capacity

Azalea flavonoids regulate endogenous antioxidant capacity and enhance antioxidant capacity mainly through activation of Keap1-Nrf2/ARE and MAPK_S signalling pathways^[10].

Antioxidant of Keap1-Nrf2/ARE Signalling Pathway

Nuclear transcription factor E2-related factor 2 (Nrf2) is a member of the Cap'n'collar (CNC) family of regulatory proteins, and is a transcription factor with a basic leucine zip structure, which is widely found in various organs of the body, and is mainly responsible for regulating cellular redox reactions^[41]. Keap1 is a Cullin3 (Cul3)-dependent substrate articulating protein of the E3 ubiquitin ligase complex that assembles

with Cul3 and Rbx1 to form a functional E3 ubiquitin ligase complex (Keap1-Cul3-E3), which in turn regulates Nrf2^[42]. As illustrated in *Fig. 3-a*, under normal physiological conditions, Nrf2 binds to its endogenous inhibitor Kelch ECH-related protein 1 (Keap1), exists in an inactive state in the cytosol and is rapidly degraded via the ubiquitin proteasome pathway in order to maintain the low transcriptional activity of Nrf2 under physiological conditions^[43]. As illustrated in *Fig. 3-b*, upon stimulation of cells by reactive oxygen species (ROS) or other nucleophiles, Nrf2 dissociates from Keap1, undergoes activation and translocates into the nucleus. In the nucleus, Nrf2 binds to Maf proteins to form heterodimers, which then recognize and bind to antioxidant response elements (AREs)-enhancer sequences located in the regulatory regions of Nrf2 target genes that are critical for the recruitment of transcriptionally essential factors^[44]. In this way, the Nrf2-Maf heterodimer activates the expression of target genes and regulates the transcriptional activity of phase II metabolic enzymes, antioxidant enzymes or drug transporters, thus exerting antioxidant damage and counteracting the effects of oxidative stress^[45].

Flavonoids, as antioxidants, can act as antioxidant damages and restore cellular homeostasis by activating the Keap1-Nrf2/ARE signalling pathway and modulating the transcriptional activity of antioxidant enzymes or drug transporters. This signalling pathway plays a key role in cellular resistance to external oxidative stress and is an important defence system against oxidative damage^[46]. As illustrated in *Fig. 3-c*, quercetin and apigenin in the flavonoids of genus Azalea can inhibit the degradation of Nrf2, promote the translocation of Nrf2 into the nucleus, and increase the level of Nrf2 in




Fig 3. Flavonoids of Azalea plants are antioxidant by regulating Keap1-Nrf2 and MAPKS signalling pathways. (a) Under normal conditions, Keap1-Nrf2 signalling pathway, Nrf2 signalling molecules are degraded, (b) Suffering from oxidative stress, Nrf2 signalling molecules are released into the nucleus to bind with Maf proteins to form heterodimers and exert antioxidant effects, (c) Quercetin and apigenin promote Nrf2 translocation into the nucleus and increase the level of Nrf2 in the nucleus, (d) Flavonoids activate Nrf2 through MAPKS and PI3K/AKT signalling pathways, which in turn enhance the antioxidant response

the nucleus as a means of up-regulating the endogenous antioxidant capacity of cells^[47,48].

Antioxidant MAPK_s Signalling Pathway

The mitogen-activated protein kinase (MAPK_s) signalling pathway consists of three major signalling pathways, JNK, ERK and p38, which play an important role in cellular responses to external stimuli (e.g. oxidative stress)^[49]. It has been shown that ROS activate Nrf2 through MAPKs and phosphatidylinositol-3-kinase/protein kinase B (PI3K/AKT) cell signalling pathways in response to cellular damage caused by oxidative stress^[50].

As illustrated in Fig. 3-d, flavonoids activate Nrf2 via MAPKs signalling pathway in conjunction with PI3K/AKT signalling pathway and thus enhance the antioxidant response.

MECHANISM OF ACTION OF ANTI-INFLAMMATORY

The anti-inflammatory properties of flavonoids from the genus *Azalea* are achieved through four main areas: inhibition of the production of inflammatory mediators, modulation of cytokines and receptors, scavenging of oxygen free radicals and cell signalling pathways.

Inhibition of the Production of Inflammatory Mediators

Prostaglandin E2 (PGE2) and NO are the two main pro-inflammatory mediators. Arachidonic acid (AA) produces

PGE2 via the cyclooxygenase pathway, which contributes to inflammation by increasing vascular permeability and vasodilation causing redness, swelling, stiffness and pain^[51]. Cyclooxygenase 2 (COX-2) mainly catalyses the production of large amounts of PGE2, which is involved in the inflammatory response, and inducible NO synthase (iNOS) catalyses the production of large amounts of NO from L-arginine, resulting in cellular damage^[52]. Thus, inhibition of COX-2 and iNOS expression reduces cellular inflammation and alleviates cellular damage.

As illustrated in Fig. 4-a, quercetin and lignans in the flavonoids of *Rhododendron* can inhibit the expression of iNOS and COX-2 at the mRNA and protein levels to reduce the production of NO and PGE2 and exert their anti-inflammatory effects^[52,53].

Regulation of Cytokines and Receptors

In the inflammatory response, proliferating cells such as macrophages, neutrophils, eosinophils and epithelial cells can synthesise and release a variety of cytokines such as interleukin 1 β (IL-1 β), interleukin 6 (IL-6), interleukin 8 (IL-8), and tumour necrosis factor (TNF- α), etc. Flavonoids can be used to inhibit these inflammatory cytokines by inhibiting their production and hindering their association with the Receptor binding, play an anti-inflammatory role^[54].

The flavonoids of the genus *Rhododendron*, including kaempferol, quercetin and lignans, can all have anti-




Fig 4. Azalea flavonoids are anti-inflammatory by inhibiting the production of inflammatory mediators, modulating cytokines and receptors and scavenging oxygen radicals. (a) Quercetin and lignans inhibit the expression of iNOS and COX-2 and reduce NO and PGE2 production, (b) Kaempferol, quercetin and lignans regulate cytokines and exert anti-inflammatory effects, (c) Quercetin and catechin scavenge oxygen free radicals and inhibit the activation of inflammatory mediators by oxygen free radicals

inflammatory effects by modulating cytokines. Kaempferol and quercetin can rapidly scavenge DPPH and ABTS free radicals, and can effectively inhibit the secretion of IL-6, IL-1 β and TNF- α [55]. A dose of quercetin (10-25 mol/L) reduces NO and TNF- α in lipopolysaccharide (LPS)-induced mouse glioma cells [56]. As illustrated in Fig. 4-b, mucuna pruriens has a highly significant inhibitory effect on both TNF- α and IL-6 [57].

Radical Scavenging

Oxygen free radicals are non-specific damage factors widely present in phagocytes of the body and are one of the important pathological mechanisms of inflammation. When the body is stimulated by inflammation, macrophages will produce a large number of oxygen free radicals, disrupting the dynamic balance in the body; at the same time, oxygen free radicals will also activate inflammatory factors, exacerbating the inflammatory response. Therefore, elimination of oxygen free radicals can slow down the inflammatory response [58]. The chemical structure of flavonoids, especially those with a catechol structure, tends to have a large number of phenolic hydroxyl groups, a structural advantage that gives them good free radical scavenging properties [35].

As illustrated in Fig. 4-c, quercetin and catechin in the flavonoids of *Rhododendron* contain o-diphenol hydroxyl structure, which can effectively scavenge oxygen free radicals [59].

Cell Signalling Pathways

NF- κ B Signalling Pathway: NF- κ B is a dimer consisting of five mono-peptide proteins, P50 (also known as NF- κ B1), p52 (also known as NF- κ B2), RelA (also known as p65), RelB and c-Rel [60]. It is also a widely distributed and functional eukaryotic transcription factor. Functional NF- κ B binding sequences are present in promoters and enhancers of genes, so NF- κ B can bind to fixed nucleotide sequences in the promoter regions of many genes to initiate gene transcription, and control a variety of targets such as cytokines, adhesion factors, growth factors, enzymes (COX-2, iNOS), neuropeptides, and so on. It plays an important role in immune response, inflammation and cell growth regulation [61]. Flavonoids inhibit IKK (IK kinase) production and increase I κ B (inhibitory factor) expression by preventing NF- κ B from entering the nucleus for transcription. The anti-inflammatory effects of these compounds are exerted in three ways: by preventing NF- κ B from entering the nucleus for transcription, by inhibiting IKK (IK kinase) production and by increasing I κ B expression [58].

Quercetin inhibits the activation of the NF- κ B signalling pathway and prevents the entry of NF- κ B into the nucleus for transcription, thereby reducing the inflammatory response [62]. As illustrated in Fig. 5-a, lignans can significantly inhibit the phosphorylation of I κ K β and I κ B α in inflammatory cells, reduce the expression of P65 in the nucleus, and then better inhibit the NF- κ B signalling pathway to exert anti-inflammatory activity [57].




Fig 5. Flavonoids of the genus *Azalea* are antioxidants through the signalling pathway. NF- κ B regulates the inflammatory response through both classical and non-classical pathways. The classical pathway is dependent on the activation of the I κ B kinase complex (IKK), which leads to the phosphorylation of the I κ B protein, which in turn triggers ubiquitination and proteasomal degradation of I κ B, and after I κ B degradation, the dimer of p50/RelA is released and enters the nucleus. The non-classical pathway is dependent on NF- κ B-inducible kinase (NIK), activation of which leads to partial degradation of p100 into p52, which forms a dimer with RelB and enters the nucleus. (a) Quercetin and lignans inhibit the NF- κ B signalling pathway and prevent its dimerisation into the nucleus, (b) Flavonoids reduce the activity of NF- κ B dimer and AP-1 in the nucleus of the cell nucleus to achieve cellular anti-inflammation

Transcription Factor Activator Protein 1 (AP-1): AP-1 is an intracellular transcriptional activator that is a heterodimer of c-Fos and c-Jun. Often works with NF- κ B to regulate the expression of inflammatory factors in mediating inflammatory responses [63]. NF- κ B is mainly involved in the regulation of cytokine expression and other inflammatory mediators, whereas AP-1 is involved in the synthesis of innate immune effector molecules and cytokine responses [64]. As illustrated in Fig. 5-b, quercetin Significantly Reduces High Glucose-Induced NF- κ B and AP-1 Activity [65].

APPLICATIONS OF RHODODENDRON FLAVONOIDS

Maintain Blood Pressure Homeostasis

Flavonoids exert anti- MAPKs inflammatory and antioxidant effects by modulating the Keap1-Nrf2/ARE, NF- κ B, and AP-1 signalling pathways, playing a significant role in maintaining cardiovascular homeostasis and tissue protection in animals. Research indicates that flavonoids may assist in maintaining blood pressure homeostasis, mitigate abnormal vascular lipid deposition, and safeguard myocardial structural and functional integrity, thereby demonstrating positive physiological protective effects on the cardiovascular system in animals [3].

Flavonoids exert a blood pressure-stabilising regulatory effect, with quercetin exerting a significant influence on blood pressure by reducing diastolic pressure in hypertensive animal models and systolic pressure in normotensive animal models [66]. Nitric oxide (NO), as the primary vasodilator, plays a crucial role in blood pressure regulation. In rat models, kaempferol administered orally at 10 mg kg⁻¹ day⁻¹ for 6 weeks significantly enhanced aortic eNOS expression, elevated serum NO levels, and reduced systolic blood pressure (approximately ↓18 mmHg) in a dose-dependent manner [67]. In spontaneously hypertensive rat models, the *Rhododendron*-derived flavonoid falarone (at doses of 10–40 mg kg⁻¹ over 8 weeks) reduced systolic blood pressure in a dose-dependent manner by inhibiting angiotensin II-mediated NADPH oxidase activation. This alleviated vascular oxidative stress, restored endothelium-dependent relaxation in the aorta, and reduced intimal hypertrophy [68]. In dogs, intravenous administration of rhoifolin (5 mM/kg) and vitexin significantly reduced mean aortic pressure (by 8%), arterial and pulmonary capillary pressures, and heart rate through vasodilation and negative inotropic effects. Oral administration of *Elsholtzia blanda* total flavonoids (25–100 mg/kg) dose-dependently reduced myocardial infarction area (from 19.42% to 8.87%), decreased serum creatine kinase-MB (CK-MB) and malondialdehyde (MDA) levels, and inhibited lipid peroxidation in a coronary artery occlusion model [69,70].

In horses, administration of 50–75 mg horse chestnut seed extract (containing 2% aescin and flavonoids) every 12 h improved venous tone, reduced transcapillary filtration, and enhanced venous return, with clinical application in chronic venous insufficiency [71].

Lowering blood lipid levels is the optimal approach for preventing and treating atherosclerotic diseases, whilst flavonoids counteract atherosclerosis by inhibiting inflammation in adipose tissue, enhancing cholesterol reverse transport, and reducing blood lipid levels. In a high-fat diet rat model, continuous oral administration of *Rhododendron* methanolic extract (dose: 300–200 mg/kg-d) for 63 days reduced plasma total cholesterol by approximately 28%, triglycerides by approximately 32%, and low-density lipoprotein cholesterol by approximately 35%, while increasing high-density lipoprotein cholesterol by approximately 25%. Its efficacy in reducing the atherosclerosis index surpassed that of lovastatin (10 mg/kg), with no significant toxic reactions observed [72]. In a study by Shubhi Agarwal and colleagues on a high-cholesterol New Zealand rabbit model, continuous oral administration of *Rhododendron* extract reduced serum total cholesterol, triglycerides, and low-density lipoprotein cholesterol levels, increased high-density lipoprotein cholesterol, and decreased the atherosclerosis index [73].

Restriction of cardiac contraction and relaxation constitutes a common clinical manifestation of cardiomyopathy. Calcium ions represent a primary factor in cardiac contraction whilst flavonoid compounds can inhibit Ca²⁺-sensitivity dysregulation induced by troponin I phosphorylation [74,75]. In isolated rat hearts and cardiomyocytes, total *Rhododendron* flavonoids (5–20 µg/mL) inhibited potassium-induced calcium influx and sarcoplasmic reticulum calcium release, thereby reducing intracellular Ca²⁺ levels. This mitigated hypoxia-induced injury and diminished infarct size, revealing a calcium-regulated cardioprotective mechanism [76]. In an autoimmune myocarditis rat model, green tea catechins (400 mg/kg-d administered for 4 consecutive weeks) mitigated left ventricular dysfunction, reduced inflammatory cell infiltration and myocardial fibrosis, and shifted the cytokine profile towards anti-inflammatory characteristics. This demonstrates that tea flavonoids can simultaneously alleviate inflammatory responses and adverse remodelling in experimental myocarditis [77].

Anticancer

Flavonoids, as natural radiosensitising agents and chemotherapeutic sensitising agents, can be integrated into comprehensive cancer management across four levels: primary prevention, synergistic radiotherapy, overcoming drug resistance, and secondary chemoprevention. They demonstrate significant potential particularly in enhancing radiotherapy sensitivity, improving the efficacy of

chemotherapeutic drugs, and reversing tumour multidrug resistance ^[78].

Regarding radiotherapy sensitisation, multiple studies provide clear dose-response evidence. In colon cancer models, 24 h pretreatment of HT-29 and DLD-1 cells with 50 μ M quercetin significantly increased sensitivity to 6 Gy X-rays by downregulating the Notch-1 signalling pathway, reducing colony survival rates by approximately 40% ^[79]. In mouse solid tumour studies, daily intraperitoneal administration of 25 mg/kg apigenin combined with 10 mg/kg cryptotanshinone for 14 consecutive days, alongside 2 Gy local irradiation, achieved a tumour volume suppression rate of 78% in Ehrlich tumours. This effect markedly exceeded the 44% observed in the radiotherapy-alone group, fully demonstrating the advantages of flavonoid compounds in synergising radiotherapy ^[80].

Flavonoids also demonstrate remarkable efficacy in enhancing chemotherapy outcomes. In a prostate cancer xenograft model, daily oral administration of 50 mg/kg quercetin concurrently with weekly docetaxel (5 mg/kg) over four weeks elevated tumour weight inhibition to 72%, substantially surpassing the 38% achieved by docetaxel monotherapy. Its mechanism of action is associated with downregulating P-glycoprotein (P-gp) expression, inhibiting the PI3K/Akt signalling pathway, and modulating androgen receptor signalling ^[81]. In the hormone-refractory PC-3 prostate cancer model, combining 40 mg/kg quercetin with 10 mg/kg paclitaxel (administered every 3 days for 5 consecutive cycles) prolonged tumour doubling time by 1.8-fold through inducing endoplasmic reticulum (ER) stress and reactive oxygen species (ROS) bursts, offering novel insights for enhancing chemotherapy efficacy ^[81]. In addition to synergising with chemotherapeutic agents, certain flavonoid compounds themselves exhibit distinct *in vivo* antitumour activity. For instance, research by Ma et al. ^[82] on the *in vivo* antitumour evaluation of *Rhododendron decorum* flavonoids demonstrated that in a Kunming mouse model transplanted with S180 sarcoma, oral administration of total flavonoids from *Rhododendron grandiflorum* (at doses of 50, 100, 200 mg/kg/d) via oral gavage for 10 consecutive days. The 200 mg/kg group achieved a tumour volume inhibition rate of 54.7% ($P < 0.01$). Mechanistic studies revealed that this dose significantly elevated serum interleukin-2 (IL-2) and TNF- α levels while reducing VEGF levels, suggesting a synergistic anticancer effect through immune enhancement and anti-angiogenesis. No hepatotoxicity, nephrotoxicity, or abnormal body weight changes were observed during the experiment. Consequently, it is proposed that *Rhododendron decorum* flavonoids exert a dose-dependent antitumour effect in animals via a dual immune-antiangiogenic mechanism.

In the field of reversing tumour multidrug resistance (MDR), the efficacy of flavonoid compounds has also been demonstrated. In a colon cancer model resistant to 5-fluorouracil (5-FU), co-treatment with 20 μ M kaempferol and 5 μ M 5-FU for 48 hours produced a significant synergistic inhibitory effect on LS174-Resistant cells (combination index $CI \approx 0.6$), elevating apoptosis rates to 3.7 times that of monotherapy, primarily through inhibiting ABCB1 transporter function and inducing G₂/M phase arrest in the cell cycle ^[81]; In the HL-60/NB4 acute myeloid leukaemia resistance model, 40 μ M kaempferol effectively restored cell sensitivity to doxorubicin by downregulating ABCB1 and ABCC1 transporter expression alongside Akt and BCL2 signalling molecules, reducing the resistance index from 8.2 to 2.1 ^[83].

The aforementioned studies consistently demonstrate that flavonoid compounds such as quercetin, kaempferol, and total flavonoids from *Rhododendron anthopogon*, within the dose range of 20-200 μ M (*in vitro*) or 25-200 mg/kg (*in vivo*), can significantly enhance radiotherapy/chemotherapy efficacy, reverse ABC transporter-mediated resistance, or exert direct antitumour effects through diverse mechanisms. These well-defined dose-response relationships in preclinical studies provide a foundation for incorporating flavonoid compounds into clinical anticancer combination therapies or for their standalone use. Chemotherapy efficacy, reverse ABC transporter-mediated resistance, or exert direct antitumour effects. This clear dose-response relationship in preclinical evidence provides substantial support for the clinical application of flavonoids in combined anticancer therapies or as monotherapy.

Animal Diabetes Mellitus

Flavonoids intervene in type 2 diabetes mellitus (T2DM) and its microvascular and macrovascular complications by lowering blood glucose, improving insulin sensitivity, and preventing complications. Regarding hypoglycaemic effects and insulin resistance improvement, high-dose quercetin (≥ 100 mg/kg/day via gastric lavage in db/db mice for 8 weeks) inhibits protein tyrosine phosphatase 1B (PTP1B), elevates insulin receptor substrate 2 (IRS-2) phosphorylation, and markedly reduces fasting blood glucose. Administered via gastric lavage to db/db mice for 8 weeks) inhibits protein tyrosine phosphatase 1B (PTP1B), elevates insulin receptor substrate 2 phosphorylation levels, significantly reduces fasting blood glucose, and improves oral glucose tolerance. At equivalent doses, kaempferol mitigated inflammatory insulin resistance by blocking IKK β /NF- κ B signalling, thereby downregulating inflammatory mediators such as IL-6 and TNF- α ^[84].

Flavonoids exert effects by promoting insulin secretion and modulating the intestinal-pancreatic axis. In a

C57BL/6 high-fat diet model, 50 $\mu\text{mol/L}$ quercetin or 25 $\mu\text{mol/L}$ kaempferol effectively stimulated L-cell secretion of glucagon-like peptide-1 (GLP-1) in the ileum, inhibiting dipeptidyl peptidase-4 (DPP-4) activity and elevating total GLP-1 levels by 1.8-fold, thereby enhancing glucose-dependent insulin release [85]. In preventing diabetic complications, flavonoids exhibit multi-target protective effects: For retinopathy, quercetin at 50 mg/kg/d (intraperitoneal injection, STZ-induced rats, 12 weeks) suppressed retinal vascular endothelial growth factor (VEGF) and intercellular adhesion molecule-1 (ICAM-1) expression, reducing blood-retinal barrier leakage; For diabetic nephropathy, kaempferol at 75 mg/kg/day (oral administration, db/db mice for 16 weeks) reduced urinary albumin excretion and fibronectin deposition, a mechanism associated with inhibition of the TGF- β 1/Smad3 signalling pathway; For cardiomyopathy and osteoporosis, quercetin alleviates myocardial oxidative stress by activating the Nrf2/HO-1 pathway and protects the bone matrix by inhibiting receptor for advanced glycation end-products (RAGE) expression [86].

CONCLUSION

In this paper, we review the past research progress on the mechanism and clinical application of antioxidant and anti-inflammatory activities of flavonoids from *Rhododendron* plants. We have identified 17 flavonoids in *Rhododendron* plants with significant antioxidant and anti-inflammatory activities. These compounds act through a variety of mechanisms, including direct scavenging of free radicals, inhibition of oxidative enzymes (e.g. NADPH oxidase and xanthine oxidase), chelation of metal ions, provision of hydrogen ions and enhancement of the antioxidant system to stabilise or reduce reactive oxygen species (ROS) and scavenging of free radicals, as well as modulation of the endogenous antioxidant capacity by the Keap1-Nrf2/ARE and MAPK_s signalling pathways. The antioxidant is achieved by means of the Keap1-Nrf2/ARE and MAPK_s signalling pathways. Its anti-inflammatory effects are achieved by inhibiting the production of inflammatory mediators, modulating cytokines and receptors, scavenging oxygen free radicals, and affecting cell signalling pathways such as NF- κ B and AP-1. At present, it has become a prevailing trend to study the antioxidant and anti-inflammatory properties of flavonoids at home and abroad. *Rhododendron* is found all over the world and is rich in flavonoids, so isolating the specific composition and structure of flavonoids from *Rhododendron* will help to design and develop more effective natural antioxidant protective agents and animal health supplements [87]. An in-depth elucidation of its antioxidant and anti-inflammatory mechanisms holds promise for advancing the development of novel natural

antioxidants for protecting animal organisms against oxidative damage and maintaining health homeostasis [88].

In terms of clinical applications, flavonoids have been shown to have potential therapeutic effects on cardiovascular diseases such as hypertension, atherosclerosis, cardiomyopathy and myocarditis [3]. It also shows great therapeutic potential in the fields of diabetes and anticancer [78,86]. A deeper understanding of their antioxidant and anti-inflammatory mechanisms will advance research into the application of flavonoids in animal health management, thereby fully harnessing their physiological protective and supportive efficacy.

Flavonoids and terpenoids can act synergistically to enhance each other's pharmacological activities, especially in the fight against cancer. Terpenoids modulate caspase-3 activity and flavonoid circuit affects enzyme activity [89]. Flavonoids and terpenoids collectively influence the regulation of ATP-binding cassette (ABC) transporter protein efflux function, and semi-synthetic nitrogen-containing flavonoids and terpenoids derivatives possess potential as multidrug resistance (MDR) reversal agents designed to be effective in cancer [90]. Azalea plants are rich in flavonoids and terpenoids, the study of Azalea plants can not only study flavonoids and terpenoids, but also study the synergistic effect of the two, the development of new anti-cancer drugs.

Although some progress has been made on the antioxidant and anti-inflammatory activities of flavonoids in the genus *Rhododendron*, there are still some limitations, such as the absence of systematic studies on their mechanistic mechanisms, small sample sizes, and geographical bias. Future studies need to consider a wider range of samples and different geographical regions to validate the existing findings and further systematically explore the pharmacological mechanisms of action of flavonoids [10]. Moreover, in-depth research into flavonoids holds promise for providing further scientific rationale for developing novel approaches to maintaining and safeguarding animal health, particularly demonstrating significant research and application value in antioxidant and anti-inflammatory physiological protection.

In conclusion, significant progress has been made in research concerning the antioxidant and anti-inflammatory physiological activities of flavonoids from *Rhododendron* species. This has laid a solid foundation for subsequent development of natural physiological protective agents and their application in animal health management. Future studies should continue to explore the physiological protective and supportive potential of these compounds, overcome existing research limitations, and further expand their application scenarios in safeguarding animal organisms and maintaining health.

DECLARATIONS

Availability of Data and Materials: The data and materials used to support the findings of this study are available from the first author (Z.Z.) and the primary corresponding author (Z.C.) upon reasonable request. All relevant data have been properly curated and can be provided to facilitate reproducibility of the research results.

Acknowledgements: The authors sincerely thank Xizang Agricultural and Animal Husbandry University, School of Animal Science and Tibetan Key Laboratory of Clinical Veterinary Medicine for equipment and financial support. We also appreciate the reviewers and editors for their valuable comments that improved this manuscript.

Funding Information: This study was supported by the grants from Science and Technology Programme of Linzhi City, Tibet Autonomous Region (XDHZ-2025-07)

Conflict of Interest: The authors declare no conflicts of interest.

Declaration on Generative Artificial Intelligence (AI): Generative artificial intelligence was only used for language polishing in this study; it was not involved in any other aspects of the research. All authors have reviewed the final content and take full responsibility for its authenticity and scientificity.

Author Contributions: All authors made substantial contributions to this study. Ziqin Zhang was responsible for conceptualization, methodology and original draft writing; Sheng Kang and Hailong Dong engaged in data curation and formal analysis; Meixue Ni and Xiaoqing Guo participated in investigation, resources collection and validation. As corresponding authors, Qingxia Wu was in charge of supervision, funding acquisition and manuscript review & editing; Zhenyu Chang, the primary corresponding author, undertook supervision, conceptualization, methodology and manuscript review & editing.

REFERENCES

- Liu XJ, Su HG, Peng XR, Bi HC, Qiu MH: An updated review of the genus *Rhododendron* since 2010: Traditional uses, phytochemistry, and pharmacology. *Phytochemistry*, 217:113899, 2024. DOI: 10.1016/j.phytochem.2023.113899
- Cheng J, Li M, Yuan T, Huang H, Yang G, Bai X: A dataset on wild *Rhododendron* and geographical distribution information in China. *Biodivers Sci*, 29 (9): 1175-1180, 2021. DOI: 10.17520/biods.2021152
- Liu Y, Luo J, Peng L, Zhang Q, Rong X, Luo Y, Li J: Flavonoids: Potential therapeutic agents for cardiovascular disease. *Heliyon*, 10 (12):E32563, 2024. DOI: 10.1016/j.heliyon.2024.e32563
- Pooja G, Shweta S, Patel P: Oxidative stress and free radicals in disease pathogenesis: A review. *Discov Med*, 2:104, 2025. DOI: 10.1007/s44337-025-00303-y
- Sarkar C, Chaudhary P, Jamaddar S, Janmeda P, Mondal M, Mubarak MS, Islam MT: Redox activity of flavonoids: Impact on human health, therapeutics, and chemical safety. *Chem Res Toxicol*, 35 (2): 140-162, 2022. DOI: 10.1021/acs.chemrestox.1c00348
- Pathology Teaching and Research Group, Lanzhou Medical College, Lanzhou, China: Pharmacological study and toxicity observation of loquatoin. *Chin Med J*, 54 (5): 279-283, 1974.
- Yang HX, Xu W, Huang ZH: Chemical constituents of *Anoectochilus roxburghii* based on UPLC-Q-TOF-MS/MS. *Chin J Mod Appl Pharm*, 42 (6): 948-957, 2025. DOI: 10.13748/j.cnki.issn1007-7693.20232655
- Tan C, Chen X, Liu X, Hui JM, Huo XY, Gao SJ, Wang YH, Wang JR, Zhang XM, Yang Y: Progress in the study of *Rhododendron*. *China Wild Plant Resour*, 40 (9): 29-36, 2021.
- Verma K, Kumar I: A complete review on *Rhododendron arboreum*: Phytochemistry, pharmacological activities and future prospects. *Int J Pharm Pharm Res*, 19 (3): 161-176, 2020.
- Wang L, Zeng X, Yao HZ, Yuan T: Studies on flavonoids from *Rhododendron tigrinum* and their antioxidant and anti-inflammatory activities. *Nat Prod Res Dev*, 36 (8): 1350-1356, 2024.
- Hu M, Liu YL, Xiao PG: Chemical composition of *Rhododendron angustifolium*. *J Integr Plant Biol*, 32 (10): 777-782, 1990.
- Li ZL, Xue DY, Chen YZ, Shi ZX, Zhang JX: Studies on the chemical composition of the volatile oil of *Rhododendron schoenanthus*. *Nat Prod Res Dev*, 2, 11-16, 1991. DOI: 10.16333/j.1001-6880.1991.02.003
- Zhao X, Che WS: Progress of chemical composition and extraction method of Manzan Hong. *J Anhui Med Univ*, 48 (5): 580-582, 2013. DOI: 10.19405/j.cnki.issn1000-1492.2013.05.038
- Duan WT, Cheng Q, Bi XX, Wang Y, Cheng B, Luan J, Zhang GH, He JB: Studies on the chemical composition of *Rhododendron* on cloud and its no production inhibiting activity. *Nat Prod Res Dev*, 36 (1): 63-71, 2023. DOI: 10.16333/j.1001-6880.2024.1.007
- Zhou XL, Qin CH, Mei Y, Huang S: Studies on the chemical composition of *Rhododendron barbadensis* leaves. *Chin Herb Med*, 41 (2): 206-208, 2010.
- Zhou SY, Li RT: Progress in the study of chemical composition and biological activity of *Rhododendron* plants. *Yunnan J Tradit Chin Med*, 30 (5): 51-53, 2008. DOI: 10.16254/j.cnki.53-1120/r.2008.05.036
- Xiao YC, Hu FZ, Du YZ, Wei LX: Determination of seven flavonoid constituents in different parts of Tibetan medicine *Rhododendron simsii* by HPLC method. *J Pharm Anal*, 38 (8): 1325-1330, 2018. DOI: 10.16155/j.0254-1793.2018.08.06
- Xia X, Gong R, Zhang C: Integrative analysis of transcriptome and metabolome reveals flavonoid biosynthesis regulation in *Rhododendron pulchrum* petals. *BMC Plant Biol*, 22 (1):401, 2022. DOI: 10.1186/s12870-022-03762-y
- Zhang M, Pan DR: Progress in the study of flavonoids in *Rhododendron* plants. *Sci Across Taiwan Straits*, 28 (5): 3-4+14, 2012.
- Oliveira THB, Gusmao NB, Silva LAO, Coelho LCBB: Free radicals and actinobacteria as a misexplored goldmine of antioxidant compounds. *An Acad Bras Cienc*, 93 (Suppl.-3): E20201925, 2021. DOI: 10.1590/0001-3765202120201925
- Shen N, Wang T, Gan Q, Liu S, Wang L, Jin B: Plant flavonoids: Classification, distribution, biosynthesis, and antioxidant activity. *Food Chem*, 383:132531, 2022. DOI: 10.1016/j.foodchem.2022.132531
- Valko M, Leibfritz D, Moncol J, Cronin MTD, Mazur M, Telser J: Free radicals and antioxidants in normal physiological functions and human disease. *Int J Biochem Cell Biol*, 39 (1): 44-84, 2007. DOI: 10.1016/j.biocel.2006.07.001
- Yang X, Wang H, Lu K, Ma X, Tan Z, Long B, Chen X, Li C, Zhai T, Li Y, Qu K, Xia Y, Zhang Y, Li X, Chen S, Dong H, Zeng L, Zhang YH: Reactive aldehyde chemistry explains the missing source of hydroxyl radicals. *Nat Commun*, 15 (1):1648, 2024. DOI: 10.1038/s41467-024-45885-w
- Guo CM, Liu SQ, Guo YM, Yu HS, Sun MZ: Scavenging function of hydroxyl radicals (Ho·) by flavonoids and their conformational relationships. *Nat Prod Res Dev*, 24 (12): 1724-1728, 2012. DOI: 10.16333/j.1001-6880.2012.12.007
- Lu L, Luan Y, Wang H, Gao Y, Wu S, Zhao X: Flavonoid as a potent antioxidant: quantitative structure-activity relationship analysis, mechanism study, and molecular design by synergizing molecular simulation and machine learning. *J Phys Chem A*, 128 (30): 6216-6228, 2024. DOI: 10.1021/acs.jpca.4c03241
- Garcia JG, Ansorena E, Izal I, Zalba G, De Miguel C, Milagro FI: Structure, regulation, and physiological functions of NADPH oxidase 5 (NOX5). *J Physiol Biochem*, 79 (2): 383-395, 2023. DOI: 10.1007/s13105-023-00955-3
- Ullah Z, Yue P, Mao G, Zhang M, Liu P, Wu X, Zhao T, Yang L: A Comprehensive review on recent xanthine oxidase inhibitors of

- dietary based bioactive substances for the treatment of hyperuricemia and gout: Molecular mechanisms and perspective. *Int J Biol Macromol*, 278:134832, 2024. DOI: 10.1016/j.ijbiomac.2024.134832
28. **Balázs O, Dombi Á, Zsidó BZ, Hetényi C, Valentová K, Vida RG, Poór M:** Inhibition of xanthine oxidase-catalyzed xanthine and 6-mercaptopurine oxidation by luteolin, naringenin, myricetin, ampelopsin and their conjugated metabolites. *Biomed Pharmacother*, 167:115548, 2023. DOI: 10.1016/j.biopha.2023.115548
 29. **Yan J, Zhang G, Hu Y, Ma Y:** Effect of luteolin on xanthine oxidase: Inhibition kinetics and interaction mechanism merging with docking simulation. *Food Chem*, 141 (4): 3766-3773, 2013. DOI: 10.1016/j.foodchem.2013.06.092
 30. **Liu Y, Liang S, Shi D, Zhang Y, Bai C, Ye RD:** A predicted structure of NADPH oxidase 1 identifies key components of ROS generation and strategies for inhibition. *Plos One*, 18 (5):E0285206, 2023. DOI: 10.1371/journal.pone.0285206
 31. **Yu Y, Xiong Y, Tong S, Li Y, Cai R, Zhang X, Gao F:** Inhibitory activity of quercetin, rutin, and hyperin against xanthine oxidase: Kinetics, fluorescence, and molecular docking. *Curr Pharm Biotechnol*, 25, 1-12, 2024. DOI: 10.2174/0113892010297269240427055003
 32. **Porfirio DA, Ferreira RDQ, Malagutti AR, Valle EMA:** Electrochemical study of the increased antioxidant capacity of flavonoids through complexation with iron(II) ions. *Electrochim Acta*, 141, 33-38, 2014. DOI: 10.1016/j.electacta.2014.07.046
 33. **Zhu L, Ma LL, Li XS, Lu ZK, Lv WH, Le XY:** Preparation and *in vitro* antioxidant activity of two copper(II) complexes of flavonoids. *Food Ind Sci Technol*, 40 (4): 51-55+61, 2018. DOI: 10.13386/j.issn1002-0306.2019.04.008
 34. **Mira L, Fernandez MT, Santos M, Rocha R, Florencio MH, Jennings KR:** Interactions of flavonoids with iron and copper ions: A mechanism for their antioxidant activity. *Free Radic Res*, 36 (11): 1199-1208, 2002. DOI: 10.1080/1071576021000016463
 35. **Zheng YZ, Deng G, Zhang YC:** Multiple free radical scavenging reactions of flavonoids. *Dyes Pigments*, 198:109877, 2022. DOI: 10.1016/j.dyepig.2021.109877
 36. **Zhou W, Peng C, Wang D, Li J, Tu Z, Zhang L:** Interaction mechanism between ova and flavonoids with different hydroxyl groups on B-ring and effect on antioxidant activity. *Foods*, 11 (9): 1302, 2022. DOI: 10.3390/foods11091302
 37. **Jomova K, Alomar SY, Alwasel SH, Nepovimova E, Kuca K, Valko M:** Several lines of antioxidant defense against oxidative stress: Antioxidant enzymes, nanomaterials with multiple enzyme-mimicking activities, and low-molecular-weight antioxidants. *Arch Toxicol*, 98 (5): 1323-1367, 2024. DOI: 10.1007/s00204-024-03696-4
 38. **Rudenko NN, Vetoshkina DV, Marenkov TV, Borisova-Mubarakshina MM:** Antioxidants of non-enzymatic nature: Their function in higher plant cells and the ways of boosting their biosynthesis. *Antioxidants*, 12 (11):2014, 2023. DOI: 10.3390/antiox12112014
 39. **Gao Y, Dong X, Wang R, Hao F, Zhang H, Zhang Y, Lin G:** Exogenous calcium alleviates oxidative stress caused by salt stress in peanut seedling roots by regulating the antioxidant enzyme system and flavonoid biosynthesis. *Antioxidants*, 13 (2):233, 2024. DOI: 10.3390/antiox13020233
 40. **Sheng YI, Liu H, Yu J:** Influence of structural and non-structural factors on the role of flavonoids in antioxidant activity. *Chin Pharm*, 24 (7): 660-663, 2013.
 41. **Morgenstern C, Lastres-Becker I, Demirdöğen BC, Costa VM, Daiber A, Foresti R, Motterlini R, Kalyoncu S, Ariozi BI, Genc S, Jakubowska M, Trougakos IP, Piechota-Polanczyk A, Mickael M, Santos M, Kensler TW, Cuadrado A, Coppole IM:** Biomarkers of NRF2 signalling: Current status and future challenges. *Redox Biol*, 72:103134, 2024. DOI: 10.1016/j.redox.2024.103134
 42. **Horie Y, Suzuki T, Inoue J, Iso T, Wells G, Moore TW, Mizushima T, Dinkova-Kostova AT, Kasai T, Kamei T, Koshiha S, Yamamoto M:** Molecular basis for the disruption of keap1-NRF2 interaction via Hinge & Latch mechanism. *Commun Biol*, 4 (1):576, 2021. DOI: 10.1038/s42003-021-02100-6
 43. **Dewanjee S, Bhattacharya H, Bhattacharyya C, Chakraborty P, Fleishman J, Alexiou A, Papadakis M, Jha SK:** Nrf2/keap1/ARE regulation by plant secondary metabolites: A new horizon in brain tumor management. *Cell Commun Signal*, 22 (1):497, 2024. DOI: 10.1186/s12964-024-01878-2
 44. **Ma K, Miao L, Li B, Yu W, Liu F, Liu K, Li Y, Huang C, Yang Z:** Mechanism of action of Nrf2 and its related natural regulators in rheumatoid arthritis. *J Orthop Surg Res*, 19 (1):759, 2024. DOI: 10.1186/s13018-024-05221-w
 45. **Katsoka F, Motohashi H, Engel JD, Yamamoto M:** Nrf2 transcriptionally activates the mafg gene through an antioxidant response element. *J Biol Chem*, 280 (6): 4483-4490, 2005. DOI: 10.1074/jbc.m411451200
 46. **Speisky H, Shahidi F, Costa De Camargo A, Fuentes J:** Revisiting the oxidation of flavonoids: Loss, conservation or enhancement of their antioxidant properties. *Antioxidants*, 11 (1):133, 2022. DOI: 10.3390/antiox11010133
 47. **Dai C, Zhang Q, Shen L, Sharma G, Jiang H, Wang Z, Shen J:** Quercetin attenuates quinocetone-induced cell apoptosis *in vitro* by activating the P38/Nrf2/HO-1 pathway and inhibiting the ros/mitochondrial apoptotic pathway. *Antioxidants*, 11 (8):1498, 2022. DOI: 10.3390/antiox11081498
 48. **Cheng YJ, Quan ZJ, Zheng N, Xie YQ, Chen J:** Apigenin alleviates high glucose-induced renal cell damage and oxidative stress by activating the keap1-Nrf2-ARE signalling pathway. *Chin Pharmacist*, 27 (6): 975-983, 2024.
 49. **Liu TT, Zhang SP, Tang XY, Pan RY:** Advances in MAPK signal transduction pathway and nerve injury research. *Chin Public Health*, 32 (2): 248-254, 2015.
 50. **Cheng C, Yin YY, Wu XF, Wu RR, Xu YY:** Progress in activation of Nrf2 by reactive oxygen species through MAPKs and PI3k/AKT pathways. *Chin Public Health*, 32 (6): 870-873, 2016.
 51. **Kawahara K, Hohjoh H, Inazumi T, Tsuchiya S, Sugimoto Y:** Prostaglandin E2-induced inflammation: Relevance of prostaglandin E receptors. *Biochim Biophys Acta Mol Cell Biol Lipids*, 1851 (4): 414-421, 2015. DOI: 10.1016/j.bbali.2014.07.008
 52. **Jin K, Qian C, Lin J, Liu B:** Cyclooxygenase-2-prostaglandin E2 pathway: A key player in tumor-associated immune cells. *Front Oncol*, 13:1099811, 2023. DOI: 10.3389/fonc.2023.1099811
 53. **Hussain MS, Gupta G, Goyal A, Thapa R, Almalki WH, Kazmi I, Alzarea SI, Fuloria S, Meenakshi DU, Jakhmova V, Pandey M, Singh SK, Dua K:** From nature to therapy: luteolin's potential as an immune system modulator in inflammatory disorders. *J Biochem Mol Toxicol*, 37 (11):E23482, 2023. DOI: 10.1002/jbt.23482
 54. **Al-Khayri JM, Sahana GR, Nagella P, Joseph BV, Alessa FM, Al-Msalleem MQ:** Flavonoids as potential anti-inflammatory molecules: A review. *Molecules*, 27 (9):2901, 2022. DOI: 10.3390/molecules27092901
 55. **Lee KJ, Oh YC, Cho WK, Ma JY:** Antioxidant and anti-inflammatory activity determination of one hundred kinds of pure chemical compounds using offline and online screening HPLC assay. *Evid Based Complement Alternat Med*, 2015:65457, 2015. DOI: 10.1155/2015/165457
 56. **Rather MA, Khan A, Alshahrani S, Rashid H, Qadri M, Rashid S, Alsaffar RM, Kamal MA, Rehman MU:** Inflammation and Alzheimer's disease: Mechanisms and therapeutic implications by natural products. *Mediators Inflamm*, 2021:982954, 2021. DOI: 10.1155/2021/982954
 57. **Wang W, He P, Jiang XM:** Anti-inflammatory and antioxidant effects of lignans and their flavonoid glycosides. *Food Sci*, 41 (17): 208-215, 2020.
 58. **Zhang P, Yang J, Gao X, Ling, Guo YR, Tian CL, Liu MC:** Progress in the study of anti-inflammatory effects of flavonoids and their mechanism of action. *Livest Vet Med*, 50 (2): 134-137, 2018.
 59. **Zhao XW, Liu PY, Liu D, Sun SS, Li Z, Yu KX, Zhang ML, Shi QW:** Advances in the study of conformational relationships of flavonoids. *Chin Herb Med*, 46 (21): 3264-3271, 2015.

60. **Msweli S, Pakala SB, Syed K:** NF- κ B Transcription factors: Their distribution, family expansion, structural conservation, and evolution in animals. *Int J Mol Sci*, 25 (18):9793, 2024. DOI: 10.3390/ijms25189793
61. **Geng YF, Du HB, Liu LL, Sun MY, Jiu JY, Zhang JB, Zhao CL, Tian CY:** Advances in the post-translational modification of NF- κ B family member *rela* and its physiopathological role. *Life Sci*, 32 (5): 431-438, 2020. DOI: 10.13376/j.cbcls/2020054
62. **Li L, Jiang W, Yu B, Liang H, Mao S, Hu X, Feng Y, Xu J, Chu L:** Quercetin improves cerebral ischemia/reperfusion injury by promoting microglia/macrophages M2 polarization via regulating PI3k/AKT/NF- κ B signaling pathway. *Biomed Pharmacother*, 168:115653, 2023. DOI: 10.1016/j.biopha.2023.115653
63. **Han CC, Wan FS:** Advances in AP-1 research. *Chin J Cell Biol*, 39 (10): 1357-1362, 2017.
64. **Fujioka S, Niu J, Schmidt C, Sclabas GM, Peng B, Uwagawa T, Li Z, Evans DB, Abbruzzese JL, Chiao PJ:** NF- κ B and AP-1 connection: Mechanism of NF- κ B-dependent regulation of AP-1 activity. *Mol Cell Biol*, 24 (2): 780-794, 2004. DOI: 10.1128/mcb.24.17.7806-7819.2004
65. **Qiao Y, He H, Jonsson P, Sinha I, Zhao C, Dahlman-Wright K:** AP-1 is a key regulator of proinflammatory cytokine TNF α -mediated triple-negative breast cancer progression. *J Biol Chem*, 291 (35): 18309-18320, 2016. DOI: 10.1074/jbc.a115.702571
66. **Cao Y, Xie L, Liu K, Liang Y, Dai X, Wang X, Lu J, Zhang X, Li X, Zhang Y:** The antihypertensive potential of flavonoids from Chinese herbal medicine: A review. *Pharmacol Res*, 174:105919, 2021. DOI: 10.1016/j.phrs.2021.105919
67. **Puzserova A, Bernatova I:** Blood pressure regulation in stress: Focus on nitric oxide-dependent mechanisms. *Physiol Res*, 65 (Suppl.-3): S309-S342, 2016. DOI: 10.33549/physiolres.933442
68. **Qin Xj:** Vasorelaxant effect of farrerol on rat aorta and its protection against aortic injury in spontaneously hypertensive rats. *Msc Thesis*, Shanxi Medical University, 2015.
69. **Refaat J, Desoukey SY, Ramadan MA, Kamel MS:** Rhoifolin: A review of sources and biological activities. *Int J Pharmacogn*, 2, 102-109, 2015. DOI: 10.13040/ijpsr.0975-8232.ijp.2(3).102-09
70. **Guo X, Wang Y, Zhu Z, Li L:** The role of plant extracts in enhancing nutrition and health for dogs and cats: Safety, benefits, and applications. *Vet Sci*, 11:426, 2024. DOI: 10.3390/vetsci11090426
71. **Wakshlag JJ:** Flavonoids: Not just for cancer anymore. *Compend Contin Educ Vet*, 33, E1-4, 2011.
72. **Allyn OQ, Kusumawati E, Nugroho RA:** Antimicrobial activity of terminalia catappa brown leaf extracts against *Staphylococcus aureus* ATCC 25923 and *Pseudomonas aeruginosa* ATCC 27853. *F1000res*, 7:1406, 2018. DOI: 10.12688/f1000research.15998.1
73. **Agarwal S, Rajput A:** *Rhododendron arboreum* phytochemistry, medicinal applications and pharmacology: A review. *Int J Pharmacogn Phytochem Res*, 9 (3): 1-10, 2022.
74. **Zhang Y, Chen B, Wang M, Liu H, Chen M, Zhu J, Zhang Y, Wang X, Wu Y, Liu D, Cui G, Kitakaze M, Kim JK, Wang Y, Luo T:** A novel function of claudin-5 in maintaining the structural integrity of the heart and its implications in cardiac pathology. *Biochim Biophys Acta Mol Basis Dis*, 1870 (6):167274, 2024. DOI: 10.1016/j.bbdis.2024.167274
75. **Papadaki M, Vikhorov PG, Marston SB, Messer AE:** Uncoupling of myofilament Ca²⁺ sensitivity from troponin I phosphorylation by mutations can be reversed by epigallocatechin-3-gallate. *Cardiovasc Res*, 108 (1): 99-110, 2015. DOI: 10.1093/cvr/cvv181
76. **Zhang BN, Hou YL, Liu BJ, Liu QM, Qiao GF:** The *Rhododendron dauricum* L. flavonoids exert vasodilation and myocardial preservation. *Iran J Pharm Res*, 9 (3): 303-311, 2010.
77. **Suzuki J, Ogawa M, Futamatsu H, Kosuge H, Sagesaka YM, Isohe M:** Tea catechins improve left ventricular dysfunction, suppress myocardial inflammation and fibrosis, and alter cytokine expression in rat autoimmune myocarditis. *Eur J Heart Fail*, 9 (2): 152-159, 2007. DOI: 10.1016/j.ejheart.2006.05.007
78. **Liskova A, Samec M, Koklesova L, Brockmueller A, Zhai K, Abdellatif B, Siddeeqi M, Biringer K, Kudela E, Pec M, Gadanec LK, Šudomová M, Hassan STS, Zulli A, Shakibaei M, Giordano FA, Büsselberg D, Golubnitschaja O, Kubatka P:** Flavonoids as an effective sensitizer for anti-cancer therapy: Insights into multi-faceted mechanisms and applicability towards individualized patient profiles. *Epma J*, 12 (2): 155-176, 2021. DOI: 10.1007/s13167-021-00242-5
79. **Li Y, Wang Z, Jin J, Zhu SX, He GQ, Li SH, Wang J, Cai Y:** Quercetin pretreatment enhances the radiosensitivity of colon cancer cells by targeting NOTCH-1 pathway. *Biochem Biophys Res Commun*, 523 (4): 947-953, 2020. DOI: 10.1016/j.bbrc.2020.01.048
80. **Medhat AM, Azab KS, Said MM, El Fatih NM, El Bakary NM:** Antitumor and radiosensitizing synergistic effects of apigenin and cryptotanshinone against solid Ehrlich carcinoma in female mice. *Tumor Biol*, 39 (10):101042831772848, 2017. DOI: 10.1177/1010428317728480
81. **Lu X, Yang F, Chen D, Zhao Q, Chen D, Ping H, Xing N:** Quercetin reverses docetaxel resistance in prostate cancer via androgen receptor and PI3k/AKT signaling pathways. *Int J Biol Sci*, 16 (7): 1121-1134, 2020. DOI: 10.7150/ijbs.41686
82. **Ma YZ, Cheng Q, Fan ZB, Tang XQ, Wang Y, Chen B, Luan J, Zhang GH, He JB:** Chemical constituents of *Rhododendron decorum* and their antitumor cytotoxic activities. *J Yunnan Univ (Nat Sci)*, 46 (3): 555-560, 2024. DOI: 10.7540/j.ynu.20230368
83. **Riahi-Chebbi I, Souid S, Othman H, Haoues M, Karoui H, Morel A, Srairi-Abid N, Essafi M, Essafi-Benkhadir K:** The phenolic compound kaempferol overcomes 5-fluorouracil resistance in human resistant LS174 colon cancer cells. *Sci Rep* 9 (1):195, 2019. DOI: 10.1038/s41598-018-36808-z
84. **Tiganis T:** PTP1B and TCPTP-nonredundant phosphatases in insulin signaling and glucose homeostasis. *Febs J*, 280 (2): 445-458, 2013. DOI: 10.1111/j.1742-4658.2012.08563.x
85. **Anghel SA, Badea RA, Chiritoiu G, Patriche DS, Alexandru PR, Pena F:** Novel luciferase-based glucagon-like peptide 1 reporter assay reveals naturally occurring secretagogues. *Br J Pharmacol*, 179 (19): 4738-4753, 2022. DOI: 10.1111/bph.15896
86. **Wen D, Li M:** The emerging role of flavonoids in the treatment of type 2 diabetes mellitus: Regulating the enteroendocrine system. *Explor Res Hypothesis Med*, 10 (1): 56-68, 2025. DOI: 10.14218/erhm.2024.00055
87. **Ren H, Zheng SJ, Zhang XL, Lu KK, Wu SR, Ming J:** Progress of flavonoids conformational relationships based on different antioxidant mechanisms. *Food Ind Sci Technol*, 37 (2): 384-388, 2016. DOI: 10.13386/j.issn1002-0306.2016.02.070
88. **Hussain A, Azam S, Maqsood R, Anwar R, Akash MSH, Hussain H, Wang D, Imran M, Kotwicz-Mojzycz K, Khan S, Hussain S, Ayub MA:** Chemistry, biosynthesis, and theranostics of antioxidant flavonoids and polyphenolics of genus *Rhododendron*: An overview. *Naunyn-Schmiedeberg's Arch Pharmacol*, 397 (5): 1479-1505, 2024. DOI: 10.1007/s00210-024-03428-6
89. **Masood N, Dubey V, Luqman S:** Activation of caspase-3 by terpenoids and flavonoids in different types of cancer cells. *Curr Top Med Chem*, 20 (21): 1876-1887, 2020. DOI: 10.2174/1568026620666200710101859
90. **Goncalves BME, Cardoso DSP, Ferreira MJU:** Overcoming multidrug resistance: flavonoid and terpenoid nitrogen-containing derivatives as ABC transporter modulators. *Molecules*, 25 (15):3364, 2020. DOI: 10.3390/molecules25153364

