

REVIEW ARTICLE

Fermented Chinese Herbal Medicines Improve Intestinal Health in Livestock and Poultry via Microbial Transformation of Bioactive Compounds and Anti-Inflammatory Mechanisms

Yisheng ZHANG¹ , Zhengshun WANG¹ , DAWAQIONGDA² , Hongming LIU¹ , Yefen XU¹ , Zeyan HUANG¹ , Qinghui KONG^{1(*)} 

¹ Xizang Agricultural and Animal Husbandry University, College of Animal Science, Nyingchi City, Xizang Autonomous Region, CHINA

² Zhongba County Agriculture, Animal Husbandry Integrated Service Center, Zhongba, Xizang Autonomous Region, CHINA



(*) Corresponding author:

Yu Zhang

Phone: +86-13709767423

E-mail: kqh610@sina.com

How to cite this article?

Zhang Y, Wang Z, Dawaqiongda, Liu H, Xu Y, Huang Z, Kong Q: Fermented Chinese Herbal Medicines Improve Intestinal Health in Livestock and Poultry via Microbial Transformation of Bioactive Compounds and Anti-Inflammatory Mechanisms. *Kafkas Univ Vet Fak Derg*, 2026 (Article in Press). DOI: 10.9775/kvfd.2026.37510

Article ID: KVFD-2026-37510

Received: 07.07.2026

Accepted: 06.10.2026

Published Online: 08.10.2026

Abstract

Intestinal inflammation, underpinned by gut dysbiosis, epithelial barrier dysfunction and dysregulated TLR4/NF-κB-mediated innate immunity, represents a major constraint on livestock production efficiency and animal welfare. Traditional Chinese herbal medicines (TCHMs) possess multi-component, multi-target anti-inflammatory profiles and favourable safety margins; however, their therapeutic efficacy is limited by poor oral bioavailability, extensive first-pass metabolism and substantial pharmacokinetic variability. Microbial fermentation can remodel herbal constituents through polysaccharide depolymerisation, flavonoid glycoside hydrolysis, phenolic acid liberation and alkaloid/tannin modification, potentially yielding metabolites with improved bioaccessibility and intestinal anti-inflammatory activity. This narrative review synthesises current evidence on fermentation-mediated phytochemical transformation of Chinese medicinal herbs and the intestinal health effects of the resulting preparations in production animals. Fermentation microorganisms, metabolic profiles and condition-dependent phytochemical transformations are critically examined and integrated with tissue-level protective mechanisms encompassing gut microbiota remodelling, short-chain fatty acid metabolism, epithelial tight-junction reinforcement, suppression of aberrant TLR4/MyD88/NF-κB signalling and pathological NLRP3 inflammasome activation, as well as Nrf2-dependent antioxidant defence. Evidence obtained directly from fermented herbal preparations in production animals is explicitly delineated from that extrapolated from general probiotic or short-chain fatty acid (SCFA) literature. Importantly, much of the mechanistic evidence remains preclinical, and few studies have validated efficacy, optimal dosing or safety under commercial production conditions. Finally, a "strain-ingredient-metabolite-microbiota-host" mechanistic framework is proposed to guide the standardisation of fermentation protocols, quality control and the veterinary application of fermented TCHMs.

Keywords: Antioxidant defence, Chinese medicinal herbs, Gut microbiota, Microbial fermentation, Short-chain fatty acid

INTRODUCTION

Intestinal inflammation encompasses a spectrum of disorders hallmarked by mucosal barrier impairment, gut dysbiosis, and dysregulated innate immunity. In production animals, post-weaning diarrhoea (PWD) in piglets, frequently triggered by enterotoxigenic *Escherichia coli* (ETEC) infection and weaning stress, represents a leading cause of morbidity, growth retardation and economic loss in the swine industry^[1,2]. Pathologically, ETEC-derived lipopolysaccharides and enterotoxins activate TLR4/MyD88/NF-κB signalling, driving

TNF-α, IL-1β and IL-6 production whilst concurrently disrupting tight junction localisation and increasing epithelial permeability^[3,4]. These convergent pathologies illustrate that intestinal inflammation emerges not from isolated cytokine dysregulation, but from the integrated collapse of microbiota homeostasis, barrier integrity and inflammatory control.

Traditional Chinese herbal medicines (TCHMs) offer multi-component, multi-target anti-inflammatory potential; however, their bioactive constituents -polysaccharides, flavonoid glycosides, saponins and



alkaloids- are constrained by poor oral bioavailability. High molecular weight and extensive hydrogen bonding limit their bioaccessibility, i.e., release from the herbal matrix during gastrointestinal digestion, whilst limited membrane permeability, extensive first-pass metabolism and microbiota-dependent metabolic variability further restrict the fraction reaching systemic circulation in an active form (i.e., bioavailability), resulting in negligible systemic exposure and highly inconsistent therapeutic responses [5-7]. For instance, oral ginsenosides exhibit absolute bioavailability below 5% [8], whilst most flavonoid glycosides require microbial deglycosylation prior to absorption [9]. Overcoming these bottlenecks of poor absorption, poor targeting and high metabolic variability is essential for realising the intestinal anti-inflammatory efficacy of TCHMs.

Microbial fermentation addresses these constraints through targeted bioconversion [10]. In production animals, fermented herbal preparations have demonstrated direct efficacy: a probiotic-fermented complex of *Radix isatidis*, *Radix scutellariae* and *Radix glycyrrhizae* reduces diarrhoea incidence and improves growth performance in weaned piglets [11]; fermented soybean meal attenuates ETEC K88-induced PWD by modulating inflammatory cytokines and caecal microbiota [12]; and fermented herbal feed additives enrich beneficial *Lactobacillus* whilst improving feed conversion in piglets [13]. Mechanistically, fermentation depolymerises polysaccharides into bioactive oligosaccharides, hydrolyses glycosides into membrane-permeable aglycones, and modifies alkaloids and tannins to attenuate toxicity whilst preserving bioactivity [14]. Murine dextran sulphate sodium (DSS)-colitis models further indicate that fermented *Astragalus* suppresses colonic TNF- α , IL-1 β and IL-6, upregulates IL-10, and restores tight junction protein expression [15,16]. However, such murine findings serve solely as mechanistic explorations; their extrapolation to porcine PWD should be regarded as tentative given species-dependent differences in immune architecture, microbiota composition and disease pathogenesis-specifically, DSS induces chemical epithelial injury and Th1/Th17-mediated transmural inflammation, whereas porcine PWD is primarily driven by ETEC enterotoxins, fimbrial adhesion and weaning-induced barrier collapse [17].


Despite this growing body of work, the specific knowledge gap addressed by this review remains insufficiently served by the existing literature. Prior reviews of fermented TCHMs have been written primarily from phytochemical, pharmacological or human clinical perspectives, cataloguing component-content changes and *in vitro* bioactivities; the few reviews touching on animal production have largely listed growth-promoting effects without critically appraising the strength or origin

of the underlying evidence [18-20]. Three unresolved gaps persist. First, evidence obtained directly from fermented herbal preparations in livestock and poultry has not been systematically separated from mechanisms extrapolated from rodent colitis models, cell-culture systems or the general probiotic and SCFA literature, leaving the actual strength of support for proposed intestinal mechanisms unclear. Second, fermentation variables (microbial strain, substrate, temperature, pH, moisture, oxygen availability and duration) have rarely been linked to the resulting metabolite profiles and intestinal outcomes, which hampers standardisation and reproducibility. Third, safety and quality-consistency concerns specific to veterinary use (including microbial contamination, mycotoxins, toxic metabolites and batch-to-batch variation) remain largely unaddressed.

To address these gaps, the present narrative review makes two distinctive contributions. First, it centres specifically on the production-animal intestinal anti-inflammation dimension, explicitly distinguishing evidence derived directly from Chinese fermented herbal preparations in livestock from mechanisms extrapolated from general probiotic, SCFA or murine colitis literature. Second, it advances a unified “strain-ingredient-metabolite-microbiota-host” mechanistic framework that integrates fermentation biotransformation with gut microbiota remodelling and host tissue-level pathways, providing a conceptual foundation for standardising fermentation protocols and advancing targeted microbial fermentation strategies in veterinary practice. Fig. 1 provides an overview of this process: herbal materials are biotransformed by probiotic microbes into bioavailable metabolites that remodel gut microbiota, with downstream tissue level protective mechanisms presented in Fig. 4. The following sections examine fermentation microorganisms and metabolic profiles, condition-dependent phytochemical transformations, and gut anti-inflammatory mechanisms, before evaluating therapeutic efficacy in livestock production.

LITERATURE SEARCH AND REVIEW METHODOLOGY

This article is a narrative review; no systematic protocol was registered and the PRISMA framework was not applied. Literature was retrieved from PubMed, Web of Science, Scopus and CNKI up to June 2026, using combinations of terms covering fermentation (“fermentation”, “biotransformation”, “probiotic fermentation”), subject (“traditional Chinese medicine”, “Chinese herbal medicine”) and outcomes (“intestinal inflammation”, “gut microbiota”, “intestinal barrier”, “weaned piglet”, “broiler”, “livestock”); reference lists of retrieved articles were



screened for additional sources. Peer-reviewed English or Chinese publications examining microbial fermentation of Chinese medicinal herbs and reporting phytochemical transformations, fermentation conditions, intestinal or anti-inflammatory outcomes, or production performance were included; conference abstracts, dissertations and non-fermentation studies were excluded. No formal risk-of-bias assessment was performed; instead, evidence is appraised qualitatively throughout, distinguishing data obtained directly in livestock and poultry from rodent, cell-culture and general probiotic/SCFA evidence, and demonstrated effects from proposed mechanisms.

FUNDAMENTALS OF MICROBIAL FERMENTATION AND BIOTRANSFORMATION OF HERBAL ACTIVE COMPOUNDS

Microbial Strains and Metabolic Profiles in Chinese Herbal Fermentation

Chinese herbal fermentation typically employs *Lactobacillus*, *Bacillus* species, yeasts, and mixed microbial consortia. *Lactobacillus* produce lactic acid, acetic acid, exopolysaccharides, and glycoside hydrolases, thereby acidifying the fermentation substrate and facilitating glycoside biotransformation^[21]. *Bacillus* species demonstrate robust stress tolerance and secrete proteases, amylases, cellulases, and pectinases, rendering them well suited for the fermentation of herbal residues and fibrous feedstocks. Chang et al.^[22] developed a synbiotic formulation comprising probiotic-fermented herbal extracts and biogenic selenium nanoparticles, illustrating

the field's transition from monostrain fermentation to polymicrobial synbiotic systems. Strain selection dictates the trajectory of biotransformation. Co-fermentation with *Lactobacillus acidophilus* and *Bacillus subtilis* enhances the bioactivity of honeysuckle–Cassia seed extracts^[23]. Liao et al.^[24] optimised *B. subtilis*-mediated fermentation of the 'Four Yellow' formula and assessed its anti-*Escherichia coli* efficacy and growth-promoting activity. These findings suggest that future Chinese herbal fermentation should move beyond conventional probiotics towards strain screening based on defined biotransformation capacity and targeted anti-inflammatory functionality.

Effects of Fermentation Conditions on Active Compound Biotransformation

Fermentation conditions critically govern enzyme activity and biotransformation efficiency, thereby determining product quality. Temperature and pH fluctuations directly modulate enzyme secretion profiles and catalytic stability. During solid-state fermentation, temperature gradually increases during spore germination, then rises sharply owing to metabolic heat generation during rapid growth. Concomitantly, pH initially increases due to protein deamination and NH₃ release, followed by a sharp decline as organic acids accumulate and NH₃ is consumed. This pH-temperature coupling governs the temporal dynamics of enzyme production: pectinases and invertases peak during rapid pH decline, whereas xylanase and α-galactosidase synthesis is markedly delayed^[25]. pH directly modulates enzyme activity. For instance, recombinant AnGlu04478 exhibits optimal activity at pH 5.0 and retains substantial residual activity following 2 h incubation across pH 3.0–8.0. Its thermal optimum is

40°C, with robust stability between 25 and 50°C. Metal ions differentially modulate β -glucosidase activity: Na^+ , K^+ , Ca^{2+} and Mg^{2+} serve as activators or inhibitors, whereas SDS and EDTA consistently suppress enzyme function. Substrate induction also critically regulates enzyme secretion. For example, ginsenoside supplementation in *Aspergillus niger* fermentation markedly upregulates β -glucosidase gene expression, thereby enhancing rare ginsenoside biotransformation [26].

Moisture, oxygen availability and fermentation duration likewise shape biotransformation outcomes. Moisture governs substrate swelling, nutrient diffusion and enzyme secretion, and its optimum is strain-dependent: filamentous fungi tolerate low water activity, whereas bacteria and yeasts require 40-70% moisture [27,28]. Oxygen regime directs the transformation route-aerobic fungal processes favour oxidative bioconversion, whereas anaerobic lactic fermentation redirects metabolism towards organic acid accumulation [29]. Duration determines whether conversion halts at intermediates or reaches terminal metabolites, and over-incubation may degrade products through non-specific proteolysis [30]. These parameters act interactively, and their optimal windows are strongly strain-dependent (Table 1).

Organic acid accumulation indirectly promotes polysaccharide hydrolysis by lowering environmental pH. Lactic acid, acetic acid, and other fermentation-derived organic acids alter polysaccharide molecular weight and functional properties through pH-dependent mechanisms. Distinct microbial communities generate unique acid profiles, resulting in divergent polysaccharide structural and functional modifications. Concomitantly, fermentation-generated free radicals attack glycosidic bonds and side-chain moieties, inducing polysaccharide cleavage and structural remodeling. Such modifications

may unmask cryptic active sites, thereby enhancing antioxidant and immunomodulatory capacity [35].

Fermentation conditions also contribute substantially to poor inter-study reproducibility [36]. Variations in microbial strains, fermentation duration, or drying protocols applied to identical herbal materials yield divergent polysaccharide molecular weights, organic acid contents, and low-molecular-weight phenolic acid profiles. Fan et al. [37] compared the pro-digestive effects of *Massa Medicata Fermentata* (MMF) before and after processing, integrating metabolomic and microbiome analyses to demonstrate that processing methods concurrently modulate chemical composition and gut microbiota structure. Critically, most published studies report only a subset of relevant parameters; strain identity, moisture content and oxygen regime are the most frequently omitted, which not only accounts for much of the apparent inconsistency between studies but also precludes quantitative correlation of process parameters with metabolite outcomes. Claims that a given condition “enhances” bioactive transformation should accordingly be interpreted with caution unless the full parameter set is disclosed. Consequently, evaluating fermented Chinese herbal medicines requires a continuous evidence chain linking process parameters, metabolite profiles, and biological effects. Accordingly, Fig. 2 delineates the interrelationships among microbial strains, fermentation conditions, and metabolites.


EFFECTS OF FERMENTATION ON BIOACTIVE COMPOUNDS IN CHINESE HERBAL MEDICINES

Biotransformation of Polysaccharides and Glycosides

Polysaccharides constitute a critical material basis for the intestinal anti-inflammatory activity of Chinese

Table 1. Comparison of the principal microbial groups used in herbal fermentation and their condition-dependent biotransformation profiles

Microbial Group	Representative Genera	Key Enzymes	Typical Moisture/O ₂ Regime	Major Biotransformations	Ref
Lactic acid bacteria	<i>Lactiplantibacillus</i> , <i>Lactobacillus</i> , <i>Pediococcus</i>	β -glucosidases, esterases, proteases	Higher moisture (40-70% in SSF); anaerobic/microaerophilic	Glycoside $\rightarrow$ aglycone hydrolysis; organic acid accumulation; phenolic liberation	[31]
<i>Bacillus</i> spp.	<i>B. subtilis</i> , <i>B. licheniformis</i>	Proteases, cellulases, amylases	Moderate-high moisture; aerobic	Cell-wall disruption and release of bound phenolics; glycoside hydrolysis; peptide generation	[32]
Yeasts	<i>Saccharomyces</i> , <i>Candida</i>	β -glucosidases, invertase	Moderate moisture; facultative anaerobic	Glycoside hydrolysis; flavour-active metabolite formation; ethanol/ester production	[31]
Filamentous fungi	<i>Aspergillus</i> , <i>Rhizopus</i> , <i>Monascus</i>	Cellulases, xylanases, pectinases, tannase, β -glucosidases	Low water activity tolerated; strictly aerobic, gas-exchange-dependent	Lignocellulose degradation; extensive glycoside $\rightarrow$ aglycone conversion; alkaloid modification	[26,33]
Mixed/co-culture consortia	LAB + yeast + <i>Bacillus</i> combinations	Complementary enzyme repertoires	Stepwise or simultaneous; regime depends on dominant member	Sequential transformations unattainable by single strains	[34]



herbal medicines. Fermentation deploys polysaccharide hydrolases, cellulases, and pectinases to reduce polysaccharide molecular weight and increase soluble oligosaccharide content, thereby enhancing microbial utilisation. Yang et al. [38] demonstrated the prebiotic potential of fermented *Astragalus* polysaccharides, which modulate gut microbiota composition *in vitro*. Laoxianghuang polysaccharides also promote IL-10 production through gut microbiota-dependent linoleic acid metabolism [39]. These findings indicate that polysaccharide-mediated anti-inflammation relies on a sequential cascade from polysaccharide structure through microbial fermentation and metabolite generation to immune modulation. Glycoside biotransformation represents another critical mechanism for fermentation-enhanced bioavailability. Many flavonoids, saponins, and anthraquinones occur naturally as glycosides, which are highly polar and poorly absorbed [40]. Fermentation-derived β-glucosidases liberate aglycones, a transformation

experimentally demonstrated by direct compositional analyses [41–44]; by contrast, the subsequent epithelial absorption and inflammatory target engagement of the liberated aglycones remain largely inferred from studies of isolated, non-fermented compounds and have rarely been demonstrated for fermented preparations themselves. Ma et al. [41] screened flavonoid glycoside-hydrolysing strains for the bioconversion of Shuanghuanglian medicinal residue, enabling its valorisation. This study underscores that targeted screening of biotransformation-capable strains is pivotal for elucidating fermentation-enhanced efficacy.

Biotransformation of Flavonoids, Phenolic Acids, and Alkaloids

Flavonoids and phenolic acids represent the predominant bioactive classes underpinning the anti-inflammatory efficacy of fermented Chinese herbal medicines [42]. Fermentation releases conjugated phenolic acids,

remodels flavonoid glycoside-to-aglycone ratios, and generates novel phenolic metabolites ^[43]. Metabolomic analyses have documented marked phenolic remodelling during MMF fermentation, with differential metabolites primarily categorised into lipids, amino acids, phenolic acids, organic acids, flavonoids and alkaloids; structural equation modelling further identified *Bacillus velezensis* and fungal species as key drivers of flavonoid and phenolic acid accumulation ^[44]. Zhang et al. ^[45] demonstrated that chebulagic acid ameliorates DSS-induced colitis while restoring gut microbiota homeostasis and attenuating oxidative stress. It should be emphasised that chebulagic acid evidence derives from the DSS colitis model; its direct applicability to porcine ETEC-PWD remains speculative and requires species-specific validation.

Alkaloid biotransformation is highly dependent on both the herbal matrix and the microbial strain employed. Certain alkaloids undergo deglycosylation, demethylation, redox reactions, and conjugation, thereby modulating their polarity, toxicity, and anti-inflammatory potential. Investigations into intestinal absorption and component interactions in Forsythia-Honeysuckle formulae reveal potential absorptive and metabolic interplays among constituents ^[46]. Consequently, evaluating alkaloid-containing fermented formulations necessitates assessment beyond total alkaloid content to encompass constituent ratios, metabolite profiles, and safety.

Effects of Fermentation on Active Compound Physicochemical Properties

Fermentation modulates the molecular weight, solubility, acid-base properties, stability, and bioaccessibility of herbal bioactive compounds. Polysaccharide degradation enhances solubility and microbial fermentability; glycoside hydrolysis remodels polarity and membrane permeability; organic acid generation acidifies the microenvironment and suppresses pathogenic proliferation. Bai et al. ^[47] demonstrated that MMF modulates gut microbiota composition in rats with functional dyspepsia; as detailed above, Fan et al. ^[37] and Zheng et al. ^[44] concurrently established that MMF fermentation remodels phenolic metabolite profiles and microbial community structure. Fermentation also potentiates the anti-inflammatory activity of ox bile through NLRP3 inflammasome suppression ^[48]. Collectively, these findings indicate that physicochemical remodelling ultimately translates into microecological and immunological outcomes. *Fig. 3* delineates the biotransformation pathways and functional enhancement mechanisms of key bioactive compounds during fermentation. Overall, polysaccharide depolymerisation and glycoside-to-aglycone hydrolysis are supported by direct compositional evidence, whereas most downstream functional attributions remain proposed mechanisms that await targeted experimental validation.


Research Progress on the Enhancement of the Biological Activity of Traditional Chinese Medicinal Herbs Through Fermentation

In terms of biological activity, fermented Chinese herbal medicines primarily exhibit anti-inflammatory, antioxidant, gut microbiota-regulating and gut barrier-enhancing effects. Existing research has shown that fermented herbal preparations can exert protective effects both *in vivo* and *in vitro* under LPS stimulation ^[49]; fermented Pueraria root extract can alleviate DSS-induced colitis by reducing pro-inflammatory cytokine levels ^[50]; fermented *Astragalus* and its metabolites modulate inflammatory states, improve gut microbiota structure and promote intestinal barrier repair in DSS colitis mice ^[51]. These murine mechanistic insights, while informative, should not be equated with fermented-herb efficacy in porcine PWD, given the divergent aetiologies of chemical colitis versus bacterial enterotoxigenic diarrhoea. These results indicate that the synergistic effects of fermentation are not only reflected in the transformation and enhancement of bioactive components, but are also further demonstrated by improvements in the host's inflammatory response and intestinal function.

MECHANISMS OF ACTION OF FERMENTED CHINESE HERBS IN INTESTINAL ANTI-INFLAMMATION

Regulation of Gut Microbiota Composition and Metabolic Function

The gut microbiota is a key determinant of host health. A healthy gut microbiota typically exhibits high diversity, high gene richness, and stable metabolic function, all of which are crucial for maintaining intestinal homeostasis ^[52]. Disruption of the gut microbiota following weaning is a major cause of diarrhoea in piglets. Using 16S rRNA sequencing, Konstantinov et al. ^[53] and Lu et al. ^[54] analysed gut microbiota composition and metabolic shifts in piglets during the neonatal and weaning periods. The results indicated that the gut microbiota of neonatal piglets (2-12 days) was dominated by *E. coli* and various *Lactobacillus* species, with *Lactobacillus* predominating during the pre-weaning stage. Lactic acid and acetic acid produced by these bacteria maintain a low gut pH, thereby inhibiting pathogen colonisation. However, following weaning, *Lactobacillus* abundance declined significantly, whilst levels of potential pathogens, represented by *Clostridium* and *Escherichia*, gradually increased. This indicates that weaning significantly alters gut microbiota structure and metabolic activity, potentially increasing the risk of dysbiosis and diarrhoea. Such changes weaken lactic acid- and bacteriocin-mediated antibacterial effects, thereby reducing colonisation resistance against pathogenic bacteria.



In terms of gut microbiota regulation, fermented Chinese herbal medicines selectively enrich beneficial commensals that produce short-chain fatty acids [55]. Fermented bamboo shoots increase the abundance of *Akkermansia*, *Anaerovorax* and *Bacteroides* in the gut of DSS-induced colitis mice, while reducing the disease activity index [56]. Extrapolation of such murine DSS-colitis findings to porcine post-weaning diarrhoea should be regarded as tentative, given species- and model-dependent differences in pathogenesis, immune architecture and microbiota composition [57]. *Lactobacillus helveticus* H3 restores *Lactobacillus* and *Bifidobacterium* populations suppressed by pathogenic *E. coli*, while reducing *Escherichia-Shigella* colonization [58]. Fermentation-derived polyphenolic metabolites also serve as selective growth factors for *Lactobacillus* and *Bifidobacterium*, while inhibiting pathogens such as *Clostridioides difficile* and *E. coli* [59]. *Akkermansia muciniphila*, a mucin-degrading bacterium,

produces acetate, butyrate and propionate; its abundance negatively correlates with disease activity index and pro-inflammatory factor levels [60]. It must be stressed that these microbiota shifts are largely correlational: compositional changes demonstrated by 16S rRNA profiling do not, by themselves, establish a causal contribution to improved intestinal health. Direct mechanistic evidence for fermented herbal interventions -such as faecal microbiota transplantation or germ-free validation- remains scarce; a notable exception is the transplantation-validated tryptophan-metabolite pathway discussed below [61].

At the metabolite level, SCFAs, including acetate, propionate and butyrate, are key anti-inflammatory molecules derived from dietary fibre fermentation by intestinal beneficial bacteria [61]. As the primary energy source for colonic epithelial cells, butyrate exerts anti-inflammatory effects principally through GPR109A activation and histone deacetylase (HDAC)

inhibition, while its induction of Treg differentiation and IL-10 production is largely independent of NLRP3 inflammasome activation [62,63]. Functionally, butyrate strengthens intestinal epithelial barrier integrity by upregulating tight junction proteins (claudin-1, ZO-1 and occludin) via AMPK activation and alleviates cytokine-induced barrier injury by downregulating claudin-2 [64-66]. These protective effects have been validated in weaned piglets, where supplemental butyrate improved intestinal tight junction structure and reduced the incidence of post-weaning diarrhoea [67]. Notably, direct evidence supporting the barrier-protective and anti-inflammatory functions of herbal fermentation-derived butyrate against porcine PWD remains insufficient [68]. Unlike butyrate, acetate and propionate act mainly through GPR43 to trigger physiological NLRP3 inflammasome activation and subsequent IL-18 maturation, which sustains epithelial homeostasis and differs from pathological NLRP3 activation in intestinal inflammation [69]. Additionally, butyrate facilitates inflammation resolution by inducing Foxp3 Treg cells and IL-10 secretion via GPR43/GPR109A signalling and HDAC inhibition [70]. Moreover, propionate relieves intestinal inflammation by suppressing IL-8 and VCAM-1 expression through HDAC inhibition, whereas acetate primarily mediates epithelial protection via GPR43 signalling [62].

Beyond SCFAs, tryptophan metabolites such as indole-3-lactic acid (ILA) and indole-3-acetic acid (IAA) constitute another vital mechanistic axis underlying the beneficial effects of fermented Chinese herbal medicines. Solid-state fermented *Lactocaseibacillus paracasei* cultures enrich ILA and IAA, activate aryl hydrocarbon receptor (AhR) signalling and strengthen the mucosal barrier-effects abolished by antibiotic-induced microbiota depletion yet restored by faecal microbiota transplantation or metabolite supplementation [71,72]. These tryptophan metabolites stimulate the expression of IL-22, mucins and antimicrobial peptides via AhR activation, facilitating intestinal barrier repair and homeostasis [72,73].


Intestinal Barrier Function Improvement

Intestinal barrier damage is a key step in the progression from microbial dysbiosis to clinical diarrhoea. The intestinal barrier comprises mechanical, chemical, immune and biological barriers, with the mechanical barrier serving as the first line of defence against pathogen and harmful substance invasion [74]. Normal barrier integrity is maintained by the mucus layer, epithelial cells, tight junctions and mucosal immunity. Weaning stress reduces energy intake and epithelial renewal capacity in piglets, leading to decreased villus height and crypt hyperplasia, ultimately compromising intestinal barrier integrity. Liu et al. [75] and López-Colom et al. [76] demonstrated that enterotoxins, lipopolysaccharides and

pro-inflammatory factors produced by enterotoxigenic *E. coli* (ETEC) damage the intestinal mucosal layer and tight junction structures, increasing permeability and ultimately triggering intestinal inflammation.

Fermented Chinese herbal medicines repair the intestinal mucosal barrier through multiple mechanisms, as illustrated in Fig. 4, including upregulating tight junction protein expression, suppressing aberrant immune activation and enhancing mucosal immunity. Tight junction proteins are key molecules maintaining intestinal epithelial physical barrier integrity; fermentation products promote tight junction protein expression by regulating Hippo/YAP and PI3K/Akt/mTOR signalling pathways [77]. For example, in a direct production-animal study, a fermented herbal formulation (FHF) composed of *Radix isatidis*, *Folium isatidis*, *Radix scutellariae*, *Fructus forsythiae* and *Radix glycyrrhizae* -fermented with *Enterococcus faecium* and *Saccharomyces cerevisiae*-upregulated Claudin-1, Occludin and ZO-1 expression at both mRNA and protein levels in the jejunum of weaned piglets, concomitantly reducing serum diamine oxidase (DAO) and TNF- α levels [4]. Fermented plant extract (FPE) significantly reduces intracellular ROS levels in IPEC-J2 intestinal epithelial cells, restoring barrier integrity and cell migration capacity, indicating direct barrier-protective effects of fermented Chinese herbal medicines [78]. Importantly, increased expression of barrier-related proteins alone does not necessarily demonstrate improved intestinal permeability; functional evidence is available for only a subset of preparations, and most studies lack direct permeability measurements such as FITC-dextran flux, transepithelial electrical resistance or serum D-lactate.

Concurrently, as depicted in Fig. 4-C, fermentation-derived metabolites suppress aberrant TLR4/MyD88/NF- κ B signalling and pathological NLRP3 inflammasome activation in lamina propria immune cells, thereby reducing pro-inflammatory cytokines (TNF- α , IL-1 β and IL-6) and promoting anti-inflammatory IL-10 production [4]. Notably, while these signalling pathways are well characterised in general intestinal immunology, their direct inhibition by fermented herbal metabolites has been specifically demonstrated in LPS-stimulated IPEC-J2 cells and weaned piglets for FHF, and in THP-1 macrophages for fermented plant extracts [78]; extrapolation to other fermented preparations or species should remain cautious. Moreover, activation of the Nrf2 antioxidant pathway enhances HO-1, SOD and GPX expression, further reducing ROS accumulation and attenuating intestinal inflammation (Fig. 4) [79]. At the mucosal immunity level, enrichment of beneficial bacteria such as *A. muciniphila* increases mucus layer thickness, protects intestinal mucosal integrity and maintains immune homeostasis [60]. Furthermore, fermentation products enhance intestinal



chemical barrier function by promoting sIgA secretion, and maintain lamina propria immune cell homeostasis by regulating dendritic cell maturation and Treg/Th17 balance, thereby synergistically repairing the intestinal mucosa at both structural and immune barrier levels [69]. Collectively, these coordinated actions at the epithelial and immune levels synergistically restore intestinal mucosal homeostasis (Fig. 4). Nevertheless, it must be acknowledged that strong correlation with weak causality remains a limitation: microbiota changes may be either the cause or consequence of inflammation, and increased SCFA levels may directly protect the barrier or may simply accompany beneficial bacterial enrichment without proven causality in the context of fermented herbal interventions.

Modulating Inflammatory Response and Oxidative Stress

The immune-inflammatory response is also a key factor in post-weaning diarrhoea development in piglets. LPS is recognised by TLR4 and activates MyD88, IKK and NF-κB signalling, leading to TLR4 and NF-κB upregulation and subsequent induction of pro-inflammatory cytokines including TNF-α, IL-1β and IL-6. NF-κB activation not only promotes inflammatory factor release but also affects tight junction protein localisation and epithelial

cell apoptosis [80]. Fermentation-derived metabolites suppress aberrant TLR4/MyD88/NF-κB signalling and pathological NLRP3 inflammasome activation, thereby reducing TNF-α, IL-1β and IL-6 production whilst promoting anti-inflammatory IL-10 secretion and Nrf2-mediated antioxidant responses (Fig. 4). Thus, TLR4/NF-κB signalling modulation alleviates inflammation and serves as a key pathway linking pathogen expansion, exacerbated inflammation and barrier disruption. The JAK-STAT pathway is also involved in mucosal immune regulation following weaning stress in piglets [81].

Fermented Chinese herbal medicines block the intestinal inflammatory cascade by inhibiting pro-inflammatory signalling pathways, activating antioxidant defence systems and restoring the balance between pro- and anti-inflammatory factors [69]. It should be noted that many of these anti-inflammatory mechanisms are derived from general SCFA and polyphenol literature; where effects are demonstrated specifically with fermented herbal preparations, this is indicated explicitly. Regarding pro-inflammatory pathway inhibition, fermentation products act on mitogen-activated protein kinase and nuclear factor κB signalling. Extracts from fermented and aged mountain-cultivated ginseng shoots inhibit phosphorylation of extracellular signal-regulated kinase,

p38 and c-Jun N-terminal kinase in lipopolysaccharide-stimulated RAW 264.7 macrophages, and prevent NF- κ B p65 nuclear translocation, thereby reducing production of pro-inflammatory mediators including TNF- α , IL-1 β , IL-6 and inducible nitric oxide synthase [82]. In THP-1 macrophages, fermented plant extracts inhibit nuclear factor κ B activation via a Toll-like receptor 4-independent pathway, downregulating pro-inflammatory factors including tumour necrosis factor- α and monocyte chemotactic protein, while upregulating anti-inflammatory IL-10 [78]. *Lactobacillus* fermented ginseng sprout extract also inhibits lipopolysaccharide-induced nitric oxide and prostaglandin E2 production, and downregulates cyclooxygenase-2 and pro-inflammatory cytokine expression.

Regarding antioxidant stress, fermentation products activate the nuclear factor E2-related factor 2 (Nrf2)/antioxidant response element (ARE) signalling pathway. Under oxidative stress, Kelch-like ECH-associated protein 1 (Keap1) undergoes conformational changes, enabling Nrf2 dissociation, stabilisation and nuclear translocation. Following heterodimerisation with small Maf proteins, Nrf2 binds to ARE, thereby initiating transcription of downstream antioxidant enzymes including haem oxygenase-1 (HO-1), NAD(P)H:quinone oxidoreductase 1, superoxide dismutase and glutathione peroxidase [83]. Direct *in vivo* evidence for fermented herbal preparations in production animals demonstrates that one fermented herbal formulation upregulates Nrf2 and downstream antioxidant enzymes while decreasing MDA concentrations in weaned piglets [84]. Nevertheless, much of the canonical Nrf2/ARE signalling framework described above originates from general oxidative stress research and investigations using non-fermented herbal compounds. Traditional herbal formulations such as Gegen Qinlian Decoction counteract oxidative stress by activating the Nrf2/ARE pathway, with puerarin, berberine and glycyrrhizin as the primary active drivers [85]. A distinction is warranted between direct antioxidant effects -radical scavenging by fermentation-liberated phenolics, demonstrated largely in cell-free or cell-culture assays- and activation of endogenous antioxidant defence via the Nrf2/HO-1 axis, for which *in vivo* production-animal evidence remains limited to a small number of trials [84]. The former cannot be assumed to translate into *in vivo* redox modulation.

Notably, Nrf2 and NF- κ B pathways exhibit reciprocal cross-regulation [86]. The NF- κ B p65 subunit inhibits the Nrf2 pathway by competitively binding the transcriptional co-activator cyclic adenosine monophosphate response element-binding protein (CREB), or by recruiting histone deacetylase 3 (HDAC3) to induce local deacetylation [87]. Conversely, Nrf2 activation also suppresses NF- κ B

transcriptional activity. By simultaneously regulating these two pathways, fermented Chinese herbal medicines reduce pro-inflammatory factor production (TNF- α , IL-1 β , IL-6), while promoting anti-inflammatory interleukin-10 expression, thereby restoring the dynamic balance between pro- and anti-inflammatory factors. Based on microbiota, barrier, inflammatory and oxidative stress mechanisms, Fig. 4 summarises the key mechanisms by which fermentation-derived metabolites alleviate intestinal inflammation.

APPLICATION OF FERMENTED CHINESE HERBAL MEDICINES IN INTESTINAL INFLAMMATION

Applications in Livestock and Poultry Production

Available evidence from production-animal studies indicates that specific fermented herbal preparations exert species- and formulation-dependent effects on intestinal health and production performance, though the breadth of evidence varies considerably across livestock classes (Table 2).

In pigs, fermentation of Chinese herbal mixtures and composite preparations has been associated with improved growth performance and reduced diarrhoea incidence in weaned piglets challenged by weaning stress or enterotoxigenic *E. coli* [93,94]. A recent study demonstrated that a fermented herbal formulation (FHF) containing *Radix isatidis*, *Folium isatidis*, *Radix scutellariae*, *Fructus forsythiae* and *Radix glycyrrhizae* -fermented with *Enterococcus faecium* and *Saccharomyces cerevisiae*- improved average daily gain, reduced diarrhoea rate, enhanced jejunal tight junction protein expression (Claudin-1, Occludin, ZO-1), elevated acetic and butyric acids, and enriched *Limosilactobacillus* and *Lactobacillus* in weaned piglets [85]. These effects appear to be mediated through concurrent modulation of gut microbiota composition, enhancement of tight junction protein expression and attenuation of systemic inflammatory markers such as TNF- α [84,94]. Fermented herbal extracts have also been reported to improve nutrient digestibility during critical growth phases and to reduce environmental ammonia and hydrogen sulphide emissions in commercial pig production. Consistent with these findings, dietary supplementation with fermented Chinese herbal medicine feed additives improved growth performance and intestinal microbiota composition in weaned piglets [95], and probiotics-fermented *Massa Medicata Fermentata* ameliorated weaning stress by restoring intestinal homeostasis [96].

In poultry, *Clostridium perfringens*-induced necrotic enteritis (NE) remains a major enteric disease, with

Table 2. Comparative effects of fermented herbal preparations in production animals

Fermented Preparation	Fermentation Strain/Inoculum	Animal Model (challenge, if any)	Primary Endpoint(s)	Key Outcome(s)	Ref.
Chinese herbal mixture (4 kg/t diet)	Not specified	Weaned piglets	Growth performance, diarrhoea incidence	↑ Final weight, ↑ ADG, ↓ F/G ratio, ↓ diarrhoea incidence	[85]
FHF (Banlangen, Lianqiao, Huangqin, Gancao; 0.6% diet)	Probiotic consortium	Weaned piglets	Growth, serum inflammation, intestinal barrier, caecal microbiota	↑ Final body weight, ↑ ADG, ↓ FCR, ↓ diarrhoea, ↓ serum TNF-α and DAO, ↑ jejunal tight junction proteins, ↑ <i>Limosilactobacillus</i> and <i>Lactobacillus</i>	[4]
Fermented herbal extract (FHE)	Not specified	Weaning and fattening pigs	Nutrient digestibility, faecal gas emissions	Improved digestibility, ↓ ammonia and hydrogen sulphide emissions	[88]
Fermented cottonseed meal (8% diet)	Not specified	Broilers	Weight gain	↑ Weight gain by 3-4%	[89]
Fermented rapeseed meal (50% soybean meal replacement)	Not specified	Broilers	Weight gain, FCR	↑ Weight gain by 6-24%, ↓ FCR by 6-18%	[89]
Thymol/carvacrol (plant-derived additives)	Not specified	NE-challenged broilers	NE lesion scores, mortality	↓ Lesion scores, ↓ mortality	[90,91]
Fermented Chinese herbal additive	Probiotic consortium	Beef cattle	ADG, FCR, faecal pathogens	↑ ADG, ↓ FCR, ↓ faecal pathogen loads	[92]

estimated annual economic impacts exceeding several billion US dollars [95,96]. Fermented feed ingredients -including cottonseed meal and rapeseed meal- have shown potential in ameliorating NE-associated growth depression and intestinal lesions [95]. Additionally, plant-derived bioactive compounds such as thymol and carvacrol, whether delivered directly or via fermentation matrices, appear to reduce NE lesion scores and mortality through antimicrobial and immunomodulatory actions [91]. It should be noted that most poultry-related findings were obtained using fermented feed ingredients, not fermented Chinese herbal medicines as the test material. Caution is therefore required when extrapolating results to other plant species. More directly, dietary supplementation with fermented *Astragalus* powder improved growth performance and health status in broiler chickens [97].

In ruminants, research on fermented Chinese herbal medicines remains comparatively limited. Existing studies suggest that probiotic-fermented herbal additives may improve average daily gain and feed conversion efficiency in weaned lambs, whilst favourably modulating serum biochemistry and the intestinal microbiome [98]. Butyrate-producing fermentation consortia have also been implicated in enhancing ruminal epithelial barrier function and attenuating systemic inflammatory markers. However, the evidence base is insufficient to draw definitive conclusions regarding rumen-gut axis inflammation, and further species-specific trials are warranted.

Collectively, these studies demonstrate that the efficacy of fermented herbal interventions is highly dependent on the specific botanical matrix, fermentation strain, animal model and endpoint examined (Table 2). Rather than

supporting a generalised claim that “fermented herbs are effective,” the data highlight the need for standardised, comparative evaluation of distinct preparations across defined production contexts.

Research Progress in Experimental Models of Enteritis

Experimental colitis models provide controlled conditions for mechanistic studies. The DSS colitis model is commonly used to evaluate epithelial damage, dysbiosis and inflammatory pathways [97]. Fermented *Astragalus* and its metabolites modulate inflammatory status and gut microbiota in DSS colitis mice, whilst repairing the intestinal barrier [51]. Fermented Pueraria root extract reduces pro-inflammatory cytokines and alleviates DSS colitis [50]. Zhang et al. [45] found that chebulagic acid ameliorates oxidative stress, inflammation and gut microbiota dysbiosis in DSS colitis.

Beyond the DSS model, the 2,4,6-trinitrobenzenesulfonic acid (TNBS)-induced colitis model represents a well-established chemically-induced model that mimics key features of Crohn's disease, including transmural inflammation, Th1-dominant immune responses and granuloma formation [98]. In TNBS-induced colitis, rectal administration of TNBS in ethanol disrupts the colonic mucosal barrier and haptens colonic proteins, triggering a T-cell-mediated delayed-type hypersensitivity response characterised by elevated interferon-γ (IFN-γ), TNF-α and IL-12 production [99,100]. Fermented Chinese herbal medicines have demonstrated protective efficacy in this model through multiple mechanisms. Lian-Zhi-Fan (LZF) solution, a fermented preparation of Coptidis Rhizome, Gardeniae Fructus and alum, significantly ameliorates

body weight loss, disease activity index (DAI) score and colonic histopathological injury in TNBS-induced colitis rats ^[101]. Metabolomics analysis revealed that LZF modulates amino acid metabolism (L-alanine, L-valine), primary bile acid biosynthesis (cholic acid, taurocholic acid) and nicotinate/nicotinamide metabolism, thereby restoring energy production and attenuating inflammatory responses ^[102]. Additionally, berberine, a representative alkaloid from *Coptidis Rhizome*, attenuates TNBS-induced colitis by inhibiting inflammatory responses and Th1/Th17 differentiation, concomitantly with modulation of intestinal microbiota composition ^[101]. In a *Lactobacillus brevis* AL0035 fermentation study, oral administration of the probiotic strain counteracted TNBS-induced body weight loss, reduced DAI and histological injury scores, and decreased mRNA and protein expression of pro-inflammatory cytokines (TNF- α , IL-1 β , IL-6, IL-12, IFN- γ) whilst enhancing IL-10 expression ^[103,104]. Furthermore, AL0035 restored microbial diversity depleted by TNBS administration, increased the relative abundance of *Lactobacillaceae* and *Lachnospiraceae*, and decreased *Proteobacteria* abundance, indicating that fermentation-derived metabolites alleviate TNBS colitis through coordinated regulation of inflammatory cytokines, tight junction protein expression (ZO-1) and gut microbiota composition ^[105]. These findings collectively demonstrate that fermented Chinese herbal medicines exert intestinal protective effects in the TNBS model via immunomodulation, barrier repair and microbiota restoration, consistent with mechanisms observed in DSS and other colitis models.

The LPS stimulation model, antibiotic-induced injury model, *Salmonella gastroenteritis* model and intestinal ischaemia-reperfusion model are also employed to evaluate intestinal inflammation. Bose et al. utilised LPS stimulation to validate the protective effects of fermented herbal preparations ^[49], and demonstrated in a separate study that herbal preparations counteract neomycin-induced intestinal injury ^[102]. These models illustrate the intestinal protective potential of Chinese herbal medicines and their fermentation products from various perspectives, including pathogen-induced damage, drug-induced injury and barrier ischaemia.

Comparison and Evaluation of the Application Effects of Fermented Herbal Medicines

Evaluation focus for fermented Chinese herbal medicines varies across models. Production animal studies emphasise growth performance, diarrhoea, gut health, microbiota and immune parameters, whereas experimental colitis models focus on histopathology, inflammatory cytokines, oxidative stress and barrier proteins. Piglet studies by Wang et al. ^[4] and Chen et al. ^[85] align more closely with practical animal production applications. DSS-

induced colitis research presents a more comprehensive mechanistic chain, demonstrating that fermented *Astragalus* simultaneously influences inflammation, microbiota and intestinal barrier integrity ^[51]. TNBS-induced colitis studies provide complementary evidence for Th1-mediated immune modulation and transmural inflammation, with fermented preparations such as LZF solution and berberine demonstrating efficacy through metabolic pathway regulation and microbiota restoration ^[97]. Chebulagic acid studies, meanwhile, are more helpful in confirming the anti-inflammatory basis of specific *Terminalia chebula* components ^[45].

Medicinal herbs also differ in their fermentation outcomes. *MMF* research focuses on traditional fermented medicines, digestive function and microbiome regulation ^[106], as detailed in preceding sections. Fermented *Astragalus* research, meanwhile, emphasises polysaccharides, prebiotic effects, immunomodulation and barrier repair ^[51]. Co-fermented Chinese herbal medicine by-product research demonstrates that herbal residues yield antioxidant and anti-inflammatory benefits following appropriate fermentation ^[104]. Therefore, a blanket statement that 'fermented Chinese herbal medicines are effective' is inaccurate; rather, specific herbal materials, strains, metabolites, inflammatory models and evaluation endpoints must be clearly defined. Drawing on evidence from livestock and poultry production, experimental enteritis models and cellular studies, [Fig. 5](#) summarises the distribution of research evidence for the intestinal protective effects of fermented Chinese herbal medicines.

SAFETY, STANDARDISATION AND COMMERCIAL APPLICABILITY

The safety profile of fermented Chinese herbal medicines requires explicit consideration before routine veterinary application. Herbal raw materials and improperly controlled fermentations are vulnerable to microbial contamination and mycotoxins, which remain endemic in feed supply chains and can impair animal health and enter the food chain ^[107,108]. Although fermentation can degrade antinutritional factors and certain mycotoxins ^[15,16], uncontrolled fermentation may equally generate toxic metabolites or biogenic amines, and batch-to-batch variability in herbal starting materials further complicates safety evaluation ^[92,107]. Adverse effects have rarely been reported in the studies reviewed here, but they have also rarely been monitored systematically; future trials should incorporate explicit safety endpoints, including mycotoxin screening, microbial limits and histopathological examination ^[107,108].

Standardisation is a prerequisite for reproducible efficacy. Future studies should report, as a minimum: (i) unambiguous strain identification, preferably based on




Fig 5. Overview of evidence supporting the gut-protective effects of fermented Chinese herbal medicines. (A) Studies in piglets have shown that fermented Chinese herbal medicines can improve growth performance, reduce the incidence of diarrhoea and the feed-to-meat ratio (FCR), increase villus height and SCFA levels, promote the proliferation of beneficial bacteria, alleviate oxidative stress and improve intestinal barrier function; (B) Studies in broiler chickens have shown that fermented Chinese herbal medicines can improve production performance and intestinal morphology, increase SCFA levels and the abundance of beneficial bacteria, alleviate oxidative stress and enhance barrier integrity; (C) Studies in DSS/TNBS-induced colitis models in mice have found that fermented Chinese herbal medicines can reduce disease severity and tissue damage, increase SCFA levels and the abundance of beneficial bacteria, enhance the expression of barrier proteins, and inhibit inflammation and oxidative stress; (D) Studies in ruminants have shown that fermented Chinese herbal medicines can increase feed intake, milk yield and nutrient digestibility, improve ruminal fermentation and microbial balance, and reduce methane emissions; (E) Cell and organoid studies have shown that fermentation metabolites can enhance cell viability, strengthen epithelial barrier function, promote the expression of tight junction proteins and MUC2, reduce levels of inflammatory cytokines and ROS, and activate antioxidant pathways such as Nrf2; (F) Multi-omics studies, including metabolomics, microbiomics, transcriptomics and proteomics, have further elucidated the molecular mechanisms by which fermented Chinese herbal medicines regulate the gut microbiota, host metabolism and immune responses

whole-genome sequencing with deposition in recognised culture collections; (ii) fully defined fermentation parameters, including substrate composition, inoculum size, temperature, pH, moisture, oxygen regime and duration; (iii) chemical standardisation based on marker compounds or quality markers (Q-markers) linked to the intended bioactivity^[109]; and (iv) untargeted metabolomic fingerprinting to confirm batch consistency and product stability during storage^[32,33]. Only with such controls can dose–response relationships be compared meaningfully across studies and preparations.

From a commercial perspective, the doses used in experimental studies warrant critical appraisal. Inclusion levels reported in piglet trials (for example, 0.6% dietary inclusion of a fermented herbal formulation^[4]) are within ranges considered economically feasible for commercial feed formulation, whereas high-dose extract regimens or short-term intensive administration may be impractical under field conditions^[88]. Cost-effectiveness will depend on substrate availability, fermentation scalability and product stability; solid-state fermentation of agro-industrial residues offers a low-cost production

platform in this regard^[27,30]. Expressing doses in terms of defined marker-compound content rather than raw herb equivalents would further facilitate commercial translation and regulatory evaluation.

CONCLUSION

Existing research supports the intestinal anti-inflammatory potential of fermented Chinese herbal medicines, though evidence quality varies considerably. Early studies demonstrate protective effects against LPS- and neomycin-induced intestinal injury, yet lack in-depth analysis of ingredient biotransformation and microbiota-mediated mechanisms^[49]. Trials conducted in production animals fit practical breeding scenarios, but complex dietary and environmental confounding factors restrict the interpretation of causal relationship^[84]. Although murine DSS colitis models enable relatively complete mechanistic exploration, findings cannot be reliably extrapolated to porcine post-weaning diarrhoea due to interspecies difference^[51].

Studies on flavonoid glycoside hydrolysis and fermentation-induced phenolic remodelling provide

representative evidence for active ingredient biotransformation^[41,43]. Nevertheless, numerous animal experiments lack quantitative compositional comparisons pre- and post-fermentation, obstructing the identification of truly bioactive compounds.

Mechanistically, most investigations converge on the “microbiota-barrier-inflammation” axis; however, widespread issues of strong correlation without confirmed causality remain unresolved. Alterations in gut microbiota may occur either prior to or secondary to inflammation, and elevated SCFA concentrations could exert direct barrier protection or merely accompany the enrichment of beneficial commensals. Rigorous validation via faecal microbiota transplantation, metabolite supplementation and receptor blockade experiments is required to establish definitive causal links^[45]. Future high-quality research should adopt multi-omics screening to uncover candidate pathways, followed by targeted functional verification.

Industrial standardisation constitutes a major bottleneck. Large discrepancies in starter strains, substrate proportion, fermentation humidity, temperature, duration and drying protocols hinder cross-study comparison of product composition and biological efficacy^[24]. It is urgent to build a systematic quality evaluation system covering raw material fingerprints, fermentation termination indicators, quantitative detection of active ingredients and safety assessment.

Future investigations should develop targeted fermentation strategies within the “strain-substrate-metabolite-host” conceptual framework. Researchers can screen functional strains capable of transforming specific herbal components and select tailored fermented products according to enteritis phenotypes and baseline intestinal microbiota profiles^[101]. Microbial fermentation improves the bioaccessibility of herbal constituents via polysaccharide degradation, glycoside hydrolysis and phenolic compound release. The resulting fermented preparations regulate gut microbiota, reinforce intestinal barrier integrity, inhibit NF- κ B/NLRP3 signalling and mitigate oxidative stress. To fully exploit their application prospects in livestock production, the field needs to address inconsistent fermentation protocols, ambiguous strain selection criteria, insufficient evidence of component transformation, inadequate mechanistic validation and scarce large-scale field trials. Ultimately, a complete and reproducible evidence chain spanning strain, bioactive components, metabolites, gut microbiota, host responses and *in vivo* efficacy should be established. In summary, the most reliable effects of fermented Chinese herbal medicine in production animals are the combinatorial biotransformation of herbal components and the promotion of growth performance, along with mechanisms of barrier protection, anti-inflammation, and

microbiota regulation. Although biologically plausible and increasingly supported by evidence, these effects are still at an early stage and require verification in target animals through standardized and controllable dosage trials, while also clearly addressing issues such as safety, standardization, and commercial feasibility.

DECLARATIONS

Availability of Data and Materials: As this is a review article, no original datasets were generated or analyzed. All information discussed in this review is provided within the manuscript and its cited references.

Acknowledgements: The author sincerely thanks Xizang Agricultural and Animal Husbandry University, College of Animal Science, for providing literature resources and academic support for this review. Gratitude is also extended to the journal reviewers and editors for their constructive revision suggestions, which substantially improved the manuscript.

Funding Support: This study was supported by the grants from Talent Team Building at Xizang Agricultural and Animal Husbandry University (XZNMDXRCDW-2026-05), Demonstration Base of Grassroots Agricultural Technology Extension System, Xizang Autonomous Region, and Plateau-Featured Discipline Development and Talent Training Quality Enhancement Project (YJSXK202614).

Competing Interests: The authors declare no conflicts of interest.

Declaration on Generative Artificial Intelligence (AI): Generative artificial intelligence was only used for language polishing in this study; it was not involved in any core research. All authors take full responsibility for the authenticity and scientific rigor of the manuscript.

Author Contributions: Every author took part in compiling this review and offered valuable input. Y.Z. formulated the review framework and finished the initial draft. Z.W. screened massive literatures and organized research progress data. D. supplemented localized research references. H.L. summarized domestic research outcomes. Y.X. managed and classified all collected literatures. Z.H. polished the text and standardized manuscript layout. Q.K., the corresponding author, supervised the whole writing work, obtained supporting funds, and thoroughly revised the complete manuscript.

REFERENCES

1. Xie H, Yu S, Tang M, Xun Y, Shen Q, Wu G: Gut microbiota dysbiosis in inflammatory bowel disease: Interaction with intestinal barriers and microbiota-targeted treatment options. *Front Cell Infect Microbiol*, 15:1608025, 2025. DOI: 10.3389/fcimb.2025.1608025
2. Liu H, Chen R, Wen S, Li Q, Lai X, Zhang Z, Sun L, Sun S, Cao F: Tea (*Camellia sinensis*) ameliorates DSS-induced colitis and liver injury by inhibiting TLR4/NF- κ B/NLRP3 inflammasome in mice. *Biomed Pharmacother*, 158:114136, 2023. DOI: 10.1016/j.biopha.2022.114136
3. Xing D, Zheng T, Chen X, Xie Z: Yellow teas protect against DSS-induced ulcerative colitis by inhibiting TLR4/NF- κ B/NLRP3 inflammasome in mice. *Foods*, 13 (17):2843, 2024. DOI: 10.3390/foods13172843
4. Wang X, Fan X, Li C, Chen P, Li S, Peng J, Zhou W, Zhou Z, Li X, Li J, Xiao Y: A Fermented herbal formulation improves intestinal health and growth performance in post-weaning piglets. *Animals*, 16 (8):1254, 2026. DOI: 10.3390/ani16081254
5. Qiu Y, Tang J, Wang L, Yang X, Jiang Z: Fermented corn-soybean meal improved growth performance and reduced diarrhea incidence

- by modulating intestinal barrier function and gut microbiota in weaned piglets. *Int J Mol Sci*, 25 (6):3199, 2024. DOI: 10.3390/ijms25063199
6. Wang Y, Li J, Xie Y, Zhang H, Jin J, Xiong L, Liu H: Effects of a probiotic-fermented herbal blend on the growth performance, intestinal flora and immune function of chicks infected with *Salmonella pullorum*. *Poult Sci*, 100 (7):101196, 2021. DOI: 10.1016/j.psj.2021.101196
 7. Shidore T, Buhr DL, Morano J, Dhlakama T, Baxter M, Lum J, Barton JT, Ritch MD, Westgate B, Moscato ZM, Fiandaca MJ, Sevostyanova A, Kiebler C, Zwilling MF, Copley C, Kiss MM: Detection of necrotic enteritis risk through non-invasive monitoring of *Clostridium perfringens* in feces. *Poult Sci*, 104 (2):104809, 2025. DOI: 10.1016/j.psj.2025.104809
 8. Emami NK, Dalloul RA: Centennial review: Recent developments in host-pathogen interactions during necrotic enteritis in poultry. *Poult Sci*, 100 (9):101330, 2021. DOI: 10.1016/j.psj.2021.101330
 9. Yuan S, Wang Q, Li J, Xue JC, Li Y, Meng H, Hou XT, Nan JX, Zhang QG: Inflammatory bowel disease: an overview of Chinese herbal medicine formula-based treatment. *Chin Med*, 17 (1):74, 2022. DOI: 10.1186/s13020-022-00633-4
 10. Deng Y, Song L, Huang J, Zhou W, Liu Y, Lu X, Zhao H, Liu D: *Astragalus* polysaccharides ameliorates experimental colitis by regulating memory B cells metabolism. *Chem Biol Interact*, 394:110969, 2024. DOI: 10.1016/j.cbi.2024.110969
 11. Li R, Liao X, Fu X, Li X, Liao X, Cen S, Zeng J, Huang L, Chi H, Zou Y: Microbiota-driven tryptophan metabolism and AhR triggered intestinal stem cell differentiation: mechanisms of huangqin decoction in ulcerative colitis repair. *Chin Med*, 21 (1):33, 2026. DOI: 10.1186/s13020-025-01302-y
 12. Li B, Du P, Du Y, Zhao D, Cai Y, Yang Q, Guo Z: Luteolin alleviates inflammation and modulates gut microbiota in ulcerative colitis rats. *Life Sci*, 269:119008, 2021. DOI: 10.1016/j.lfs.2020.119008
 13. Chen C, Feng F, Qi M, Chen Q, Tang W, Diao H, Hu Z, Qiu Y, Li Z, Chu Y, Tang Z: Dietary citrus flavonoids improved growth performance and intestinal microbiota of weaned piglets via immune function mediated by TLR2/NF- κ B signaling pathway. *J Agric Food Chem*, 72 (30): 16761-16776, 2024. DOI: 10.1021/acs.jafc.4c03401
 14. Jiang X, Xiao H, Huang P: Mechanism of action of plant extracts in preventing post-weaning diarrhea in piglets: A review. *Vet Sci*, 13 (4):312, 2026. DOI: 10.3390/vetsci13040312
 15. Girard M, Bee G: Invited review: Tannins as a potential alternative to antibiotics to prevent coliform diarrhea in weaned pigs. *Animal*, 14 (1): 95-107, 2020. DOI: 10.1017/S1751731119002143
 16. Biagia G, Cipollini I, Paulicks BR, Roth FX: Effect of tannins on growth performance and intestinal ecosystem in weaned piglets. *Arch Anim Nutr*, 64 (2): 121-135, 2010. DOI: 10.1080/17450390903461584
 17. Xu T, Ma X, Zhou X, Qian M, Yang Z, Cao P, Han X: Coated tannin supplementation improves growth performance, nutrients digestibility, and intestinal function in weaned piglets. *J Anim Sci*, 100 (5):skac088, 2022. DOI: 10.1093/jas/skac088
 18. Wang Y, Wang J, Zhang L, Han S, Pan X, Wen H, Yang H, Wang X, Peng D: Veterinary drug residues in food chains: Sources, exposure pathways, health impacts, mitigation, and safety assurance. *Foods*, 15 (5):840, 2026. DOI: 10.3390/foods15050840
 19. Cheng WW, Liu BH, Hou XT, Meng H, Wang D, Zhang CH, Yuan S, Zhang QG: Natural products on inflammatory bowel disease: Role of gut microbes. *Am J Chin Med*, 52 (5): 1275-1301, 2024. DOI: 10.1142/s0192415x24500514
 20. Hu QR, Hong H, Zhang ZH, Feng H, Luo T, Li J, Deng ZY, Chen F: Methods on improvements of the poor oral bioavailability of ginsenosides: Pre-processing, structural modification, drug combination, and micro- or nano- delivery system. *J Ginseng Res*, 47 (6): 694-705, 2023. DOI: 10.1016/j.jgr.2023.07.005
 21. Wang Z, Jin X, Zhang X, Xie X, Tu Z, He X: From function to metabolome: metabolomic analysis reveals the effect of probiotic fermentation on the chemical compositions and biological activities of *Perilla frutescens* leaves. *Front Nutr*, 9:933193, 2022. DOI: 10.3389/fnut.2022.933193
 22. Chang J, Zeng X, Zhang R, Shen H, Xu C: Preparation and evaluation of microecological agents incorporating probiotic-fermented traditional Chinese medicine and biogenic selenium nanoparticles for intestinal protection. *Biol Trace Elem Res*, 204 (3): 1356-1373, 2026. DOI: 10.1007/s12011-025-04753-4
 23. Zhou M, Zhang Y, Song Z, Tang S, Liu Z, Pang M, Zhang D, Wu X, Yu X, Wang P, Wei Y: Enhanced bioactivity of honeysuckle-Cassia seeds extracts through *Lactobacillus acidophilus* and *Bacillus subtilis* co-fermentation: Impact on alcoholic liver disease and gut microbiota. *Food Chem*, 486:144463, 2025. DOI: 10.1016/j.foodchem.2025.144463
 24. Liao L, Zhang W, Liu Y, Chen J, Sun Y, Liu X: Optimization of fermentation process for Sihuang compound with *Bacillus subtilis* and its efficacy against *Escherichia coli* infections and growth performance in broilers. *Poult Sci*, 104 (10):105588, 2025. DOI: 10.1016/j.psj.2025.105588
 25. Orlando M, Marchetti A, Bombardi L, Lotti M, Fusco S, Mangiagalli M: Polysaccharide degradation in an Antarctic bacterium: Discovery of glycoside hydrolases from remote regions of the sequence space. *Int J Biol Macromol*, 299:140113, 2025. DOI: 10.1016/j.ijbiomac.2025.140113
 26. Li FX, Lin DM, Yang J, Cui XM, Yang XY: The transformation pathways and optimization of conditions for preparation minor ginsenosides from *Panax notoginseng* root by the fungus *Aspergillus tubingensis*. *PLoS One*, 20 (3):e0316279, 2025. DOI: 10.1371/journal.pone.0316279
 27. Santos TC, Cavalcanti IS, Bonomo RCF, Santana NB, Franco M: Optimization of productions of cellulolytic enzymes by *Aspergillus niger* using residue of mango as a substrate. *Ciência Rural*, 41 (12): 2210-2216, 2011. DOI: 10.1590/S0103-84782011005000145
 28. Pandey A: Solid-state fermentation. *Biochem Eng J*, 13 (2-3): 81-84, 2003. DOI: 10.1016/S1369-703X(02)00121-3
 29. Rahardjo YSP, Weber FJ, le Comte EP, Tramper J, Rinzeema A: Contribution of aerial hyphae of *Aspergillus oryzae* to respiration in a model solid-state fermentation system. *Biotechnol Bioeng*, 78 (5): 539-554, 2002. DOI: 10.1002/bit.10222
 30. Kang SW, Park YS, Lee JS, Hong SI, Kim SW: Production of cellulases and hemicellulases by *Aspergillus niger* KK2 from lignocellulosic biomass. *Bioresour Technol*, 91 (2): 153-156, 2004. DOI: 10.1016/S0960-8524(03)00172-X
 31. Paventi G, Di Martino C, Coppola F, Iorizzo M: β -glucosidase activity of *Lactiplantibacillus plantarum*: A key player in food fermentation and human health. *Foods*, 14 (9):1451, 2025. DOI: 10.3390/foods14091451
 32. Chakraborty T, Kashyap SJ, Chetia PD, Purkayastha S: Gut microbiota-mediated biotransformation of dietary bioactive compounds: Inter-individual variability, determinants, and implications for personalised nutrition. *Front Syst Biol*, 6:1898890, 2026. DOI: 10.3389/fsysb.2026.1898890
 33. Li L, Wang L, Fan W, Jiang Y, Zhang C, Li J, Peng W, Wu C: The application of fermentation technology in traditional Chinese medicine: A review. *Am J Chin Med*, 48 (4): 899-921, 2020. DOI: 10.1142/S0192415X20500433
 34. Ding P, Wang X, Jiang S, Li M, He X, Peng Y: Effects of bacteria-enzyme co-fermented Chinese herbal medicine on growth performance, apparent nutrient digestibility, meat quality, and immune function in broilers. *Front Vet Sci*, 12:1676951, 2025. DOI: 10.3389/fvets.2025.1676951
 35. Sammulia SF, Suhaera S, Prayoga DK, Pitriani P, Ramadhania ZM, Fristiohady A, Milanda T: Fermentation-induced changes in phytochemical composition and pharmacological activities of *Zingiberaceae* plants: Insight from *in vitro* and *in vivo* studies. *Drug Des Dev Ther*, 20:604038, 2026. DOI: 10.2147/DDDT.S604038
 36. Wang T, Sheng K, Zhang Y, Jin S, Feng L, Wang L: Metabolomics analysis reveals the effect of fermentation on the chemical composition and antioxidant activity of *Paenonia lactiflora* root. *Heliyon*, 10 (7):e28450, 2024. DOI: 10.1016/j.heliyon.2024.e28450

37. Fan S, Zhu H, Liu W, Ha J, Liu Y, Mi M, Ren Q, Xu L, Zhang J, Liu W, Feng F, Xu J: Comparing massa medicata fermentata before and after charred in terms of digestive promoting effect via metabolomics and microbiome analysis. *J Ethnopharmacol*, 327:117989, 2024. DOI: 10.1016/j.jep.2024.117989
38. Yang P, Zhou Q, Zhang Y, Jia M, Li R, Qu Q, Li Z, Feng M, Tian Y, Ren W, Peng X, Shi X: Exploring the prebiotic potential of fermented *Astragalus* polysaccharides on gut microbiota regulation *in vitro*. *Curr Microbiol*, 82 (1):52, 2024. DOI: 10.1007/s00284-024-04035-7
39. Zhang Y, Yang Y, Song J, Yu W, Li Y, Liu D, Gao J, Fan B, Wang F, Zheng Y: Laoxianghuang polysaccharide promotes the anti-inflammatory cytokine interleukin-10 in colitis via gut microbial linoleic acid. *Phytomedicine*, 135:156136, 2024. DOI: 10.1016/j.phymed.2024.156136
40. He Y, Hu Z, Li A, Zhu Z, Yang N, Ying Z, He J, Wang C, Yin S, Cheng S: Recent advances in biotransformation of saponins. *Molecules*, 24 (13):2365, 2019. DOI: 10.3390/molecules24132365
41. Ma X, Chen Y, Li L, Wang T, Teng K, Su J, Li L, Li L, Li H, Diao W, Chen G, Wang J, Zhong J: Efficient utilization of Shuanghuanglian medicine residues by microbial transformation with flavonoid glycosides-hydrolyzing strains. *Front Microbiol*, 16:1553399, 2025. DOI: 10.3389/fmicb.2025.1553399
42. Chen L, Cao H, Huang Q, Xiao J, Teng H: Absorption, metabolism and bioavailability of flavonoids: A review. *Crit Rev Food Sci Nutr*, 62 (28): 7730-7742, 2022. DOI: 10.1080/10408398.2021.1917508
43. Kim J, Choi KB, Park JH, Kim KH: Metabolite profile changes and increased antioxidative and antiinflammatory activities of mixed vegetables after fermentation by *Lactobacillus plantarum*. *PLoS One*, 14 (5):e0217180, 2019. DOI: 10.1371/journal.pone.0217180
44. Zheng W, Li J, Yin L, Sun J, Liu L, Kong B, Zhu Y, Xu N: Fermentation-driven changes in phenolic compounds of Massa Medicata Fermentata. *Biomed Chromatogr*, 40 (3):e70394, 2026. DOI: 10.1002/bmc.70394
45. Zhang S, Ren Y, Li M, Gao X, Zhang X, Xu W, Lu Q: Chebulagic acid ameliorates DSS-induced colitis in mice by improving oxidative stress, inflammation and the gut microbiota. *Am J Transl Res*, 17 (6): 4101-4118, 2025. DOI: 10.62347/QWCQ2468
46. He JY, Xiao FL, Zheng QY, Wang CH, Tang YY, Fu JX, Huang JY, Zhou LD, Zhang QH: Intestinal absorption characteristics and reciprocal interactions of *Forsythiae fructus* and *Lonicerae japonicae* flos-containing Chinese herbal formulation with human gut microbiome. *Am J Chin Med*, 53 (2): 543-566, 2025. DOI: 10.1142/S0192415X25500211
47. Bai Y, Zheng M, Fu R, Du J, Wang J, Zhang M, Fan Y, Huang X, Li Z: Effect of Massa Medicata Fermentata on the intestinal flora of rats with functional dyspepsia. *Microb Pathog*, 174:105927, 2023. DOI: 10.1016/j.micpath.2022.105927
48. Pei Y, Cheng F, Li W, Yu Q, Ma C, Zou Y, Xu T, Liu S, Zhang S, Wang Q: Enhancement of anti-inflammatory effect of cattle bile by fermentation and its inhibition of neuroinflammation on microglia by inhibiting NLRP3 inflammasome. *J Biosci Bioeng*, 133 (2): 146-154, 2022. DOI: 10.1016/j.jbiosc.2021.11.003
49. Bose S, Song MY, Nam JK, Lee MJ, Kim H: *In vitro* and *in vivo* protective effects of fermented preparations of dietary herbs against lipopolysaccharide insult. *Food Chem*, 134 (2): 758-765, 2012. DOI: 10.1016/j.foodchem.2012.02.175
50. Choi S, Woo JK, Jang YS, Kang JH, Jang JE, Yi TH, Park SY, Kim SY, Yoon YS, Oh SH: Fermented *Pueraria lobata* extract ameliorates dextran sulfate sodium-induced colitis by reducing pro-inflammatory cytokines and recovering intestinal barrier function. *Lab Anim Res*, 32 (3): 151-159, 2016. DOI: 10.5625/lar.2016.32.3.151
51. Li J, Ma Y, Li X, Wang Y, Huo Z, Lin Y, Li J, Yang H, Zhang Z, Yang P, Zhang C: Fermented *Astragalus* and its metabolites regulate inflammatory status and gut microbiota to repair intestinal barrier damage in dextran sulfate sodium-induced ulcerative colitis. *Front Nutr*, 9:1035912, 2022. DOI: 10.3389/fnut.2022.1035912
52. Adak A, Khan MR: An insight into gut microbiota and its functionalities. *Cell Mol Life Sci*, 76 (3): 473-493, 2019. DOI: 10.1007/s00018-018-2943-4
53. Konstantinov SR, Awati AA, Williams BA, Miller BG, Jones P, Stokes CR, Akkermans ADL, Smidt H, de Vos WM: Post-natal development of the porcine microbiota composition and activities. *Environ Microbiol*, 8 (7): 1191-1199, 2006. DOI: 10.1111/j.1462-2920.2006.01009.x
54. Huang S, Cui Z, Hao X, Cheng C, Chen J, Wu D, Luo H, Deng J, Tan C: Dietary fibers with low hydration properties exacerbate diarrhea and impair intestinal health and nutrient digestibility in weaned piglets. *J Anim Sci Biotechnol*, 13 (1):142, 2022. DOI: 10.1186/s40104-022-00771-7
55. Atarashi K, Tanoue T, Shima T, Imaoka A, Kuwahara T, Momose Y, Cheng G, Yamasaki S, Saito T, Ohba Y, Taniguchi T, Takeda K, Hori S, Ivanov II, Umesaki Y, Itoh K, Honda K: Induction of colonic regulatory T cells by indigenous *Clostridium* species. *Science*, 331 (6015): 337-341, 2011. DOI: 10.1126/science.1198469
56. Lu X, Xu Y, Zhang C, Liu H, Xie J, Tian B: Bamboo shoot fermented products alleviated DSS-induced ulcerative colitis in mice by effectively controlling inflammatory reactions and adjusting the gut microbiota and its metabolites. *Front Nutr*, 12:1724148, 2025. DOI: 10.3389/fnut.2025.1724148
57. Wirtz S, Popp V, Kindermann M, Gerlach K, Weigmann B, Fichtner-Feigl S, Neurath MF: Chemically induced mouse models of acute and chronic intestinal inflammation. *Nat Protoc*, 12 (7): 1295-1309, 2017. DOI: 10.1038/nprot.2017.044
58. Zhang Z, Lv J, Wang X, Chun L, Yang Q, Zhao H, Xue S, Zhang Z, Liu X, Wang S, Sun Y: *Lactobacillus helveticus* mitigates diarrhea and inflammation induced by enterotoxigenic *E. coli* through rebalance of gut microbiota. *Curr Res Food Sci*, 11:101147, 2025. DOI: 10.1016/j.crfs.2025.101147
59. Alharbi NA: Polyphenol metabolites in fermented foods: Biotransformation, bioavailability, and functional roles. *Front Nutr*, 13:1767453, 2026. DOI: 10.3389/fnut.2026.1767453
60. Ding X, Meng PF, Ma XX, Yue JY, Li LP, Xu LR: *Akkermansia muciniphila* and herbal medicine in immune-related diseases: Current evidence and future perspectives. *Front Microbiomes*, 3:1276015, 2024. DOI: 10.3389/frmbi.2024.1276015
61. Smith PM, Howitt MR, Panikov N, Michaud M, Gallini CA, Bohlooly YM, Glickman JN, Garrett WS: The microbial metabolites, short-chain fatty acids, regulate colonic T_{reg} cell homeostasis. *Science*, 341 (6145): 569-573, 2013. DOI: 10.1126/science.1241165
62. Li M, van Esch B, Henricks PAJ, Folkerts G, Garssen J: The anti-inflammatory effects of short chain fatty acids on lipopolysaccharide- or tumor necrosis factor α -stimulated endothelial cells via activation of GPR41/43 and inhibition of HDACs. *Front Pharmacol*, 9:533, 2018. DOI: 10.3389/fphar.2018.00533
63. Furusawa Y, Obata Y, Fukuda S, Endo TA, Nakato G, Takahashi D, Nakanishi Y, Uetake C, Kato K, Kato T, Takahashi M, Fukuda NN, Murakami S, Miyauchi E, Hino S, Atarashi K, Onawa S, Fujimura Y, Lockett T, Clarke JM, Topping DL, Tomita M, Hori S, Ohara O, Morita T, Koseki H, Kikuchi J, Honda K, Hase K, Ohno H: Commensal microbe-derived butyrate induces the differentiation of colonic regulatory T cells. *Nature*, 504 (7480): 446-450, 2013. DOI: 10.1038/nature12721
64. Pérez-Reytor D, Puebla C, Karahanian E, García K: Use of short-chain fatty acids for the recovery of the intestinal epithelial barrier affected by bacterial toxins. *Front Physiol*, 12:650313, 2021. DOI: 10.3389/fphys.2021.650313
65. Peng L, Li ZR, Green RS, Holzman IR, Lin J: Butyrate enhances the intestinal barrier by facilitating tight junction assembly via activation of AMP-activated protein kinase in Caco-2 cell monolayers. *J Nutr*, 139 (9): 1619-1625, 2009. DOI: 10.3945/jn.109.104638
66. Huang X, Oshima T, Tomita T, Fukui H, Miwa H: Butyrate alleviates cytokine-induced barrier dysfunction by modifying claudin-2 levels. *Biology*, 10 (3):205, 2021. DOI: 10.3390/biology10030205

67. Feng W, Wu Y, Chen G, Fu S, Li B, Huang B, Wang D, Wang W, Liu J: Sodium butyrate attenuates diarrhea in weaned piglets and promotes tight junction protein expression in colon in a GPR109A-dependent manner. *Cell Physiol Biochem*, 47 (4): 1617-1629, 2018. DOI: 10.1159/000490981
68. Huang C, Song P, Fan P, Hou C, Thacker P, Ma X: Dietary sodium butyrate decreases postweaning diarrhea by modulating intestinal permeability and changing the bacterial communities in weaned piglets. *J Nutr*, 145 (12): 2774-2780, 2015. DOI: 10.3945/jn.115.217406
69. Macia L, Tan J, Vieira AT, Leach K, Stanley D, Luong S, Maruya M, Ian McKenzie C, Hijikata A, Wong C, Binge L, Thorburn AN, Chevalier N, Ang C, Marino E, Robert R, Offermanns S, Teixeira MM, Moore RJ, Flavell RA, Fagarasan S, Mackay CR: Metabolite-sensing receptors GPR43 and GPR109A facilitate dietary fibre-induced gut homeostasis through regulation of the inflammasome. *Nat Commun*, 6:6734, 2015. DOI: 10.1038/ncomms7734
70. Singh N, Gurav A, Sivaprakasam S, Brady E, Padia R, Shi H, Thangaraju M, Prasad PD, Manicassamy S, Munn DH, Lee JR, Offermanns S, Ganapathy V: Activation of Gpr109a, receptor for niacin and the commensal metabolite butyrate, suppresses colonic inflammation and carcinogenesis. *Immunity*, 40 (1): 128-139, 2014. DOI: 10.1016/j.immuni.2013.12.007
71. Ahmad S, Ali S, Ur Rehman A, Ul Haq I, Liaqat I, Aftab MN, Shume T: SCFAs' pleiotropic role in pathogenesis and salutogenesis: Mechanisms in exacerbation and regulation of inflammation and fibrosis from gut to host. *J Med Microbiol*, 75 (1):002117, 2026. DOI: 10.1099/jmm.0.002117
72. Zhang H, Sun J, Zheng X, Yang H, Xie A, Ding Y, Mei Y, Li J, Hu Y, Ren M, Liu Y, Liang Y: Fermented *Lactocaseibacillus paracasei* cultures ameliorate colitis by modulating microbiota-derived tryptophan metabolism and macrophage polarization. *Adv Sci*, 13 (13):e13920, 2026. DOI: 10.1002/advs.202513920
73. Bernardi F, D'Amico F, Bencardino S, Faggiani I, Fanizza J, Zilli A, Parigi TL, Allocca M, Danese S, Furfaro F: Gut microbiota metabolites: Unveiling their role in inflammatory bowel diseases and fibrosis. *Pharmaceutics*, 17 (3):347, 2024. DOI: 10.3390/ph17030347
74. Turner JR: Intestinal mucosal barrier function in health and disease. *Nat Rev Immunol*, 9 (11): 799-809, 2009. DOI: 10.1038/nri2653
75. Liu G, Gu K, Wang F, Jia G, Zhao H, Chen X, Wu C, Zhang R, Tian G, Cai J, Tang J, Wang J: Tryptophan ameliorates barrier integrity and alleviates the inflammatory response to enterotoxigenic *Escherichia coli* K88 through the CaSR/Rac1/PLC- γ 1 signaling pathway in porcine intestinal epithelial cells. *Front Immunol*, 12:748497, 2021. DOI: 10.3389/fimmu.2021.748497
76. López-Colom P, Castillejos L, Rodríguez-Sorrento A, Puyalto M, Mallo JJ, Martín-Orúe SM: Impact of in-feed sodium butyrate or sodium heptanoate protected with medium-chain fatty acids on gut health in weaned piglets challenged with *Escherichia coli* F4. *Arch Anim Nutr*, 74 (4): 271-295, 2020. DOI: 10.1080/1745039X.2020.1726719
77. Neurath MF: Strategies for targeting cytokines in inflammatory bowel disease. *Nat Rev Immunol*, 24 (8): 559-576, 2024. DOI: 10.1038/s41577-024-01008-6
78. Heckmann M, Sadova N, Sandner G, Neuhauser C, Blank-Landeshammer B, Schwarzingen B, König A, Liang M, Spitzer M, Weghuber J, Stadlbauer V: Herbal extract fermented with inherent microbiota improves intestinal health by exerting antioxidant and anti-inflammatory effects *in vitro* and *in vivo*. *J Anim Sci Biotechnol*, 16 (1):52, 2025. DOI: 10.1186/s40104-025-01178-w
79. Li Z, Chen L, Huang Z, Jia G, Zhao H, Liu G, Chen X: Supplementation with L-theanine promotes intestinal antioxidant ability via Nrf2 signaling pathway in weaning piglets and H₂O₂-induced IPEC-J2 cells. *J Funct Foods*, 121:106433, 2024. DOI: 10.1016/j.jff.2024.106433
80. Wen X, Wan F, Wu Y, Liu L, Liu Y, Zhong R, Chen L, Zhang H: Caffeic acid supplementation ameliorates intestinal injury by modulating intestinal microbiota in LPS-challenged piglets. *Food Funct*, 14 (16): 7705-7717, 2023. DOI: 10.1039/d3fo02286b
81. Heneghan AF, Pierre JF, Kudsk KA: JAK-STAT and intestinal mucosal immunology. *JAK STAT*, 2 (4):e25530, 2013. DOI: 10.4161/jkst.25530
82. Cao DL, Woo MS, Kim EJ, Ahn B, Prayoga AH, Kang SS, Cho KM, Kang D: Anti-inflammatory effects of fermented and aged mountain-cultivated ginseng sprouts via suppression of MAPK-NF- κ B pathway in lipopolysaccharide-stimulated RAW264.7 macrophages. *Food Sci Nutr*, 12 (11): 9566-9576, 2024. DOI: 10.1002/fsn.3.4518
83. He F, Ru X, Wen T: Nrf2, a transcription factor for stress response and beyond. *Int J Mol Sci*, 21 (13):4777, 2020. DOI: 10.3390/ijms21134777
84. Ngo V, Duennwald ML: Nrf2 and oxidative stress: A general overview of mechanisms and implications in human disease. *Antioxidants*, 11 (12):2345, 2022. DOI: 10.3390/antiox11122345
85. Khan MZ, Li S, Ullah A, Li Y, Abohashrh M, Alzahrani FM, Alzahrani KJ, Alsharif KF, Wang C, Ma Q: Therapeutic agents targeting the Nrf2 signaling pathway to combat oxidative stress and intestinal inflammation in veterinary and translational medicine. *Vet Sci*, 13 (1):25, 2025. DOI: 10.3390/vetsci13010025
86. Chen G, Li Z, Liu S, Tang T, Chen Q, Yan Z, Peng J, Yang Z, Zhang G, Liu Y, Zheng M: Fermented chinese herbal medicine promoted growth performance, intestinal health, and regulated bacterial microbiota of weaned piglets. *Animals*, 13 (3):476, 2023. DOI: 10.3390/ani13030476
87. Kobayashi EH, Suzuki T, Funayama R, Nagashima T, Hayashi M, Sekine H, Tanaka N, Moriguchi T, Motohashi H, Nakayama K, Yamamoto M: Nrf2 suppresses macrophage inflammatory response by blocking proinflammatory cytokine transcription. *Nat Commun*, 7:11624, 2016. DOI: 10.1038/ncomms11624
88. Gao W, Guo L, Yang Y, Wang Y, Xia S, Gong H, Zhang BK, Yan M: Dissecting the crosstalk between Nrf2 and NF- κ B response pathways in drug-induced toxicity. *Front Cell Dev Biol*, 9:809952, 2022. DOI: 10.3389/fcell.2021.809952
89. Metzler-Zebeli BU, Nöllenburg N, Koger S, Schobersberger K, Vötterl J, Leeb C: Potential of fermented herbal extracts to modulate digestion and gut microbiota during the weaner and fattening period on commercial pig farms. *Front Vet Sci*, 12:1620045, 2025. DOI: 10.3389/fvets.2025.1620045
90. Katu JK, Tóth T, Ásványi B, Hatvan Z, Varga L: Effect of fermented feed on growth performance and gut health of broilers: A review. *Animals*, 15 (13):1957, 2025. DOI: 10.3390/ani15131957
91. Yin D, Du E, Yuan J, Gao J, Wang Y, Aggrey SE, Guo Y: Supplemental thymol and carvacrol increases ileum *Lactobacillus* population and reduces effect of necrotic enteritis caused by *Clostridium perfringens* in chickens. *Sci Rep*, 7 (1):7334, 2017. DOI: 10.1038/s41598-017-07420-4
92. Du E, Wang W, Gan L, Li Z, Guo S, Guo Y: Effects of thymol and carvacrol supplementation on intestinal integrity and immune responses of broiler chickens challenged with *Clostridium perfringens*. *J Anim Sci Biotechnol*, 7:19, 2016. DOI: 10.1186/s40104-016-0079-7
93. Künili İ E, Akdeniz V, Akpınar A, Budak Ş, Curiel JA, Guzel M, Karagözü C, Berkel Kasicki M, Caruana GPM, Starowicz M, Humblot C, Keyvan E, Chassard C, Pracer S, Vergères G, Kesenkaş H: Bioactive compounds in fermented foods: A systematic narrative review. *Front Nutr*, 12:1625816, 2025. DOI: 10.3389/fnut.2025.1625816
94. Gresse R, Chaucheyras-Durand F, Fleury MA, Van de Wiele T, Forano E, Blanquet-Diot S: Gut microbiota dysbiosis in postweaning piglets: Understanding the keys to health. *Trends Microbiol*, 25 (10): 851-873, 2017. DOI: 10.1016/j.tim.2017.05.004
95. Su J, Zhu Q, Zhao Y, Han L, Yin Y, Blachier F, Wang Z, Kong X: Dietary supplementation with chinese herbal residues or their fermented products modifies the colonic microbiota, bacterial metabolites, and expression of genes related to colon barrier function in weaned piglets. *Front Microbiol*, 9:3181, 2018. DOI: 10.3389/fmicb.2018.03181
96. Naeem M: Combating necrotic enteritis in poultry through nutritional interventions: A review of alternatives to antibiotics. *Anim Dis*, 6 (1):16, 2026. DOI: 10.1186/s44149-026-00225-x
97. Zhang X, Miao Q, Pan C, Yin J, Wang L, Qu L, Yin Y, Wei Y: Research advances in probiotic fermentation of Chinese herbal medicines. *iMeta*, 2 (2):e93, 2023. DOI: 10.1002/imt2.93

98. Gao C, Zhou Y, Chen Z, Li H, Xiao Y, Hao W, Zhu Y, Vong CT, Farag MA, Wang Y, Wang S: Turmeric-derived nanovesicles as novel nanobiologics for targeted therapy of ulcerative colitis. *Theranostics*, 12 (12): 5596-5614, 2022. DOI: 10.7150/thno.73650
99. Bose S, Han KW, Lee MJ, Kim H: Intestinal protective effects of herbal-based formulations in rats against neomycin insult. *Evid Based Complement Alternat Med*, 2013:161278, 2013. DOI: 10.1155/2013/161278
100. Han S, Xu G, Zhang K, Ahmad S, Wang L, Chen F, Liu J, Gu X, Li J, Zhang J: Fermented *Astragalus* powder, a new potential feed additive for broilers to improve the growth performance and health. *Animals*, 14 (11):1628, 2024. DOI: 10.3390/ani14111628
101. Xiao R, Wang L, Tang Z, Qian X, Wang J, Lian Y, Tang J, Xu J, Lin Y, Shi B, Xu P, Xiong Q: Effects of fermented Chinese herbal medicine feed additives on growth performance and intestinal microbiota of piglets. *PLoS One*, 19 (10):e0308196, 2024. DOI: 10.1371/journal.pone.0308196
102. Fan J, Cui H, Mu Z, Yao C, Yang M, Jin Y, Ning C, Zhang H: Non-targeted metabolomics analysis of fermented traditional Chinese medicine and its impact on growth performance, serum biochemistry, and intestinal microbiome of weaned lambs. *Sci Rep*, 14 (1):20385, 2024. DOI: 10.1038/s41598-024-71516-x
103. Li C, Xi Y, Li S, Zhao Q, Cheng W, Wang Z, Zhong J, Niu X, Chen G: Berberine ameliorates TNBS induced colitis by inhibiting inflammatory responses and Th1/Th17 differentiation. *Mol Immunol*, 67 (2 Pt B): 444-454, 2015. DOI: 10.1016/j.molimm.2015.07.013
104. Vesci L, Tundo G, Soldi S, Galletti S, Stoppoloni D, Bernardini R, Modolea AB, Luberto L, Marra E, Giorgi F, Marini S: A novel *Lactobacillus brevis* fermented with a vegetable substrate (AL0035) counteracts TNBS-induced colitis by modulating the gut microbiota composition and intestinal barrier. *Nutrients*, 16 (7):937, 2024. DOI: 10.3390/nu16070937
105. Wang Y, Xie Q, Sun S, Huang B, Zhang Y, Xu Y, Zhang S, Xiang H: Probiotics-fermented *Massa Medicata Fermentata* ameliorates weaning stress in piglets related to improving intestinal homeostasis. *Appl Microbiol Biotechnol*, 102 (24): 10713-10727, 2018. DOI: 10.1007/s00253-018-9438-y
106. Strober W, Fuss IJ, Blumberg RS: The immunology of mucosal models of inflammation. *Annu Rev Immunol*, 20, 495-549, 2002. DOI: 10.1146/annurev.immunol.20.100301.064816
107. Cui R, Zhang C, Pan ZH, Hu TG, Wu H: Probiotic-fermented edible herbs as functional foods: A review of current status, challenges, and strategies. *Compr Rev Food Sci Food Saf*, 23 (2):e13305, 2024. DOI: 10.1111/1541-4337.13305
108. Xu R, Kiarie EG, Yiannikouris A, Sun L, Karrow NA: Nutritional impact of mycotoxins in food animal production and strategies for mitigation. *J Anim Sci Biotechnol*, 13 (1):69, 2022. DOI: 10.1186/s40104-022-00714-2
109. Yang W, Zhang Y, Wu W, Huang L, Guo D, Liu C: Approaches to establish Q-markers for the quality standards of traditional Chinese medicines. *Acta Pharm Sin B*, 7 (4): 439-446, 2017. DOI: 10.1016/j.apsb.2017.04.012
110. Lin C, Zhou Z, Zhang L, Wang H, Lu J, Wang X, An R: Gegen Qinlian Decoction relieves ulcerative colitis via adjusting dysregulated Nrf2/ARE signaling. *Evid Based Complement Alternat Med*, 2022:2934552, 2022. DOI: 10.1155/2022/2934552