

Investigations of *Listeria* Species in Milk and Silage Produced in Burdur Province ^[1]

Fulya TAŞÇI *  Hülya TÜRÜTOĞLU ** Hatice ÖĞÜTÇÜ ***

[1] This study was supported by the Scientific Research Projects Fund of the Akdeniz University (Project No: 2005.01.0108.001)

* Mehmet Akif Ersoy University, Faculty of Veterinary Medicine, Department of Food Hygiene and Technology, TR-15100 Burdur - TURKEY

** Mehmet Akif Ersoy University, Faculty of Veterinary Medicine, Department of Microbiology, TR-15100 Burdur - TURKEY

*** Ahi Evran University, Faculty of Sciences and Arts, Department of Biology, TR-40100 Kırşehir - TURKEY

Makale Kodu (Article Code): KVFD-2010-1744

Summary

The aim of this study was to investigate the presence of *Listeria* species in the milk and silage samples obtained from fifteen different farms in Burdur. A total of 250 samples (silage and cow's milk obtained from animals fed and not fed with silage) were analyzed. *L. monocytogenes* was isolated in 6 (2.4%) out of the 250 samples. Five (6.66%) of the 75 silage samples and 1 (1.17%) of the 85 milk samples obtained from cows fed with silage were contaminated with *L. monocytogenes*, whereas no *Listeria* spp. were isolated from the 90 milk samples from cows not fed with silage. The isolation of *L. monocytogenes* from milk and silage samples in Burdur indicates that these products could create a serious risk to the public health.

Keywords: Cow's milk, Silage, *Listeria* spp.

Burdur Yöresinde Üretilen Süt ve Silajlarda *Listeria* Türlerinin Araştırılması

Özet

Bu çalışmada, Burdur yöresinde onbeş farklı çiftlikte üretilen süt ve silajlarda *Listeria* türlerinin varlığının araştırılması amaçlanmıştır. Toplam 250 örnek (silaj ile beslenen ve beslenmeyen inek sütleri ve silaj) analize alınmıştır. İki yüz elli örneğin 6'sında (%2.4) *L. monocytogenes* izole edilmiştir. Yetmiş beş silajın 5 (%6.66)'inde, 85 silaj verilen inek sütünün 1 (%1.17)'inde *L. monocytogenes* olduğu belirlenmiştir. Silaj ile beslenmeyen 90 inekten alınan sütlerde ise *Listeria* spp. izole edilememiştir. Sonuç olarak, Burdur'da üretilen süt ve silajlarda *L. monocytogenes*'in izole edilmesi bu ürünlerin halk sağlığı açısından bir risk oluşturabileceğini göstermektedir.

Anahtar sözcükler: İnek sütü, Silaj, *Listeria* spp.

INTRODUCTION

Listeria spp. are widely distributed in nature and found in soil, silage, decaying vegetation, animal feces, sewage water, and other environmental sources ¹. *Listeria monocytogenes* may contaminate milk because of mastitis, encephalitis, or abortion related to *Listeria* spp. in animals ^{1,2}. Listeriosis is a severe and often fatal illness

with clinical manifestations such as sepsis or meningitis in immunocompromised patients or neonatal babies and flu-like illness or abortion during pregnancy in women. The major outbreaks of listeriosis have been associated with the consumption of foods of animal origin ³. The genus *Listeria* contains 6 species: *L.*



İletişim (Correspondence)



+90 248 2344500



fulya_tasci22@hotmail.com

monocytogenes, *L. innocua*, *L. seeligeri*, *L. welshimeri*, *L. ivanovii*, and *L. grayi*⁴. In addition to *L. monocytogenes*, *L. seeligeri* and *L. ivanovii* may be pathogenic in humans^{1,5}. *L. monocytogenes* can frequently be isolated from unpasteurized milk and milk products⁶⁻¹⁰. Except in modern cheese production plants, raw milk is widely used in cheese production by small and medium domestic and commercial plants in Turkey. Also, isolation rate of *L. monocytogenes* in cheese samples in Turkey has been reported from 2% to 5%^{6-8,11}.

Silage is produced by harvesting a forage crop with a high moisture content (greater than 50%) and subsequently fermenting. In general, good silage remains stable, with no change in composition or heat, once air is eliminated and the silage has achieved a low pH¹². *Listeria* spp. are most commonly recovered from improperly fermented silage^{13,14}. It has been reported that listeriosis in cattle is mainly feed-borne¹ and *Listeria* spp. have been detected from 1.2% to 60% of the silage samples¹⁵⁻¹⁷. Also, *Listeria* spp. have been isolated from 2% to 6.1% milk samples from cows fed with silage¹⁷. In a study by Fenlon et al.¹⁸ has been stated that 29-31% of cattle started to shed *L. monocytogenes* after silage feeding.

The aim of this study was to investigate the presence of *Listeria* species in the milk and silage samples obtained from fifteen different farms in Burdur. The milk obtained from cows fed and not fed with silage were compared in terms of *Listeria*, and its importance in contamination of silage was put forward.

MATERIAL and METHODS

Sampling

Five research centers in Burdur were determined for sampling. Three different farms in every research center were visited every month between December 2007 and May 2008. In fifteen farms, seventy five silage samples, 85 milk samples obtained from cows fed with silage and 90 milk samples obtained from cows not fed with silage were collected. The samples were collected in sterile plastic bags and transported to the laboratory in boxes containing ice.

Isolation and Identification of *Listeria* spp.

All procedures were applied according to the FDA-Bacteriological Analytical Manual¹⁹. All media used were obtained from Oxoid (Oxoid Ltd., Hampshire, UK). Each sample (25 g/ml) was taken and placed in a stomacher bag to which 225 ml of sterile *Listeria* Selective Enrichment Broth (Oxoid) was added and homogenized

with a stomacher (Masticator, IUL Instruments-Spain) for 1-3 min and incubated at 30°C for 48 h. A loopful of homogenate was surface streaked in duplicate on Palcam agar (Oxoid) and Oxford agar (Oxoid). The Palcam plates were incubated at 37°C for 48 h under microaerophilic conditions and Oxford plates at 35°C for 48 h under aerobic conditions. All colonies surrounded by a brownish green and/or black halo were taken as possible *Listeria* spp. One suspected *Listeria* spp. colony from each plate was chosen and purified on tryptic soy agar (Oxoid CM 131) with 0.6% yeast extract (Oxoid L 21) and incubated at 30°C for 24-48 h for further biochemical characterization. Presumptive *Listeria* isolates were confirmed and identified at the species level based on Gram staining, typical umbrella motility in SIM medium (Oxoid CM 435), H₂S production, indole, urease, catalase, oxidase reaction, β-hemolysis, nitrate reduction, methyl-red/voges-proskauer (Oxoid CM 43), CAMP tests and fermentation of mannitol, L-rhamnose, D-ksilose, sorbitol, dextrose, maltose, esculin, dulcitol and salicin^{4,20,21}. Serotyping of isolates was performed with Bacto-*Listeria*-O-antisera types 1 and 4 and poly (Difco Laboratories, Detroit, MI) by the slide agglutination test^{4,21}.

Measurement of pH Values of the Samples

After the samples were collected for microbiologic analysis, the pH values of the milk samples were measured with an electronic pH meter (Metrohm 704 pH Meter). A 25-g aliquot silage sample was blended with 100 ml of deionized water for 2 min and filtered through four layers of cheesecloth. Then the pH of the extract was measured²².

Statistical Analysis: The results were analyzed using Minitab-15 with the chi-square analysis.

RESULTS

Overall, *L. monocytogenes* was found in 6 (2.4%) out of 250 samples. Five (6.66%) of the 75 silage samples and 1 (1.17%) of the 85 milk samples obtained from fed with silage were contaminated with *L. monocytogenes*, whereas no *Listeria* spp. were isolated from the 90 milk samples from cows not fed with silage. The differences between isolation rates of *L. monocytogenes* were statistically significant ($\chi^2=8.02$; $P=0.018$; $P<0.05$) (Table 1). Two selective plating media Palcam and Oxford were compared for isolating *L. monocytogenes* from the samples, and the isolation rates from these media were found to be equal.

In the present study, the pH values of the milk samples varied between 6.6 and 7.1, and the pH values

Table 1. The isolation rate of *L. monocytogenes* isolated from milk and silage^a**Tablo 1.** Süt ve silajlarda *L. monocytogenes* izolasyon oranı^a

Sample Type and Number (n)	<i>L. monocytogenes</i>	
	n	%
Silage (n: 75)	5	6.66
The milk of cows fed with silage (n: 85)	1	1.17
The milk of cows not fed with silage (n: 90)	-	-
Total (n: 250)	6	2.4

^a Chi-square statistic is significant, $\chi^2=8.02$; $P=0.018$; $P<0.05$

of the silage varied between 4.1 and 8.7. In the silage samples contaminated with *L. monocytogenes*, the pH values varied between 5.1 and 8.3, and the pH value of the milk sample contaminated with *L. monocytogenes* was 6.9.

As the collection period of the milk and silage samples was compared in isolation, the contamination of *L. monocytogenes* was found higher in March (3 silage samples) than in January (2 silage samples) and February (1 milk sample from the cows fed with silage).

In this study, for the serotype determination of 6 isolates defined as *L. monocytogenes* Difco Bacto O Antiserum type 1 and type 4, and type poly were used. The results were as follows: 5 isolates (1 milk and 4 silage samples) type poly and type 4, 1 isolate (1 milk sample) type poly.

DISCUSSION

The isolation rates of *Listeria* spp. in silage has been demonstrated in several studies carried out in Turkey and in other countries^{15-17,23,24}. In this study, *L. monocytogenes* was detected in 6.66% of the 75 silage samples. This percentage is lower than the results reported by Oliveira et al.²⁵ and Grønstøl²⁶, but similar to the 6.1% obtained by Vilar et al.¹⁷ In Turkey, Aslantaş and Yıldız²³ isolated *L. monocytogenes* from 1 of 11 silage samples. However, Şahin et al.²⁴ did not isolate *L. monocytogenes* from the silage, but isolated *L. welshimeri* and *L. grayi*. In this study, the low isolation rate of *L. monocytogenes* in silage may be accounted that high-quality silage is produced by mostly producers. However, in our study, silage samples contaminated with *L. monocytogenes* was obtained only from wet silage. In the illumination of this result, we could say that and rainy weather conditions are the cause of this result.

Many researchers have investigated *L. monocytogenes* contamination of milk^{6,9,11,23,27,28} and *Listeria* species have been detected from 0.40% to 10% of milk samples^{6,9,11,23,29}. In Turkey, the isolation rates from raw milk samples have been reported 0.45% in İstanbul⁷, 0.94% in Ankara²⁷, 1.20% in Van⁶, 3% in West Anatolia²⁸ and 5% in Ankara²⁹. In other countries, the reported isolation rates from bulk tank milk samples were 1.2% in Pennsylvania³⁰, 4.9% in Ireland³¹ and 6.5% in the United States¹⁰. The sources of *Listeria* spp. in raw milk have been reported to be fecal³² and environmental contamination during the milking, storage, and transport of infected cows on dairy farms, and poor silage quality³³.

In the present study, *Listeria* species were not found from cow's milk samples not fed silage. But, 1.17% of the milk samples obtained from cows fed with silage were contaminated with *L. monocytogenes*. However, Şahin et al.²⁴ have reported that *L. monocytogenes* was not isolated from the silage and milk samples of cows fed with silage, but *L. welshimeri* and *L. grayi* were isolated. Vilar et al.¹⁷ detected *Listeria* spp. in 33.7% of silage samples and in 16.3% of milk samples. Donnelly³⁴ observed that 8 of 44 Holstein cows fed *Listeria*-contaminated silage shed the organism in their milk. Furthermore, milk from these animals was free of *L. monocytogenes* one month after feeding of contaminated silage ceased.

In our study, two selective plating media Palcam and Oxford were compared for isolating of *L. monocytogenes* from the samples, and the isolation rates from these media were found to be equal, which is consistent with the reports by Art and Andre³⁵, Capita et al.³⁶ and Uysal and Anđ⁷.

L. monocytogenes has thirteen serotypes, but, only three serotypes-4b, 1/2a and 1/2b-are responsible for the majority of veterinary and human listeriosis cases³⁷. In this study, for the serotype determination of 6 isolates defined as *L. monocytogenes*, O Antiserum type 1 and type 4, and type poly were used. The results were as follows: 5 isolates (1 milk and 4 silage samples) type poly and type 4, 1 isolate (1 milk sample) type poly. Van Kessel et al.¹⁰ isolated *L. monocytogenes* from 56 (6.5%) of 861 bulk-tank milk samples, and serotyping of these isolates yielded 5 serotypes (1/2a, 1/2b, 3b, 4b, and 4c). Jayarao and Henning³⁸ reported isolating *L. monocytogenes* in 6 (4.6%) of 131 bulk-tank milk samples and all isolates of *L. monocytogenes* belonged to O antigen type 1.

Multiple studies have reported seasonal variations of *Listeria* spp. isolation, some report that contamination rates increase during the summer months³⁹, while others⁴⁰ reported increased rates during winter. Gaya et

al.⁴¹ found that raw caprine milk contamination by *Listeria* spp. was seasonal; the incidence in the autumn (9.33%) and winter (5.14%) samples was higher than the incidence in the spring (0.85%) and summer (0.85%) samples. Uraz and Yücel²⁷ isolated 1 of the *L. monocytogenes* in winter whereas the other one was isolated in the spring season. Two factors may explain the increased isolation rate during March in our study: (1) March is usually very rainy in Burdur, and water is moisturised silage. Therefore, the quality silage is changed. (2) Seasonal differences in the incidence of *Listeria* spp. in raw milk may also be related to breeding practices. Dairy cattle typically bear their young in late winter or early spring. During winter gestation, dairy cattle develop a weakened immune system as a direct result of pregnancy, which, in turn, makes these animals more susceptible to listerial infections and abortions⁴².

The pH values of the silage samples from which *Listeria* spp. were isolated ranged from 5.1 to 8.3. Different range from those observed in other studies were 3.8 to 5.2 in Rea et al.³¹, 5.78 to 5.89 in Ryser et al.¹⁶, and 4.47 to 6.97 in Vilar et al.¹⁷. A variety of studies have confirmed that *L. monocytogenes* contamination is most frequently associated with poor-quality silage¹⁷. Poorly fermented silage, which has a pH greater than 5.5, is ideal for *Listeria* growth^{16,17}. However, Fensterbank et al.⁴³ identified *Listeria* spp, including *L. monocytogenes*, in 11 of 31 high-quality silage samples with pHs of 3.6 to 4.0. In our study, the pH value of the milk and silage samples contaminated with *L. monocytogenes* was greater than 6.6. We believe that the contamination sources of *Listeria* spp. are the consumption of bad-quality silage, subjected to inadequate fermentation, with pH values higher than 4.0, which allows the multiplication of *Listeria* spp.

As a conclusion, the isolation of *L. monocytogenes* from milk and silage samples in Burdur indicates that these products could create a serious risk to the public health and could have a potential risk for animals. Correct practices with respect to silage production and milking are essential for preventing introduction of *Listeria* into the herd, its spread within the herd, and its entry into milk. The risk of contamination of milk by *Listeria* spp. increased when animals were fed low-quality silage, notably silage with pH \geq 4.5. Although the contamination ratio is very low in this research, *Listeria* contamination must be obstructed or minimized to achieve standard conditions.

REFERENCES

1. Jay JM: Modern Food Microbiology. 6th ed. Aspen Publishers, Inc, Gaithersburg, 2000.

2. Öcal N, Babür C, Yağcı BB, Macun HC, Çelebi B, Kılıç S, Yağcı İP: Kırıkkale yöresinde süt sığırlarında brusellozis, listeriozis ve toksoplazmozis'in seroprevalansı ve birlikte görülme sıklığı. *Kafkas Univ Vet Fak Derg*, 14 (1): 75-81, 2008.

3. Iida T, Kanzaki M, Nakama A, Kokubo Y, Maruyama T, Kaneuchi C: Detection of *Listeria monocytogenes* in human, animals and foods. *J Vet Med Sci*, 60, 1341-1343, 1998.

4. Anonymous: Bacteriological Analytical Manual Online. FDA, Revision AOAC International, USA 2003. <http://www.cfsan.fda.gov/~ebam/bam-10.htm>. Accessed: 30.04.2009.

5. Wiedmann M: An integrated science based approach to dairy food safety. *Listeria monocytogenes* as a model system. *J Dairy Sci*, 86, 1865-1875, 2003.

6. Sağun E, Sancak YC, İşleyici O, Ekici K: Van ve çevresi süt ve otlı peynirlerinde *Listeria* türlerinin varlığı ve yaygınlığı üzerine bir araştırma. *Türk J Vet Anim Sci*, 25, 15-19, 2001.

7. Uysal HK, Anğ O: Süt ve süt ürünlerinden izole edilen *Listeria* türleri. *Türk Mikrobiyol Cem Derg*, 33, 163-169, 2003.

8. Gülmez M, Güven A: Beyaz ve çeçil peynirlerinde *Campylobacter*, *Salmonella* ve *Listeria* türlerinin araştırılması. *Kafkas Univ Vet Fak Derg*, 7 (2): 155-161, 2001.

9. Dhanashree B, Otta SK, Karunasagar I, Goebel W, Karunasagar I: Incidence of *Listeria* spp. in clinical and food samples in Mangalore, India. *Food Microbiol*, 20, 447-453, 2003.

10. Van Kessel JS, Karns JS, Gorski L, McCluskey BJ, Perdue ML: Prevalence of Salmonellae, *Listeria monocytogenes*, and fecal coliforms in bulk tank milk on US dairies. *J Dairy Sci*, 87, 2822-2830, 2004.

11. Aygün O, Pehlivanlar S: *Listeria* spp. in the raw milk and dairy products in Antakya, Turkey. *Food Control*, 17 (8): 676-679, 2006.

12. McDonald P, Henderson AR, Heron SJE: The Biochemistry of Silage. Chalcombe Publications, Marlow, England, 1991.

13. Grant MA, Eklund CA, Shields SC: Monitoring dairy silage for five bacterial groups with potential for human pathogenesis. *J Food Protect*, 58, 879-883, 1995.

14. Gazyacı S, Yıldırım M, Babür C, Kılıç S: Ankara yöresindeki koç ve koyunlarda *Listeria monocytogenes*'e karşı oluşan antikorların varlığının araştırılması. *Kafkas Univ Vet Fak Derg*, 15 (6): 975-977, 2009.

15. Fenlon DR: Rapid quantitative assessment of the distribution of *Listeria* in silage implicated in a suspected outbreak of listeriosis in calves. *Vet Rec*, 118, 240-242, 1986.

16. Ryser E, Arimi SM, Donnelly CW: Effects of pH on distribution of *Listeria* ribotypes in corn, hay, and grass silage. *Appl Environ Microb*, 63 (9): 3695-3697, 1997.

17. Vilar MJ, Yus E, Sanjuán ML, Diéguez JL, Rodríguez-Otero FJ: Prevalence of and risk factors for *Listeria* species on dairy farms. *J Dairy Sci*, 90, 5083-5088, 2007.

18. Fenlon DR, Wilson J, Donachie W: The incidence and level of *Listeria monocytogenes* contamination of food sources at primary production and initial processing. *J Appl Bacteriol*, 81, 641-650, 1996.

19. Hitchens AD: *Listeria monocytogenes*. In, FDA Bacteriological Analytical Manual. 8th ed. pp. 10.1-10.13, AOAC International, USA, 1995.

- 20. Andrews W:** Manual of Food Quality Control Microbiological Analysis. Washington, DC, USA: Food and Drug Administration, 1992.
- 21. Lovett J, Francis DW, Hunt JM:** *Listeria monocytogenes* in raw milk: Detection, incidence and pathogenicity. *J Food Protect*, 50, 188-192, 1987.
- 22. Hart SP, Horn FP:** Ensiling characteristics and digestibility of combinations of turnips and wheat straw. *J Anim Sci*, 14, 1790-1800, 1987.
- 23. Aslantaş O, Yıldız P:** Kars ilinde çiğ süttten *Listeria monocytogenes* izolasyonu. *Firat Univ Sag Bil Vet Derg*, 17 (1): 11-15, 2003.
- 24. Şahin K, Çerci IH, Güler T, Özcan C, Şahin N:** Silaj ve kuru ot katılan rasyonlarla beslenen süt ineklerinin kaba yem ve sütlerinde *Listeria* türlerinin araştırılması. *Firat Univ Sağlık Bil Vet Derg*, 10, 245-249, 1996.
- 25. Oliveira M, Guerra M, Bernardo F:** Occurrence of *Listeria monocytogenes* in silages assessed by fluorescent in situ hybridization. *Arq Bras Med Vet Zootec*, 60 (1): 267-269, 2008.
- 26. Grønstøl H:** Listeriosis in sheep: Isolation of *Listeria monocytogenes* from grass silage. *Acta Vet Scand*, 20, 492-497, 1979.
- 27. Uraz G, Yücel N:** *Listeria monocytogenes*, *Yersinia enterocolitica* ve benzerlerinin (*Yersinia kristensenii*, *Yersinia frederiksenii*, *Yersinia intermedia*) çiğ süt örneklerinden izolasyonu. *Turk J Agri Forestry*, 22, 463-468, 1998.
- 28. Soyutemiz E, Çetinkaya F, Özakın C, Gedikoğlu S:** Presence of *Listeria monocytogenes* in raw milk samples from West Anatolia. *Turk J Infect*, 15 (1): 5-9, 2001.
- 29. Erol İ, Şireli UT:** Occurrence and contamination levels of *Listeria* spp. in milk and dairy products in Ankara. *FEMS Symposium on the Versatility of Listeria species* (10-11 October, No. 44, pp. 44-45). İzmir, Turkey: Publication of The Turkish Microbiological Society, 2002.
- 30. Jayarao BM, Donaldson SC, Straley BA, Sawant AA, Hegde NV, Brown JL:** A survey of foodborne pathogens in bulk tank milk and raw milk consumption among farm families in Pennsylvania. *J Dairy Sci*, 89, 2451-2458, 2006.
- 31. Rea MC, Cogan TM, Tobin:** Incidence of pathogenic bacteria in raw milk in Ireland. *J Appl Bacteriol*, 73, 331-336, 1992.
- 32. Griffiths MW:** *Listeria monocytogenes*: Its importance in dairy industry. *J Sci Food Agri*, 47, 133-158, 1989.
- 33. Bemrah N, Sanaa M, Cassin MH, Griffiths MW, Cerf O:** Quantitative risk assessment of human listeriosis from consumption of soft cheese made from raw milk. *Prev Vet Med*, 37, 129-145, 1998.
- 34. Donnelly CW:** Listeriosis and dairy products: Why now and why milk? *Hoards Dairyman*, 131, 663-687, 1986.
- 35. Art D, Andre P:** Comparison of three selective isolation media for the detection of *L. monocytogenes* in foods. *Zbl Bakteriol*, 275, 79, 1991.
- 36. Capita R, Alonso Calleja C, Prieto M, Garcia Fernandez MC, Moreno B:** Comparison of PALCAM and modified Oxford plating media for isolation of *Listeria* species in poultry meat following UVM II or Fraser secondary enrichment broths. *Food Microbiol*, 18, 555-563, 2001.
- 37. Wesley IV, Borucki M, Call DR, Larson D, Schroeder-Tucker L:** Detection and diagnosis of *Listeria* and listeriosis. In, Torrence ME, Issacson RE (Eds): *Microbial Food Safety in Animal Agriculture-Current Topics*. pp. 233-241, Iowa State Press, Iowa, USA, 2003.
- 38. Jayarao BM, Henning DR:** Prevalence of foodborne pathogens in bulk tank milk. *J Dairy Sci*, 84, 2157-2162, 2001.
- 39. Farber JM, Sanders GW, Malcom SA:** The presence of *Listeria* spp. in raw milk in Ontario. *Can J Microbiol*, 34, 95-100, 1988.
- 40. Fernández-Garayzábal JF, Domínguez L, Vázquez JA, Gómez-Lucia A, Rodríguez-Ferri ER, Suárez G:** Occurrence of *Listeria monocytogenes* in raw milk. *Vet Rec*, 120, 258-259, 1987.
- 41. Gaya P, Saralegui C, Medina M, Nuñez M:** Occurrence of *Listeria monocytogenes* and other *Listeria* spp. in raw caprine milk. *J Dairy Sci*, 79, 1936-1941, 1996.
- 42. Ryser ET:** Incidence and behavior of *Listeria monocytogenes* in unfermented dairy products. In, Ryser ET, Marth EH (Eds): *Listeria, Listeriosis, and Food Safety*. 3rd ed., Chapter 11, pp. 357-404, CRC Press, Taylor & Francis Group, Boca Raton, 2007.
- 43. Fensterbank R, Audurier A, Godu J, Guerrault P, Malo N:** Study of *Listeria* strains isolated from sick animals and from the silage consumed. *Ann Rech Vet*, 15 (1): 113-118, 1984.