


Prevalence of Fungi in the Conjunctival Sac of Clinically Normal Sheep

Abdullah ARAGHI-SOOREH¹  Vahid HASSANPOUR²

¹ Department of Clinical Sciences, Faculty of Veterinary Medicine, Urmia Branch, Islamic Azad University, Urmia, IRAN
² Graduated in Veterinary Medicine, Faculty of Veterinary Medicine, Urmia Branch, Islamic Azad University, Urmia, IRAN

KVFD-2014-12550 Received: 03.11.2014 Accepted: 01.12.2014 Published Online: 09.12.2014

Abstract

Conjunctival swabs were obtained from both eyes of 50 healthy sheep to identify the fungal flora. Data were analyzed for effect of age and sex. Out of 100 samples cultured, 18 (18%) showed fungal growth with predominance of the genus *Cladosporium* (38.89%). Other isolated fungal genera were *Penicillium* (16.67%), *Rhodotorula* (16.67%), *Aspergillus* (16.67%) and *Curvularia* (11.10%). There was no significant effect of sex and age on frequency of fungal isolation. Results showed that fungi are not prevalent on the ocular surface of healthy sheep.

Keywords: Fungal flora, Eye, Sheep, *Cladosporium* spp.

Klinik Olarak Normal Koyunların Konjunktival Kesesinde Mantarların Yaygınlığı

Özet

Elli adet koyunda yapılan bu çalışmada hayvanların gözlerindeki konjunktival keselerden alınan sıvap örnekleme ile normal fungal floranın belirlenmesi amaçlandı. Çalışmada yaş ve cinsiyetin etkisi analiz edildi. Kültürü yapılan 100 örneğin 18 (%18)'inde fungal üreme görüldü. *Cladosporium* cinsi (%38.89) en çok üreyen mantar idi. Diğer üreyen mantarlar; *Penicillium* (%16.67), *Rhodotorula* (%16.67), *Aspergillus* (%16.67) ve *Curvularia* (%11.10) idi. Cinsiyet ve yaşın fungal izolasyon üzerine etkisi saptanmadı. Sonuçlar sağlıklı koyunların göz yüzeyinde mantarların yaygın olarak bulunmadığını gösterdi.

Anahtar sözcükler: Mantar florası, Göz, Koyun, *Cladosporium* spp.

INTRODUCTION

In many animal species fungi are considered as a part of the normal ocular flora, but their existence can turn to pathogenic state when defense mechanisms of the outer eye are damaged. Knowledge on the fungal species which are most likely encountered in the conjunctival sac is important to select an antifungal drug as initial empirical treatment of corneal mycoses^[1,2]. There is a paucity of information on the ocular fungal flora of ruminants^[3-5], especially sheep^[6]. Reportedly, *Cladosporium* spp, *Penicillium* spp and *Mucor* spp. are the predominant species. In order to increase knowledge of the ovine ocular fungal flora and to determine the effect of sex and age on the prevalence of isolates, we conducted present study on healthy sheep.

MATERIAL and METHODS

Fifty clinically healthy fat-tailed sheep presented at the

Urmia abattoir (Iran) from May to July 2011 were swabbed. Animals were from both sexes (26 males and 24 females) and divided into two age groups based on dental formula; A: Under 2 years old (34.68%) and B: above 2 years old (16.32%).

Samples were taken from the lower conjunctival sac of both eyes (n=100) using dry sterile swabs per eye, avoiding the eyelid margins and eyelashes. Swabs were placed into sterile tubes containing 2 ml normal saline and immediately transferred to the laboratory in a chilled box. Samples were plated onto Sabouraud dextrose agar (Merck, Darmstadt, Germany) and malt extract agar (Quelab, Montreal, Canada) and incubated at 25°C for 3 weeks. Identification of isolates was achieved to the genus level on the basis of macroscopic and microscopic features^[7].

The effect of age and sex on the frequency of fungal isolation was determined using Fischer's exact test. Significance was set at P<0.05.



İletişim (Correspondence)



+98 937 8819856; Fax: +98 443 2622043



a.araghi@iaurmia.ac.ir

RESULTS

Five genera of fungi were cultured from a total of 13 sheep (26%) and 18 eyes. The fungi isolated and the isolation rates are listed in [Table 1](#). *Cladosporium* spp (38.89%) was the most frequent isolate. Single fungi were isolated from each eye.

According to statistical analysis ([Table 2](#), [Table 3](#)), the prevalence of fungal isolates did not show significant differences between sexes and age groups ($P>0.05$).

Table 1. Fungal genera, frequency of species isolated and number of positive culture eyes and sheep

Tablo 1. Mantar türleri, izole edilen türlerin bulunma sıklığı ile göz ve koyunlarda pozitif kültür sayıları

Fungi	No. of Positive Sheep (%)	No. of Positive Eyes (%)	No. of Isolates (%)
<i>Cladosporium</i> spp.	6 (12)	7 (7)	7 (38.89)
<i>Penicillium</i> spp.	3 (6)	3 (3)	3 (16.67)
<i>Rhodotorula</i> spp.	3 (6)	3 (3)	3 (16.67)
<i>Aspergillus</i> spp.	3 (6)	3 (3)	3 (16.67)
<i>Curvularia</i> spp.	2 (4)	2 (2)	2 (11.10)

Table 2. Frequency analysis of the conjunctival fungal isolates in relation to sex of sheep

Tablo 2. Koyunların cinsiyet ile ilişkili olarak konjuktival mantar izolatlarının sıklık analizi

Fungi	No. of Isolates (%)		P-value
	Males n=26	Females n=24	
<i>Cladosporium</i> spp.	2	5	0.239
<i>Penicillium</i> spp.	2	1	0.999
<i>Rhodotorula</i> spp.	1	2	0.602
<i>Aspergillus</i> spp.	1	2	0.602
<i>Curvularia</i> spp.	2	-	0.491
Total	8	10	0.557

DISCUSSION

To our knowledge, this is the second published report of the ocular fungal flora in healthy sheep. In the current study, 26% of the sheep demonstrated positive cultures for fungi. This finding was consistent with those reported for goats (37.14%) [5], dogs (22%) and cats (40%) [3]. In the large animals this amount were reported to be 100% for cattle, 95% for horses [3], and 79.04% for donkeys [8]. It seems that results of the ocular fungal culture could be influenced by size of eye and its surface area exposed to the environmental fungi.

Table 3. Frequency analysis of the conjunctival fungal isolates in relation to age of sheep

Tablo 3. Koyunların yaş ile ilişkili olarak konjuktival mantar izolatlarının sıklık analizi

Fungi	No. of Isolates (%)		P-value
	<2 years old n=34	>2 years old n=16	
<i>Cladosporium</i> spp.	3	4	0.190
<i>Penicillium</i> spp.	2	1	0.999
<i>Rhodotorula</i> spp.	2	1	0.999
<i>Aspergillus</i> spp.	1	2	0.236
<i>Curvularia</i> spp.	1	1	0.542
Total	9	9	0.060

In the majority of studies on equine species *Aspergillus* spp is the most common isolate of the normal conjunctiva [2,8-10], but in domestic carnivores [3] and ruminants including cattle [3,4] and goats [5] generally *Cladosporium* spp and *Penicillium* spp were reported to be predominant. In our study, genus *Cladosporium* (38.89%) also reported as the most frequent isolates of conjunctiva. This finding was not similar to that reported for sheep in Italy [6], in which *Mucor* spp predominated (49%). It has been suggested that fungi are transitory inhabitants of the eye surface, and their prevalence can thus be influenced by geographic conditions [1,11].

In the present study there was no significant difference in isolation rates of fungi between sexes and age groups. In various studies have been showed that prevalence of the normal eye flora could be affected by these two host factors, as in male horses [1], male goats [5], female pigs [12] and younger horses [11] frequency of isolates were significantly higher.

Fungi are not prevalent on the ocular surface of healthy sheep. This could be considered as a reason for paucity of keratomycosis in this species.

REFERENCES

- Moore CP, Heller N., Majors LJ, Whitley D, Burgess EC, Weber J: Prevalence of ocular microorganisms in hospitalized and stabled horses. *Am J Vet Res*, 49, 773-777, 1988.
- Rosa M, Cardozo LM, Pereira JS, Brooks DE, Martins ALB, Florido PSS, Stussi GSP: Fungal flora of normal eyes of healthy horses from the State of Rio de Janeiro, Brazil. *Vet Ophthalmol*, 6, 51-55, 2003. DOI: 10.1046/j.1463-5224.2003.00267.x
- Samuelson DA, Andresen TL, Gwin RM: Conjunctival fungal flora in horses, cattle, dogs, and cats. *J Am Vet Med Assoc*, 184, 1240-1242, 1984.
- Sgorbini M, Barsotti G, Nardoni S, Brombin M, Sbrana A, Mancianti F, Corazza M: Seasonal prevalence of fungi in the conjunctival fornix of healthy cows during a 2-year study. *Vet Ophthalmol*, 13, 227-234, 2010. DOI: 10.1111/j.1463-5224.2010.00788.x
- Araghi-Sooreh A, Mokhber-Dezfuli MR, Mohammadi-Chorsi M: Identification of fungal isolates from the conjunctival sac of healthy goats. *J Vet Res*, 68, 337-332, 2013.

-
- 6. Bonelli F, Barsotti G, Attili AR, Mugnaini L, Cuteri V, Preziuso S, Corazza M, Preziuso G, Sgorbini M:** Conjunctival bacterial and fungal flora in clinically normal sheep. *Vet Rec Open*, 2014; 0:e000017. DOI: 10.1136/vropen-2013-000017.
- 7. Davise HL:** Medically Important Fungi: A Guide to Identification. 4th ed., ASM, New York, 2002.
- 8. Nardoni S, Sgorbini M, Barsotti G, Corazza M, Mancianti F:** Conjunctival fungal flora in healthy donkeys. *Vet Ophthalmol*, 10, 207-210, 2007. DOI: 10.1111/j.1463-5224.2007.00537.x
- 9. Barsotti G, Sgorbini M, Nardoni R, Corazza M, Mancianti F:** Occurrence of fungi from conjunctiva of healthy horses in Tuscany, Italy. *Vet Res Commun*, 30, 903-906, 2006. DOI: 10.1007/s11259-006-3366-5
- 10. Araghi-Sooreh A, Navidi M., Razi M:** Conjunctival bacterial and fungal isolates in clinically healthy working horses in Iran. *Kafkas Univ Vet Fak Derg*, 20, 625-627, 2014. DOI: 10.9775/kvfd.2013.10649
- 11. Andrew SE, Nguyen A, Jones GL, Brooks DE:** Seasonal effects on the aerobic bacterial and fungal conjunctival flora of normal thoroughbred brood mares in Florida. *Vet Ophthalmol*, 6, 45-50, 2003. DOI: 10.1046/j.1463-5224.2003.00265.x
- 12. Davidson HJ, Rogers DP, Yeary TJ, Stone GG, Schoneweis DA, Chengappa MM:** Conjunctival microbial flora of clinically normal pigs. *Am J Vet Res*, 55, 949-951, 1994.