

# Comparison of Three Fixation Methods for the Prevention of Wound Contractions in Diabetic and Non-Diabetic Mice with Full-Thickness Skin Excision <sup>[1]</sup>

Turgut ANUK <sup>1</sup>  Savaş ÖZTÜRK <sup>2</sup> İsa ÖZAYDIN <sup>2</sup> Şahin KAHRAMANCA <sup>3</sup>  
Sadık YAYLA <sup>2</sup> Özgür AKSOY <sup>2</sup> Zafer OKUMUŞ <sup>4</sup>  
Uğur AYDIN <sup>2</sup> Bülent Çağlar BİLGİN <sup>1</sup> İbrahim DEMİRKAN <sup>5</sup>

<sup>[1]</sup> Presented at 1<sup>st</sup> International Turkey Veterinary Surgery Congress (oral presentation), 11-14 May 2016, Erzurum, Turkey  
<sup>1</sup> Kafkas University, Faculty of Medicine, Department of General Surgery, TR-36100 Kars - TURKEY; <sup>2</sup> Kafkas University, Faculty of Veterinary Science, Department of Surgery, TR-36100 Kars - TURKEY; <sup>3</sup> Kars State Hospital, Department of General Surgery, TR-36300 Kars - TURKEY; <sup>4</sup> Atatürk University, Faculty of Veterinary Science, Department of Surgery, TR-25240 Erzurum - TURKEY; <sup>5</sup> Afyon Kocatepe University, Faculty of Veterinary Medicine, Department of Surgery, TR-03100 Afyon - TURKEY

Article Code: KVFD-2015-14837 Received: 14.12.2015 Accepted: 24.04.2016 Published Online: 25.04.2016

## Abstract

The most crucial problem in experimentally induced excisional wounds in animals with slim-skinned and poor subcutaneous attachment is wound contraction formed during recovery period. The contraction is mainly induced by myofibroblasts leading to early wound closure. This may be associated with false outcomes of the study. Recently, various trials have been made to reduce the contraction in full-thickness excisional models of wounds. This study aimed at comparing different fixation methods on the minimization of wound contraction. Study was carried out in two main groups as non-diabetic (Group I, n=24) and diabetic (Group II, n=24) and one control group. The animals in each group were divided into four equal subgroups (Group I-A, I-B, I-C, I-D; Group II-A, II-B, II-C, II-D) and 1.2x1.2 cm-sized full-thickness of skin excisions were generated on the dorsum of animals. No fixations were applied in subgroups I-A and II-A however, interrupted suture in subgroups I-B and II-B, suture passing from a couple of points through skin and subcutaneous tissues placed in each side of wound tied only in corners in subgroups I-C and II-C and using a polyethylene fixatives that fixed in wound edges with interrupted sutures in subgroups I-D and II-D were achieved to produce wound fixation methods. Digital photographs of the wound were taken on the 3<sup>rd</sup>, 7<sup>th</sup>, 14<sup>th</sup> and 21<sup>st</sup> days of the study, reduction ratios of the wounds were compared and the efficiency of methods for the prevention of contraction was examined. When wounds were evaluated comparatively after fixation, it was observed that three different applied fixation methods were efficient in prevention of wound contraction in all phases in comparison to control groups. Moreover, it was observed that contraction was highly impeded in interrupted suture groups (I-B, II-B) and polyethylene-reinforced interrupted suture groups (I-D, II-D) when compared to suture groups tied only in corners. The results revealed that wound surface area shrinkage should be taken into consideration in experimental studies to avoid misleading consequences.

**Keywords:** Wound, contraction, Full-thickness skin, Excision, Fixation, Mice

## Tam Katmanlı Deri Eksizyonu Oluşturulmuş Diyabetik ve Non-Diyabetik Farelerde Yara Kontraksiyonunun Önlenmesi İçin Üç Farklı Fiksasyon Yönteminin Karşılaştırılması

## Özet

İnce derili ve zayıf derialtı bağlantılı deney hayvanlarında oluşturulan eksizyonel yara modellerinde en önemli sorun iyileşme döneminde şekillenen yara kontraksiyonudur. Daha çok myofibroblastların neden olduğu bu durum yaranın erken kapanmasına ve yanıtıcı sonuçların ortaya çıkmasına neden olabilmektedir. Son zamanlarda, tam katmanlı eksizyonel yara modellerinde kontraksiyonu redukte etmek için çeşitli denemeler yapılmıştır. Bu çalışma ile yara kontraksiyonunun minimize edilmesine yönelik üç farklı fiksasyon yönteminin karşılaştırılması amaçlandı. Çalışma, non-diyabetik (Grup I, n=24) ve diyabetik (Grup II, n=24) olmak üzere 2 ana grupta yürütüldü. Her bir gruptaki hayvanlar 4 eşit alt gruba ayrılarak (Grup I-A, I-B, I-C, I-D; Grup II-A, II-B, II-C, II-D) hayvanların sırt bölgesinde 1.2x1.2 cm ölçülerinde tam katmanlı deri eksizyonu oluşturuldu. I-A ve II-A gruplarında herhangi bir fiksasyon uygulanmazken, I-B ve II-B gruplarında ayrı dikiş, I-C ve II-C gruplarında deri ve deri altı dokulardan birkaç noktadan geçecek şekilde her bir yara kenarına yerleştirilip sadece köşelerde düğümlenen dikiş, I-D ve II-D gruplarında ise polietilen bir fiksasyon materyalinin ayrı dikişlerle yara kenarına tespit edildiği fiksasyon modelleri oluşturuldu. Çalışmanın 3, 7, 14 ve 21. günlerinde yara bölgesinin dijital fotoğrafları alınarak yaraların küçülme oranları karşılaştırıldı ve kullanılan yöntemlerin kontraksiyonu engellemedeki etkinlikleri değerlendirildi. Fiksasyon sonrası yara alanları karşılaştırmalı olarak değerlendirildiğinde, uygulanan her üç fiksasyon yönteminin bütün aşamalarda kontrol gruplarına göre yara kontraksiyonunu engellemede etkili olduğu gözlemlendi. Ayrıca, tüm kontrol günlerinde ayrı dikiş (I-B, II-B) ve polietilen destekli ayrı dikiş (I-D, II-D) gruplarında, sadece köşelerde düğümlenen dikiş gruplarına göre kontraksiyonun daha fazla engellendiği de saptandı. Çalışmanın sonuçları, yara alanının küçülmesine dayalı deneylerde, biyolojik bir süreç olan yara kontraksiyonunun dikkate alınması ve yanıtıcı sonuçlardan kaçınmak için deney protokollerinin uygun bir fiksasyon eşliğinde yürütülmesi gerektiğini ortaya koymuştur.

**Anahtar sözcükler:** Yara kontraksiyonu, Tam katmanlı deri eksizyonu, Yara fiksasyonu, Fare



İletişim (Correspondence)



+90 532 6974498



turgutanuk@gmail.com

## INTRODUCTION

Wound healing can be defined as tissue regeneration of mechanic, cellular and biochemical process initiated after trauma. The process continues at three successive stages: hemostasis and inflammation phase, proliferation (granulation) phase and maturation phase [1,2]. In human and veterinary medicine, experimental studies on animal models are widely used as an important biological approach to clarify the pathophysiology of wound healing and to identify new strategies [3,4]. Even if different animal species are used for this purpose, mice and other rodents are extensively preferred because their care and feeding are easy, they reproduce and grow quickly, suitable for application of surgical procedures and their biology is thoroughly investigated [3,5]. On the other hand, an excessive skin contraction is a reality and edges of the wound move rapidly towards the centre. While the wound closes, the central portion undergoes keratinization; thus premature and bad wound healing occurs. This may be associated with that these animals' skin has thin and loosed skin, weak subcutaneous connection and little adipose tissue. It is known that the wound contraction accomplished by fibroblasts, particularly myofibroblasts is more effective in the first 2-8 weeks of wound healing [6]. Since the studies on the contraction-based early wound closure especially those investigating various medical drugs on wound healing may produce misleading and erroneous results. It has been advocated that the experiment regarding full-thickness excisional wounds should be carried out with skin fixation [7]. A few fixation materials and methods were reported to fix the edges of the wound [2,6-8]. Nevertheless, it is difficult to emphasize an ideal fixation method that can be used for all the wound models.

In this study, the efficiency of skin fixation simplified from previous complex models in the prevention of wound contraction after full-thickness excisional skin wounds were evaluated in mouse model. Also, to determine whether diabetes affects wound contraction, the study was carried

out in two groups as a diabetic and a non-diabetic. The wound area and the wound area reduction rate were digitally measured.

## MATERIAL and METHODS

### Ethical Approval

The study was performed after the approval by the local ethical committee (No: KAÜ-HADYEK 2014/060).

### Animals

A total of 48 male, Swiss Albino mice, aged 8-12 week were included in the study. Mice were homed in separate cages under standard laboratory conditions (12 h dark/12 h daylight, 45%-55% of humidity rate and room temperature at 20-22°C). Animals were fed with a standard feed and water was supplied as *ad libitum*.

### Study Groups


Animals were allocated into two main groups as non-diabetic (group I,  $n=24$ ) and diabetic (group II,  $n=24$ ). Each group were further divided into four equal subgroups (groups I-A, I-B, I-C, I-D; II-A, II-B, II-C, II-D). The following procedures were applied;

- *Control groups (I-A and II-A):* Skin was incised but sutures were not applied (Fig. 1-A),

- *Simple interrupted suture groups (I-B and II-B):* Wound edges were fixed using 12 sutures in equal numbers for each wound side after skin excision (Fig. 1-B),

- *Groups tied of sutures in the corners of wound (I-C and II-C):* Sutures were tied in the corners of wound by firmly stretching the knots (Fig. 1-C).

- *Polyethylene-supported suture groups (I-D and II-D):* Wound was fixed with 12 sutures using a polyethylene material made from a fluid infusion bag according to the wound size (Fig. 1-D).



**Fig 1.** Excision and fixation models. A- Excised wound (control groups), B- Excision + simple interrupted suture, C- Excision + suture tied in the corners of the wound, D- Excision + polyethylene-supported simple interrupted suture  
**Şekil 1.** Eksizyon ve fiksasyon modelleri. A- Yara eksizyonu (kontrol grupları), B- Eksizyon + basit ayrı dikiş, C- Eksizyon + yara köşelerinde düğümlenen dikiş, D- Eksizyon + polietilen materyal destekli basit ayrı dikiş

### Induction of Diabetes

To produce diabetic animals, freshly prepared streptozotocin in 0.1M sodium citrate buffer solution (pH 4.5) [9] was intraperitoneally administered in a single dose (200 mg/kg). On the 3<sup>rd</sup> day of injection, preprandial blood glucose level was measured by application of a capillary tube to medial eye canthus. Mice showing over 250 mg/mL (13.9 mMol/dL) level were considered as diabetic [9,10].

### Excisional Wound and Sutures

General anesthesia was induced by injecting the mixture of 10 mg/kg xylazine HCl (Rompun, 2%, Bayer) and 100 mg/kg ketamine HCl (Ketalar, 50 mg/ml, Pfizer) intraperitoneally. Dorsal regions of animals were clipped and disinfected with povidone iodine + 70% ethanol. Then, square shaped skin incision in size of 12x12 mm was made by a surgical knife. Skin and adipose tissues were dissected with the help of forceps and surgical scissors creating full-thickness skin excision (Fig. 1). In control subgroups, no surgical intervention was applied (subgroups I-A and II-A) (Fig. 1-A). In simple interrupted suture subgroups (I-B and II-B), 12 sutures were placed on wound edges in 3 mm apart from incision line. Skin was fixed to subcutaneous tissues and *panniculus carnosus* (Fig. 1-B). In suture tied in the corners of wound subgroups (I-C and II-C; Fig. 1-C), sutures located on each wound edge (in a way to pass through skin and subcutaneous tissues) were tied to their adjacent sutures stretched enough for secure fixation. In polyethylene-supported groups (I-D and II-D), a squarely-prepared polyethylene material was placed on skin after its central part was spread out more widely than the size of wound and then it was fixed with 12 simple interrupted sutures (Fig. 1-D). Thus, the polyethylene material generated more force for fixation by exerting pressure on the skin. In all suture groups, 4/0 polyglactin 910 (Vicryl - Ethicon) was used as suture material. Wounds were left open in all groups and any local and/or systemic treatment was not administered during the entire study.

### Wound Area Measurement and Wound Area Reduction Calculation

On the 0<sup>th</sup>, 3<sup>rd</sup>, 7<sup>th</sup>, 14<sup>th</sup> and 21<sup>st</sup> days of the study, wound area of all the subjects were logged on graph papers via a digital camera (Samsung WB2100). Digital image analyses were performed using a software program (Stereo Investigator 7.0; MBF Bioscience US). All image measurements were calibrated for each image with the help of graph paper therefore error probability originating from digital systems was eliminated. Wound areas were specified and they were measured with a 0.5 mm-sized point grid. Data were figured in mm<sup>2</sup>. Wound area reduction rate (%) was calculated by means of following equation [2]:

$$\frac{\text{Created wound area} - \text{Measured wound area}}{\text{Created wound area}} \times 100$$

### Statistical Analysis

Data generated in the study were evaluated by Anova, one-way test after being subjected to normality test (Minitab 17 Packed Programme). The significance level was set at  $P < 0.05$ .

## RESULTS

Measurements of wound sizes on the 0<sup>th</sup> day, and after excision and fixation days of 3, 7, 14 and 21 in non-diabetic and diabetic groups were given in Table 1. On the 3<sup>rd</sup> day after surgery showed that wound area in both non-diabetic and diabetic control groups decreased significantly ( $P < 0.05$ ) compared to fixation groups. In non-diabetic simple interrupted suture group the wound area was larger with respect to other fixation groups. While wound area in non-diabetic control group decreased significantly ( $P < 0.05$ ), on the 7<sup>th</sup> day, no significant difference was seen among fixation groups.

In diabetic group, wound areas in subgroups II-C and II-D were significantly larger than not only in II-A and II-B

**Table 1.** Non-diyabetik ve diyabetik deney gruplarında kontrol günlerinde saptanan yara yüzey alanı değerleri (ortalama± standart sapma)

**Table 1.** The values of wound area determined in non-diabetic and diabetic experimental groups on control days (mean± standard deviation)

Day	Wound Area (mm <sup>2</sup> )							
	Non-diabetic (n=24)				Diabetic (n=24)			
	I-A (n=6)	I-B (n=6)	I-C (n=6)	I-D (n=6)	II-A (n=6)	II-B (n=6)	II-C (n=6)	II-D (n=6)
0	143.8±1.8 <sup>a</sup>	144.0±1.4 <sup>a</sup>	143.5±0.9 <sup>a</sup>	144.01±0.9 <sup>a</sup>	143.08±1.2 <sup>a</sup>	144.01±1.4 <sup>a</sup>	143.07±1.6 <sup>a</sup>	144.27±2.2 <sup>a</sup>
3	54.31±4.8 <sup>a</sup>	103.66±11.3 <sup>b</sup>	77.69±6.7 <sup>c</sup>	83.63±7.4 <sup>bc</sup>	60.94±6.2 <sup>a</sup>	86.38±7.6 <sup>bc</sup>	89.94±5.2 <sup>bc</sup>	83.88±1.6 <sup>bc</sup>
7	29.31±2.2 <sup>a</sup>	43.06±3.8 <sup>b</sup>	41.94±3.8 <sup>b</sup>	40.42±4.6 <sup>b</sup>	40.56±4.6 <sup>b</sup>	45.38±5.3 <sup>b</sup>	61.94±4.9 <sup>c</sup>	68.25±5.6 <sup>c</sup>
14	3.25±0.75 <sup>a</sup>	11.88±7.0 <sup>b</sup>	15.22±7.1 <sup>b</sup>	11.93±0.5 <sup>b</sup>	4.50±2.00 <sup>a</sup>	14.81±4.7 <sup>b</sup>	20.06±1.3 <sup>c</sup>	31.25±2.3 <sup>d</sup>
21	1.43±1.5 <sup>a</sup>	3.68±0.5 <sup>a</sup>	2.81±1.4 <sup>a</sup>	3.43±1.5 <sup>a</sup>	3.00±1.4 <sup>a</sup>	7.50±1.0 <sup>b</sup>	3.44±1.4 <sup>a</sup>	16.06±1.4 <sup>c</sup>


<sup>a,b,c,d</sup> Different letters on each line show statistical differences ( $P < 0.05$ )

subgroups but also those in non-diabetic subgroups. Wound area reduction in non-diabetic control group showed significant difference in comparison to fixation groups on the 14<sup>th</sup> day as for the 7<sup>th</sup> day ( $P < 0.05$ ). The difference among in subgroups I-B, I-C and I-D were not significant, while wound area in diabetic control group decreased significantly in comparison to other diabetic subgroups at the same period. Wound area values in subgroup II-C were significantly larger with respect to subgroup II-B. Moreover wound area value in subgroup II-D was significantly larger than subgroups II-B and II-C ( $P < 0.05$ ). According to wound area measurements in non-diabetic and diabetic groups, on the 21<sup>st</sup> day, values in diabetic subgroups II-B and II-D were significantly different between in-groups and inter-groups (Table 1).

Reduction ratios in wound areas were presented respectively in Fig. 2 and Fig. 3 with respect to non-diabetic and diabetic groups. While the least reduction ratio in wound area was observed in subgroup I-B on the 3<sup>rd</sup> day. Reduction rate on the 7<sup>th</sup> and 14<sup>th</sup> day was higher only in control subgroup than other groups. However there was no significant difference between groups on the 21<sup>st</sup> day. Despite no significant difference in diabetic groups with regard to reduction rates of wound areas on the 3<sup>rd</sup> day, it was noted that subgroups II-C and II-D on the 7<sup>th</sup> and 14<sup>th</sup> day and group II-D on the 21<sup>st</sup> day showed less reduction in comparison to other groups.


## DISCUSSION

Since *in vitro* models remain insufficient to generate conclusive wound healing process the *in vivo* settings mandatory to be performed. Different aspects of wound healing and pathophysiology have to be understood at least for designing new therapeutic strategies. It is necessary to create models that incision, tissue loss, flap, graft and diabetes for rational approach towards different clinical problems. The use of experimental animals is highly crucial biological tools in human and veterinary medicine and it seems to remain reliable option as it was. Rodents are mostly chosen as experimental animals for acute and chronic wound model because their care and feeding are easy, they reproduce and grow quickly and their biological characteristics are well understood [3-5,11,12]. On the other hand, they create a significant problem during experimental studies in wound healing due to a loose connection between skin and subcutaneous tissues in these highly thin-skinned animals. Panniculus carnosus muscle is responsible for the wound contraction [2,6]. This biological event where myofibroblasts generally take an active role causing the wound to close earlier by pulling wound edges rapidly towards its centre [2]. For this reason, in full-thickness models of open wound in rodents, experimental protocols have to be implemented after providing skin fixation to achieve correct measurements. The adequate



**Fig 2.** Reduction rates of wound area in Group-I (non-diabetic) (%)  
**Şekil 2.** Group-I (non-diabetic) yara alanı küçülme oranları (%)

**Fig 3.** Reduction rates of wound area in Group-II (diabetic) (%)  
**Şekil 3.** Group-II (diabetic) yara alanı küçülme oranları (%)



and correct models are one of the fundamental aspects of studies on the formulation of therapeutic strategies [3,13]. This study, without consideration of any therapeutic goal, aimed at measuring how full-thickness excisional acute wounds in mice can be affected by no treatment or fixation by various options in terms of wound contraction. Since wound contraction generally progress together with inflammation (1-4 days) and proliferation (4-21 days) process, this was taken into consideration while controlling animals. Different methods have been described with the purpose of wound fixation in rodents. The basic approach of these studies is based on developing fixation materials that are glued or stitched to wound edges producing resistance to wound contraction. Metals (titanium, steel), splints prepared in shape of silicon-like circle or variants [6-8,14], adhesive materials such as Tegaderm, Duoderm, and polyvinylchloride are some of the materials used for this purpose. It was shown that some methods were successful but others inadequate for the prevention of contraction [2]. Therefore, it is difficult to explain an ideal fixation material and method for all wound management, now. The aims of the experiment or wound characteristics are all important aspects for determining fixation methods. An ideal experimental model of wound should sufficiently reflects new tissue formation process and re-epithelization associated with biology of wound healing however it should have a minimum effect on wound contraction. The cost of fixation method should be affordable, easily applicable and effective in minimization of contraction [8].

In experimental animal models, some coworkers prefer to study on auricle (rabbit), scalp and tail skin where wound contraction is minimum instead of skin and subcutaneous tissues of other parts of the body that create strong contraction [3,13]. Three different fixation methods used in this study were significantly effective in non-diabetic and diabetic models for reduction of contraction in comparison with corresponding control groups. These effects were observed more specifically on 3<sup>rd</sup>, 7<sup>th</sup> and 14<sup>th</sup> days of wound area measurements in particular, whereas any superiority was not identified between groups where suture and suture-supported polyethylene material fixation were used. Wound areas in diabetic subgroups where simple uninterrupted suture (II-B) and polyethylene material-supported suture (II-D) used failed to display considerable wound closure compared to other groups on the 21<sup>st</sup> day. The wound reduction rate in diabetic control group was almost similar to non-diabetic groups. This may suggest that the diabetes may not play significant role in wound

contraction. Hence, this condition may be largely explained that the wound healing is disruption due to diabetes but not wound contraction.

This study showed that three different employed fixation methods are considered as efficient in reduction of wound contraction in full-thickness excisional wound models in rodent. Thus, it was concluded that the wound fixation models presented here were simple and easy to apply for the determination of duration and degree of granulation formation, tissue proliferation and re-epithelialization for wound healing studies.

## REFERENCES

1. Orsted HL, Keast D, Forest-Lalande L, Marie Françoise M: Basic principles of wound healing. *Wound Care Canada*, 9 (2): 4-12, 2011.
2. Bae SH, Bae YC, Nam SB, Choi SJ: A skin fixation method for decreasing the influence of wound contraction on wound healing in a rat model. *Arch Plast Surg*, 39, 457-462, 2012. DOI: 10.5999/aps.2012.39.5.457
3. Ansell DM, Holden KA, Hardman MJ: Animal models of wound repair: Are they cutting it? *Exp Dermatol*, 21, 581-585. 2012. DOI: 10.1111/j.1600-0625.2012.01540.x
4. Perez R, Davis SC: Relevance of animal models for wound healing. *Wounds*, 20 (1): 3-8, 2008.
5. Dunn L, Prosser HC, Tan JT, Vanags LZ, Ng MK, Bursill CA: Murine model of wound healing. *J Vis Exp*, 28 (75): e50265, 2013. DOI: 10.3791/50265
6. Davidson JM, Yu F, Opalenik SR: Splinting strategies to overcome confounding wound contraction in experimental animal models. *Adv Wound Care (New Rochelle)*, 2, 142-148, 2013. DOI: 10.1089/wound.2012.0424
7. Carlson MA, Longaker MT, Thompson JS: Wound splinting regulates granulation tissue survival. *J Surg Res*, 110, 304-309, 2003. DOI: 10.1016/S0022-4804(02)00098-7
8. Ren L, Zhou B, Chen L: Silicone ring implantation in an excisional murine wound model. *Wounds*, 24 (2): 36-42, 2012.
9. Graham ML, Janecek JL, Kittredge JA, Hering BJ, Schuurman HJ: The streptozotocin-induced diabetic nude mouse model: Differences between animals from different sources. *Comp Med*, 61, 356-360, 2011.
10. Mendes JJ, Leandro CI, Bonaparte DP, Pinto AL: A rat model of diabetic wound infection for the evaluation of topical antimicrobial therapies. *Comp Med*, 62 (1): 37-48, 2012.
11. Gul Satar NY, Cangul IT, Topal A, Oktay A, Inan K, Akgul MB: Effects of ankaferd blood stopper (ABS) and topical tripeptide copper complex (TCC) on wound healing in rats: An experimental study. *Kafkas Univ Vet Fak Derg*, 20, 545-551, 2014. DOI: 10.9775/kvfd.2013.10555
12. Wong VW, Sorkin M, Glotzbach JP, Longaker MT, Gurtner GC: Surgical approaches to create murine models of human wound healing. *J Biomed Biotechnol*, 2011, Article ID: 969618, 2011. DOI: 10.1155/2011/969618
13. Davidson JM: Animal models for wound repair. *Arch Dermatol Res*, 290 (Suppl.): S1-S11, 1998.
14. Abercrombie M, James DW, Newcombe JF: Wound contraction in rabbit skin, studied by splinting the wound margins. *J Anat*, 94 (Pt 2): 170-182, 1960.