

Determination of the Stages of Deep Pectoral Myopathy Induced in Broilers Fed with Supplemental Coenzyme Q₁₀ ^[1]

Zlatko JOJKIC ¹ Slobodan STOJANOVIC ¹ Dragan ZIKIC ² Sinisa BJEDOV ²
Verica MILOSEVIC ³ Marko MILER ³ Gordana USCEBRKA ¹

^[1] This research has been financially supported by the Ministry of Education, Science and Technology Development of Republic of Serbia as a part of the Project No. TR 31033

¹ University of Novi Sad, Faculty of Agriculture, Department of Veterinary Medicine, Dositeja Obradovica 8, 21000 Novi Sad - SERBIA

² University of Novi Sad, Faculty of Agriculture, Department of Animal Science, Dositeja Obradovica 8, 21000 Novi Sad - SERBIA

³ University of Belgrade, Institute for Biological Research Sinisa Stankovic, Despota Stefana 142, 11000 Belgrade - SERBIA

Article Code: KVFD-2018-20103 Received: 11.05.2018 Accepted: 20.08.2018 Published Online: 20.08.2018

How to Cite This Article

Jojkic Z, Stojanovic S, Zikic D, Bjedov S, Milosevic V, Miler M, Uscebrka G: Determination of the stages of deep pectoral myopathy induced in broilers fed with supplemental coenzyme Q₁₀. *Kafkas Univ Vet Fak Derg*, 24 (6): 859-865, 2018. DOI: 10.9775/kvfd.2018.20103

Abstract

The aim of this study was to examine the structural characteristics and incidence of different stages of deep pectoral myopathy (DPM) that was induced in broilers fed a coenzyme Q₁₀ (CoQ₁₀) supplemented diet. A total of 288 1-day-old chicks (Cobb 500) were equally divided among 8 pens (pens 1 to 8). The diet was the same for all chicks until day 35 post-hatching. Subsequently, broilers in pens 5 to 8 were fed the 20 mg of CoQ₁₀/kg finisher diet until the end of the experiment (day 42 post-hatching). To induce DPM, 5 male birds from each of the pens 1 to 8 were subjected to encouraged wing flapping (EWF) at the end of their 37th day. At the end of the trial, the incidence of DPM stages in broilers was determined and an analysis of the histological parameters of deep pectoral muscles was performed. Results showed that, in the groups subjected to EWF, broilers with the CoQ₁₀ supplement had a lower average DPM stage and volume density of necrotic muscle cells, as well as a higher volume density of non-necrotic muscle cells. These results can be related to the antioxidant properties of CoQ₁₀, which, in chickens subjected to EWF, reduced the effects of DPM on cell necrosis and muscle tissue damage.

Keywords: Deep pectoral myopathy, Coenzyme Q₁₀, Broiler

İlave Koenzim Q₁₀ İle Beslenerek Derin Pektoral Miyopati Oluşturulan Broiler Piliçlerde Miyopatinin Devrelerinin Belirlenmesi

Öz

Bu çalışmanın amacı; koenzim Q₁₀ (CoQ₁₀) ilaveli diyet ile beslenerek derin pektoral miyopati (DPM) oluşturulan broiler piliçte meydana gelen miyopatinin farklı devrelerini ve yapısal özelliklerini araştırmaktır. Toplam 288 adet 1 günlük civciv (Cobb 500) 8 kafese eşit olarak dağıtıldı. Tüm civcivler için yumurtadan çıkmayı takiben 35. güne kadar diyet aynı tutuldu. Takibinde, 5'den 8'e kadar olan kafeslerdeki civcivler çalışmanın son gününe kadar (yumurtadan çıkmayı takiben 42. gün) 20 mg CoQ₁₀/kg bitirme diyeti ile beslendi. DPM oluşturmak amacıyla, 1'den 8'e kadar olan kafeslerin her birinde 5 erkek civciv 37. günlerinde kanat çırpmaya zorlandı. Deneme sonunda, broiler piliçlerdeki DPM devreleri belirlendi ve derin pektoral kaslarda histolojik parametreler analiz edildi. Elde edilen sonuçlar kanat çırpmaya zorlanan gruplarda CoQ₁₀ ilavesi ile beslenenlerde daha az DPM devresi ve daha az nekrotik kas hücresi hacim yoğunluğu ile daha fazla nekrotik olmayan kas hücresi yoğunluğunun oluştuğunu gösterdi. Bu sonuçlar CoQ₁₀'nın antioksidan özellikleri ile ilişkili olup, kanat çırpmaya zorlanan piliçlerde DPM'nin hücre nekrozu ve kas dokusu hasarı etkilerini azaltabilir.

Anahtar sözcükler: Derin pektoral miyopati, Koenzim Q₁₀, Broiler

INTRODUCTION

Several myopathies which occur in broilers, such as deep

pectoral myopathy (DPM), white striping (WS), and wooden breast (WB), are thought to be related to modern trends in poultry production that target increased growth rate,



İletişim (Correspondence)



+381 21 4853484



slobodan113@gmail.com

body weight and muscle mass of chickens. While WS is characterized by the occurrence of white striations [1], and WB by hardening of the breast muscle [2], DPM is associated with ischemic necrosis of the deep pectoral muscles [3].

The function of the deep pectoral muscle is related to wing activity [4]; its location near the sternum, the non-elastic fascia which surrounds the muscle, and the great size of the breasts in broiler chickens, leave insufficient space for the muscle to expand in response to wing movements. This leads to increased pressure within the muscle, blood flow obstruction and inadequate oxygen supply [5]. Tissue necrosis occurs as a result of ischemia. Described processes represent main characteristics of DPM.

Among several types of stresses (technological, environmental, nutritional, and internal) that can occur in poultry production [6], some of them caused by, for example, increased stocking density, catching or carrying the chickens by their wings, inappropriate weighing methods, etc., could result in unnecessary wing movements. This contributes to the development of more or less pronounced degrees of DPM [7]. In recent studies, methods defined as forced wing flapping [8] and encouraged wing flapping [9] have been used for the induction and examination of DPM in broilers.

Oxidative stress is related to elevated production of reactive oxygen species [10,11] that are associated with tissue necrosis [12]. Oxidative stress can be induced during hypoxia [13,14]. Considering that hypoxia is one of the characteristics of DPM (hypoxia can be induced through the occlusion of blood vessels and ischemia), it is therefore important to examine the influence of antioxidants on DPM development.

Beside the antioxidant properties of coenzyme Q₁₀ (CoQ₁₀) [15], it has an important role in oxidative phosphorylation and ATP synthesis [16]. CoQ₁₀ is already used as a supplement in broiler diets for the treatment of certain diseases and to improve immune functions. Its supplementation has been found to decrease mortality due to pulmonary hypertension syndrome in broilers exposed to low ambient temperature [17]. Furthermore, in research on Newcastle disease, chicks antibody titer was higher on day 21 post-hatching in broilers supplemented with 20 mg of CoQ₁₀/kg of diet [18].

The purpose of this study was to identify and examine the stages of DPM induced in broilers fed with supplemental CoQ₁₀.

MATERIAL and METHODS

Animals and Experimental Design

For the purpose of this study, 288 1-day-old chicks (Cobb 500) were randomly distributed between 8 floor pens

(pens 1 to 8), 36 chicks in each pen. Pens 1 to 4 represented 4 replicates, where chickens were fed with starter (days 1-14), grower (days 15-35) and finisher (days 36-42) diets (Table 1). Meanwhile pens 5 to 8 represented 4 replicates where feeding programs were the same as in pens 1 to 4, but with the additional supplementation of 20 mg of CoQ₁₀/kg finisher diet (days 36-42). To induce DPM, 5 male birds from each of the pens 1 to 4, as well as from each of the pens 5 to 8, with a body weight nearest to the mean body weight of the male birds, were subjected to encouraged wing flapping (EWF) at the end of day 37 post-hatching.

The chickens of the control group were fed a basal diet, while the 3 treatments included: 1) chickens fed the basal diet and subjected to EWF; 2) chickens fed a basal diet with CoQ₁₀ supplementation; and 3) chickens fed a basal diet supplemented with CoQ₁₀ that were subjected to EWF. Food and water were available *ad libitum*. The use of animals in this experiment was approved by the decision of the Ministry of Education, Science and Technological Development of Serbia (Decision No. 401-00-9/2011 of 25 January 2011).

For DPM induction in broiler chickens, the EWF method as described by Lien et al. [9] was applied. This method was basically performed in several steps. First, the chicken was held in the palms of the hands at a height of 1 m, after

Table 1. Composition of experimental diets

Ingredients	Starter (0-14 d)	Grower (15-35 d)	Finisher (36-42 d)
Diet composition (g/kg)			
Corn	488.0	528.0	553.0
Full fat soybean	170.0	180.0	165.0
Soybean meal (44% CP)	270.0	220.0	200.0
Soy oil	20.0	20.0	30.0
Monocalcium phosphate	15.0	15.0	15.0
Limestone	18.0	18.0	18.0
Salt	2.5	2.5	2.5
Sodium bicarbonate	3.0	3.0	3.0
L-Lysine HCl	1.0	1.0	1.0
DL-methionine	2.5	2.5	2.5
Vitamin and mineral premix	10	10	10
Calculated nutrient composition			
AME (MJ/kg)	12.87	13.11	13.43
Crude protein (g/kg)	225.5	210.6	198.4
Lysine (g/kg)	13.4	12.4	11.5
Methionine (g/kg)	5.9	5.7	5.6
Ca (g/kg)	11.5	11.4	11.3
P total (g/kg)	7.5	7.3	7.2
P available (g/kg)	4.4	4.3	4.2
CP: crude protein, AME: apparent metabolizable energy			

which it was raised to a height of 2 m. From that height, each chicken was allowed to fall freely back to a height of 1 m. Flapping of the chicken's wings occurred during the fall. One cycle of EWF consisted of the previously described procedure. In our study, broilers were treated with EWF at the end of day 37 post-hatching, with a total treatment duration of 45 seconds, which included about 20 cycles of EWF.

During the rearing period, which lasted 42 days, body weight of broilers was in accordance with performance objectives for Cobb 500, and there were no significant differences between treatments. Subsequently, the 5 male birds from each of the pens 1 to 8 that were subjected to EWF were sacrificed. From each of the pens 1 to 4, as well as from each of the pens 5 to 8, 5 male birds that were not subjected to EWF, with body weights nearest to the mean body weight of the male birds, were also sacrificed.

Detection of DPM Stages

In order to reach the deep pectoral muscle, a ventral cut was made through the breast tissue. A score of DPM 0 was used to indicate no signs of myopathy in deep pectoral muscles. Categorization of DPM into a further 4 stages was made according to Kijowski and Kupinska^[8]. Hemorrhages and clotted blood in the vessels characterized stage 1 (DPM 1). In stage 2 (DPM 2), necrosis and fibrosis of muscle tissue were present, while the color of the deep pectoral muscle was pale pink. Stage 3 (DPM 3) was related to a change in the color of muscle tissue to green, particularly in the central muscle area. A high degree of muscle tissue necrosis characterized stage 4 (DPM 4), which was manifested by an intensive green color of muscle, which in some parts turned to white and gray.

Morphometric and Stereological Examinations

For the purpose of histological analysis, samples of deep pectoral muscle (*M. supracoracoideus*) were taken after determination of macroscopic DPM stage. Muscle tissue samples were fixed in a 10% buffered formalin solution, followed by several stages of tissue processing: dehydration, clearing, and embedding in paraffin^[19,20]. Then, the samples were cut into serial 5µm thick sections using a microtome. Hematoxylin-eosin (H&E) and the Mallory trichrome staining method were performed^[21,22] to observe the following parameters: diameter and volume density of non-necrotic and necrotic muscle cells, diameter and volume density of adipose cells, and volume density of connective tissue proper cells.

Samples were analyzed by light microscopy using a Leica DMLS microscope with a Leica DC 300 digital camera and Leica IM 1000 software (Leica Imaging Systems Ltd, Cambridge, UK). The diameter of muscle as well as of adipose cells was measured as the average of the longest lines drawn across the length and width of their cross-

sections. The M42 multipurpose testing system was used to measure the volume density of certain cells (muscle cells, adipose cells, and connective tissue proper cells). This testing system represents a grid that consists of 21 straight-line segments, while the test points are located at both ends of each line, making 42 test points in total^[23,24]. To calculate the volume density of certain cells, the following formula was used:

$$Vv(c) = \frac{P(c)}{P(m)} \cdot 100 (\%)$$

where $Vv(c)$ is the volume density of certain cells, $P(c)$ is the number of test points lying over the certain cells, and $P(m)$ is the number of test points lying over the muscle.

Statistical Analysis

Analysis of variance (ANOVA) and *post hoc* Tukey's test were used to determinate the influence of different treatments on the value of the observed histological parameters. To study the influence of different treatments on the incidence of DPM stages, chi-square test followed by the Bonferroni correction was used; a significance level of 0.05 was applied. Statistical tests were carried out using the software package Statistica 13.0 (Dell Software, Round Rock, Texas, USA).

RESULTS

Measurements showed that in those groups where chickens were fed a basal diet and treated with EWF, the incidence of DPM stage 0 was lower ($P < 0.01$), while the incidence of DPM stage 3 as well as average DPM stage was higher ($P < 0.01$) compared to groups without EWF treatment (*Table 2*). Between groups with EWF treatment, average DPM stage was lower ($P < 0.05$) in groups where chickens were fed a basal diet with CoQ₁₀ supplementation.

All stages of DPM were detected during the examination of deep pectoral muscles in sacrificed birds (*Fig. 1*). Deep pectoral muscles with no signs of DPM were labeled as DPM 0. Microscopically these muscles showed standard characteristics of skeletal muscles. DPM stage 1, characterized by red colored deep pectoral muscle with excessive fluid in the damaged area, histologically showed hyperemia, edema, and numerous leucocytes. DPM stage 2 was defined macroscopically by a pale plum color of the deep pectoral muscle with an accompanying fibrotic texture, and microscopically by structural alterations such as necrosis of muscle cells with pale cytoplasm and nuclei. Certain muscle cells were swollen during the early stages of necrosis. In DPM stage 3, muscle samples taken from the central, green parts of the deep pectoral muscle, histologically demonstrated an increased quantity of fibrous tissue with the occasional presence of adipose cells. Muscle tissue classed as DPM stage 4 macroscopically

Table 2. Effects of treatments on the incidence of deep pectoral myopathy (DPM) stages in broilers at 42 days post-hatching

Item	Groups			
	Control	Basal Diet with EWF Treatment	CoQ ₁₀ Supplementation	CoQ ₁₀ Supplementation with EWF Treatment
DPM 0 (%)	87.50 ^A	37.50 ^B	87.50 ^A	62.50 ^{AB}
DPM 1 (%)	6.25	6.25	12.50	18.75
DPM 2 (%)	6.25	12.50	0	6.25
DPM 3 (%)	0 ^B	37.50 ^A	0 ^B	12.50 ^{AB}
DPM 4 (%)	0	6.25	0	0
Average DPM stage	0.19 ^{Bab}	1.69 ^{Aa}	0.12 ^{Bab}	0.69 ^{ABb}

EWF: encouraged wing flapping. CoQ₁₀: Coenzyme Q₁₀

^{A, B} Values in the same row without a common superscript capital letter differ significantly ($P < 0.01$)

^{a, b} Values in the same row without a common superscript lowercase letter differ significantly ($P < 0.05$)

Table 3. Effects of treatments on histological parameters of muscle, adipose, and proper connective tissue of *M. supracoracoideus* of broilers at 42 days post-hatching

Parameter	Groups			
	Control	Basal Diet with EWF Treatment	CoQ ₁₀ Supplementation	CoQ ₁₀ Supplementation with EWF Treatment
Muscle tissue				
Diameter of non-necrotic muscle cell (μm)	62.06±1.86	56.40±1.66	62.17±1.47	60.10±1.90
Diameter of necrotic muscle cell (μm)	54.23±3.27	50.99±2.78	55.24±3.86	52.05±2.99
Volume density of non-necrotic muscle cells (%)	82.05±0.70 ^A	47.61±1.00 ^C	83.64±0.77 ^A	64.16±1.28 ^B
Volume density of necrotic muscle cells (%)	0.68±0.15 ^C	21.55±0.98 ^A	0.62±0.18 ^C	8.79±0.84 ^B
Adipose tissue				
Diameter of adipose cell (μm)	24.74±1.75	19.07±1.68	22.61±0.56	20.45±1.73
Volume density of adipose cells (%)	0.36±0.1 ^B	3.05±0.44 ^A	0.42±0.1 ^B	2.54±0.33 ^A
Proper connective tissue				
Volume density of proper connective tissue cells (%)	12.30±0.45 ^B	22.36±0.57 ^A	11.36±0.41 ^B	20.97±0.67 ^A

Values are presented as mean ± standard error of the mean

EWF: encouraged wing flapping. CoQ₁₀: Coenzyme Q₁₀

^{A, B} Values in the same row without a common superscript capital letter differ significantly ($P < 0.01$)


contained whitish regions within areas of green necrotic tissue; in addition, partial replacement of damaged muscle tissue with fibrous tissue and in places with adipose tissue, were detected at the microscopic level.

Histological examination of muscle tissue showed that, in groups with EWF treatment, volume density of necrotic muscle cells was higher ($P < 0.01$), while the volume density of non-necrotic muscle cells was lower ($P < 0.01$) compared to other groups (Table 3). Between groups with EWF treatment, volume density of necrotic muscle cells was higher ($P < 0.01$), while volume density of non-necrotic muscle cells was lower ($P < 0.01$) in groups fed the basal diet. Microscopic observations of adipose and connective tissue showed that the volume density of adipose cells as well as connective tissue proper cells were lower ($P < 0.01$) in groups without EWF treatment compared to other groups.

DISCUSSION

Between the groups subjected to EWF treatment, the groups fed the CoQ₁₀ supplemented diet had a lower volume density of necrotic muscle cells, as well as a higher volume density of non-necrotic muscle cells. The cause of this could be related to the antioxidant properties of CoQ₁₀. Excessive wing flapping is known to lead to ischemia and hypoxia [25].

Under hypoxic conditions, oxidative stress may be induced [14], which results in cell necrosis, tissue damage, and structural modifications within skeletal muscles [12,26]. These effects of oxidative stress can be reduced by antioxidants; so CoQ₁₀ could decrease changes in tissue structure and cell necrosis and be responsible for the lower volume density of necrotic muscle cells in broilers fed the CoQ₁₀ supplemented diet.



The use of different classifications used by authors in determining the stage of DPM could lead to misunderstandings. In papers so far, 3-stage [3] and 4-stage [8] classifications can be found. In order to classify DPM stages more precisely, both macroscopically and microscopically, we applied the 4-stage classification as used by Kijowski and Kupińska [8].

In previous studies in which broilers were subjected to forced wing flapping [8] or EWF [9] at 5 days before slaughter (similar to the current experiment) results obtained were in agreement with ours. Similarly, Kijowski and Kupińska [8] also observed all 4 stages of DPM in Ross 308 and Flex line broilers that were subjected to forced wing flapping. In male broilers (Ross x Cobb 500) subjected to EWF, a 71% prevalence of DPM lesions was detected by Lien et al. [9], which supports the current findings in the group fed a basal diet and subjected to EWF.

If we surmise that the antioxidant CoQ₁₀ can reduce muscle tissue damage and cell necrosis caused by oxidative stress [10,15], it can be assumed that a consequence of that reduction could be a lower average DPM stage in groups of broilers fed a CoQ₁₀ supplemented diet and treated with EWF, as compared to groups of broilers fed a basal diet and treated with EWF. This points out the direct relationship between DPM stage and the degree of tissue damage and cell necrosis. Our results were in agreement with the above mentioned theories; between the groups that were subjected to EWF, the volume density of necrotic muscle cells was lower in groups fed the CoQ₁₀ supplemented diet.

Examination of DPM in house reared broilers [27] report similar observations as were noticed in our study, such as greenish, pale, and swollen muscles with the presence of necrotic and hemorrhagic tissue that were visible during dissection of the pectoral muscle. In addition, leucocyte infiltration, necrotic muscle fibers, large necrotic areas, and fibro-adipose tissue were detected during microscopic examinations.

The question arises, if the EWF treatment, which in our study lasted 45 seconds, is prolonged, how would this be reflected in the signs of DPM. Kijowski and Kupińska [8] show that increased duration of wing activity leads to greater fatigue of the pectoral muscle, which results in a reduced number of wing flaps in additional time. Therefore, it can be surmised that prolonged duration of EWF treatment will lead to enhanced signs of DPM, but not in proportion to the extended time.

Determination of the most suitable day of age of broilers for the application of EWF treatment is very important. In our study, broilers were subjected to EWF treatment at the end of their 37th day, based on previous findings [8,9] where treating broilers too early in their life resulted in almost complete absence of DPM; this was because of the low weight of the pectoral muscle in which symptoms

of myopathy were unable to develop. Meanwhile the application of treatment a short time before slaughter, did not allow time for the later stages of DPM to develop.

Fathi [17] showed that differences in total mortality percentage of broilers due to ascites existed between 2 groups fed diets supplemented with 20 mg and 40 mg of CoQ₁₀ per kg of feed. In previous studies, the age at which broilers were offered CoQ₁₀ supplements differed. Huang et al. [28] offered broilers a CoQ₁₀ supplement diet from 1 day post-hatching, while Fathi [17] offered CoQ₁₀ supplement at day 15. As in our experiment, broilers were treated with 20 mg of CoQ₁₀/kg of finisher diet from days 36 to 42; in further research it would be useful to apply different levels of CoQ₁₀ in treatments and/or to apply the supplemented diets to broilers of different ages.

We were mindful of animal welfare when selecting EWF in this study as the means for inducing DPM. In previous studies [29-31], the applied methods, such as surgical occlusion of the vascular supply, electrically-induced contractions of the muscle, and forced wing exercise, were more stressful and/or painful to broilers than EWF.

Intensive wing flapping is the main reason for the occurrence of DPM in broilers, and by reducing it the incidence of DPM could be decreased. In order to successfully reduce signs of DPM in broilers, several measures must be applied such as reduced human activity in broiler houses as well as animal activity around the house, refraining from catching birds by their wings, providing sufficient space for every bird during resting, moving, eating, or drinking, etc. These measures are in line with the flock management guidelines presented by Bilgili and Hess [3] in order to reduce the incidence of green muscle disease in broiler flocks.

In conclusion, results indicate that, in broilers fed supplemental CoQ₁₀, the average DPM stage as well as effects of DPM on histological parameters of deep pectoral muscle were reduced at 42 days post-hatching. These findings suggest that the antioxidant properties of CoQ₁₀ could reduce the effects of DPM on cell necrosis and change in tissue structure, considering that between the groups subjected to EWF treatment, a lower average DPM stage and volume density of necrotic muscle cells, as well as higher volume density of non-necrotic muscle cells were detected in broilers fed supplemental CoQ₁₀.

REFERENCES

1. Kuttappan VA, Shivaprasad HL, Shaw DP, Valentine BA, Hargis BM, Clark FD, McKee SR, Owens CM: Pathological changes associated with white striping in broiler breast muscles. *Poult Sci*, 92 (2): 331-338, 2013. DOI: 10.3382/ps.2012-02646
2. Bailey RA, Watson KA, Bilgili SF, Avendano S: The genetic basis of pectoralis major myopathies in modern broiler chicken lines. *Poult Sci*, 94 (12): 2870-2879, 2015. DOI: 10.3382/ps/pev304
3. Bilgili SF, Hess JB: Green muscle disease: Reducing the incidence in broiler flocks. *Ross Tech*, 8 (48): 1-8, 2008.
4. Biewener AA: Muscle function in avian flight: Achieving power and

- control. *Philos Trans R Soc Lond B Biol Sci*, 366, 1496-1506, 2011. DOI: 10.1098/rstb.2010.0353
- 5. Bilgili SF, Hess JB:** Green muscle disease in broilers increasing. *World Poult*, 18 (4): 42-43, 2002.
- 6. Surai PF, Fisinin VI:** Vitagenes in poultry production: Part 1. Technological and environmental stresses. *Worlds Poult Sci J*, 72 (4): 721-734, 2016. DOI: 10.1017/S0043933916000714
- 7. Kijowski J, Kupińska E, Kaczmarek A, Stangierski J, Popiół A:** Occurrence and characteristics of chicken breast muscles with DPM symptoms. *Med Weter*, 65 (7): 466-471, 2009.
- 8. Kijowski J, Kupińska E:** Induction of DPM changes in broiler chickens and characteristics of myopathy symptoms. *Bull Vet Inst Pulawy*, 56, 217-223, 2012. DOI: 10.2478/v10213-012-0039-8
- 9. Lien RJ, Bilgili SF, Hess JB, Joiner KS:** Induction of deep pectoral myopathy in broiler chickens via encouraged wing flapping. *J Appl Poult Res*, 21, 556-562, 2012. DOI: 10.3382/japr.2011-00441
- 10. Schieber M, Chandel NS:** ROS function in redox signaling and oxidative stress. *Curr Biol*, 24 (10): R453-R462, 2014. DOI: 10.1016/j.cub.2014.03.034
- 11. Nita M, Grzybowski A:** The role of the reactive oxygen species and oxidative stress in the pathomechanism of the age-related ocular diseases and other pathologies of the anterior and posterior eye segments in adults. *Oxid Med Cell Longev*, 2016, 2016: 3164734, 2016. DOI: 10.1155/2016/3164734
- 12. Mittal M, Siddiqui MR, Tran K, Reddy SP, Malik AB:** Reactive oxygen species in inflammation and tissue injury. *Antioxid Redox Signal*, 20 (7): 1126-1167, 2014. DOI: 10.1089/ars.2012.5149
- 13. Mayr M, Sidibe A, Zampetaki A:** The paradox of hypoxic and oxidative stress in atherosclerosis. *J Am Coll Cardiol*, 51 (13): 1266-1267, 2008. DOI: 10.1016/j.jacc.2008.01.005
- 14. Miyata T, Takizawa S, van Ypersele de Strihou C:** Hypoxia. 1. Intracellular sensors for oxygen and oxidative stress: Novel therapeutic targets. *Am J Physiol Cell Physiol*, 300, C226-C231, 2011. DOI: 10.1152/ajpcell.00430.2010
- 15. Littarru GP, Tiano L:** Bioenergetic and antioxidant properties of coenzyme Q₁₀: Recent developments. *Mol Biotechnol*, 37, 31-37, 2007. DOI: 10.1007/s12033-007-0052-y
- 16. Crane FL:** Biochemical functions of coenzyme Q₁₀. *J Am Coll Nutr*, 20 (6): 591-598, 2001.
- 17. Fathi M:** Effects of coenzyme Q₁₀ supplementation on growth performance, some hematological parameters, plasma enzymes activities in broilers with pulmonary hypertension syndrome (PHS). *Iran J Appl Anim Sci*, 5 (1): 147-153, 2015.
- 18. Gopi M, Purushothaman MR, Chandrasekaran D:** Effect of coenzyme Q₁₀ supplementation on serum protein, mineral status, blood picture and immune status in broilers. *Indian J Exp Biol*, 54, 808-815, 2016.
- 19. Howroyd P, Hoyle-Thacker R, Lyght O, Williams D, Kleymenova E:** Morphology of the fetal rat testis preserved in different fixatives. *Toxicol Pathol*, 33 (2): 300-304, 2005. DOI: 10.1080/01926230590896145
- 20. Slaoui M, Fiette L:** Histopathology procedures: From tissue sampling to histopathological evaluation. *Methods Mol Biol*, 691, 69-82, 2011. DOI: 10.1007/978-1-60761-849-2_4
- 21. Bancroft JD, Layton C:** The hematoxylin and eosin. In, Suvarna SK, Layton C, Bancroft JD (Eds): *Bancroft's Theory and Practice of Histological Techniques*. 7th edn., 173-186, Churchill Livingstone, London, 2012.
- 22. Bancroft JD, Layton C:** Connective and mesenchymal tissues with their stains. In, Suvarna SK, Layton C, Bancroft JD (Eds): *Bancroft's Theory and Practice of Histological Techniques*. 7th edn., 187-214, Churchill Livingstone, London, 2012.
- 23. Burity CHF, Pissinatti A, Mandarim-De-Lacerda CA:** Stereology of the liver in three species of *Leontopithecus* (Lesson, 1840) Callitrichidae - Primates. *Anat Histol Embryol*, 33 (3): 183-187, 2004. DOI: 10.1111/j.1439-0264.2004.00533.x
- 24. Drábeková J, Uhrín V, Pivko J, Macák V, Pavlík V, Jančová A, Faragová N:** Comparison of quantitative microscopic and submicroscopic analyses of the liver in cows bred in various polluted regions. *Bull Vet Inst Pulawy*, 49, 79-84, 2005.
- 25. Petracci M, Cavani C:** Muscle growth and poultry quality issues. *Nutrients*, 4, 1-12, 2012. DOI: 10.3390/nu4010001
- 26. Choi MH, Ow JR, Yang ND, Taneja R:** Oxidative stress-mediated skeletal muscle degeneration: Molecules, mechanisms, and therapies. *Oxid Med Cell Longev*, 2016, 2016: 6842568, 2016. DOI: 10.1155/2016/6842568
- 27. Ozmen O:** Pathological examination of deep pectoral myopathy in house reared broilers. *Kafkas Univ Vet Fak Derg*, 23 (5): 831-834, 2017. DOI: 10.9775/kvfd.2017.17989
- 28. Huang B, Guo Y, Hu X, Song Y:** Effects of coenzyme Q₁₀ on growth performance and heart mitochondrial function of broilers under high altitude induced hypoxia. *J Poult Sci*, 48 (1): 40-46, 2011. DOI: 10.2141/jpsa.010084
- 29. Siller WG, Wight PA, Martindale L, Bannister DW:** Deep pectoral myopathy: An experimental simulation in the fowl. *Res Vet Sci*, 24 (2): 267-268, 1978.
- 30. Wight PAL, Siller WG, Martindale L, Filshie JH:** The induction by muscle stimulation of a deep pectoral myopathy in the fowl. *Avian Pathol*, 8 (1): 115-121, 1979. DOI: 10.1080/03079457908418332
- 31. Harper JA, Bernier PE, Thompson-Cowley LL:** Early expression of hereditary deep pectoral myopathy in turkeys due to forced wing exercise. *Poult Sci*, 62 (12): 2303-2308, 1983. DOI: 10.3382/ps.0622303