

RESEARCH ARTICLE

Comparative Analysis of the Heart Tissue Transcriptomes Between Low-altitude Reared and High-altitude Reared Bar-headed Geese (*Anser indicus*)

Ying LI ^{1,a} Fang WANG ^{2,b} Xiaolong GAO ^{1,c} Lilin ZHU ^{3,d} Wen WANG ^{1,e (*)} Kirill SHARSHOV ^{4,f}¹ State Key Laboratory of Plateau Ecology and Agriculture, Qinghai University, Xi'ning 810016, Qinghai Province, CHINA² Northwest Institute of Plateau Biology, Chinese Academy of Sciences, Xi'ning 810008, Qinghai Province, CHINA³ Xi'ning Wild Animal Epidemic Disease Monitoring Station, Xi'ning 810001, Qinghai Province, CHINA⁴ Federal Research Center of Fundamental and Translational Medicine, Novosibirsk 630117, RUSSIAORCID: ^a 0000-0002-1590-9992; ^b 0000-0002-8384-4561; ^c 0000-0002-6698-6138; ^d 0000-0001-5569-5637; ^e 0000-0002-3156-7929^f 0000-0002-3946-9872

Article ID: KVFD-2021-25704 Received: 02.03.2021 Accepted: 17.06.2021 Published Online: 17.06.2021

Abstract

The bar-headed geese (*Anser indicus*) are renowned for high-altitude migratory flights and they must fly over the Qinghai-Tibetan Plateau for their annual migration. Through comparing the high-altitude bar-headed geese with the other closely related low-altitude species, many efforts have been made to reveal the unique adaptations at physiological, biochemical, and behavioral levels that help bar-headed geese living and flying in high-altitude conditions. Nonetheless, little is known about the transcriptome level changes of the bar-headed geese adaptation to low-altitude environment. To explore the variations of gene expression that were induced by low-altitude environment in the bar-headed geese, we conducted the first comparative transcriptomic analysis of heart tissues between bar-headed geese reared in high-altitude regions (~3000 m), and the bar-headed geese reared at the low-altitude regions (~30 m) for nearly three years. A total of 76 differentially expressed genes (DEGs) were detected in the low-altitude bar-headed geese compared with the high-altitude bar-headed geese. Gene ontology (GO) and Kyoto Encyclopedia of Genes and Genomes (KEGG) analysis showed that these DEGs were mainly involved in the focal adhesion, extracellular matrix (ECM) - receptor interaction, the mammalian target of rapamycin (mTOR) signaling pathway, wingless-type (Wnt) signaling pathway, and glycosaminoglycan degradation etc. The results will be useful for understanding the divergent adaptation of the bar-headed geese to different altitude environment, and the transcriptome data provides a valuable resource for future functional studies.

Keywords: *Anser indicus*, Transcriptome, Adaptation, Altitude, Heart

Düşük ve Yüksek İrtifalı Alanlarda Yetiştirilen Çubuk Başlı Kazlarda (*Anser indicus*) Kalp Dokusu Transkriptomlarının Karşılaştırmalı Analizi

Öz

Çubuk başlı kazlar (*Anser indicus*), yüksek irtifada göçmen uçuşlarıyla ünlüdür ve yıllık göçleri için Tibet Platosu üzerinden uçmaları gerekir. Yüksek irtifaya adapte çubuk başlı kazları, diğer yakından ilişkili alçak irtifalı türlerle karşılaştırarak, bu kazların yüksek irtifa koşullarında yaşamasına ve uçmasına yardımcı olan fizyolojik, biyokimyasal ve davranışsal seviyelerde benzersiz adaptasyonlarını ortaya çıkaracak birçok çalışma yapılmıştır. Bununla birlikte, çubuk başlı kazların alçak irtifalı ortamlara adaptasyonlarının transkriptom seviyesi değişiklikleri ile ilgili çok az şey bilinmektedir. Çubuk başlı kazlarda düşük irtifa ortamı tarafından indüklenen gen ekspresyon varyasyonlarını araştırmak için, yüksek irtifa bölgelerinde (~3000 m) yetiştirilen çubuk başlı kazlar arasındaki kalp dokularının ilk karşılaştırmalı transkriptomik analizini gerçekleştirdik ve bu kazlar yaklaşık üç yıl boyunca alçak irtifalı bölgelerde (~30 m) yetiştirildi. Yüksek irtifalı çubuk başlı kazlarla karşılaştırıldığında, alçak irtifalı çubuk başlı kazlarda farklı ekspresye edilmiş (DEG) toplam 76 gen tespit edildi. Gen ontolojisi (GO) ve Kyoto Genler ve Genom Ansiklopedisi (KEGG) analizi, bu DEG'lerin temel olarak fokal adezyon, ekstrasellüler matris (ECM) - reseptör etkileşimi, rapamisin protein kompleksinin memeli hedefi (mTOR) sinyal yolu, wingless-tip (Wnt) sinyal yolu ve glikozaminoglikan degradasyonu vb. ile ilişkili olduğunu göstermiştir. Çalışma sonuçları, çubuk başlı kazların farklı irtifa ortamlarına farklı adaptasyonlarını anlamak için faydalı olacaktır ve elde edilen transkriptom verileri gelecekteki fonksiyonel çalışmalar için değerli bir kaynak imkanı sunacaktır.

Anahtar sözcükler: *Anser indicus*, Transkriptom, Adaptasyon, İrtifa, Kalp

How to cite this article?

Li Y, Wang F, Gao X, Zhu L, Wang W, Sharshov K: Comparative analysis of the heart tissue transcriptomes between low-altitude reared and high-altitude reared bar-headed geese (*Anser indicus*). *Kafkas Univ Vet Fak Derg*, 27 (4): 455-463, 2021.
DOI: 10.9775/kvfd.2021.25704

(*) Corresponding Author

Tel: +86-971-5201533

E-mail: 279334395@qq.com (W. Wang)



This article is licensed under a Creative Commons Attribution-NonCommercial 4.0 International License (CC BY-NC 4.0)

INTRODUCTION

The bar-headed goose (*Anser indicus*) is endemic to Asia, breeding in the high-altitude plateau wetlands of central Asia [1-4], wintering mainly in the south-central Tibet (approximately 56.88% of the world populations) and India (around 27.84% of the world populations) [1,4]. This species is famous for extremely high-altitude flying over the Himalayan Mountains twice a year on the Central Asian flyway between the wintering areas in India subcontinent and the summering grounds on the Qinghai-Tibetan Plateau [5]. The bar-headed geese were tracked flying as high as 7.290 meters up [6], and mountaineers even reported seeing them fly over the summit of Mount Everest (8.850 meters above sea level) [7]. At these heights, the partial pressure of oxygen is one-third that of sea level. Incredibly, bar-headed geese are able to maintain the high metabolic and oxygen consumption rates necessary for flapping flight under severe hypoxia conditions [8]. Not surprisingly, there has been increasing interest among researchers in understanding the physiological, molecular, and behavioral adaptations of bar-headed geese that allow for the exceptional high-altitude flight performances [9,10].

Through comparing the high-altitude bar-headed geese with the other closely related low-altitude species, a growing body of literature has identified several unique adaptations that help bar-headed geese flying in low oxygen conditions. For example, physiological adaptations have evolved at every step in the oxygen transport cascade of bar-headed geese to help them accomplish the high-altitude success. Such adaptations include an effective breathing pattern [11], larger lungs [12], an enhanced hypoxic ventilatory response [13], hemoglobin (Hb) with a higher oxygen affinity [14], an increase in flight and cardiac muscle capillarity [15], and subcellular redistribution of the mitochondrial reducing oxygen diffusion distances [15]. The increased affinity between Hb and O₂ plays a critical role in adaptation of bar-headed geese to hypoxia through enhancing pulmonary O₂ loading. A few studies have found that in the bar-headed goose the major isoform of Hb has significantly higher O₂-affinity than that of the closely related low-altitude species due to the single and large-effect substitutions in the hemoglobin gene [16,17]. In addition, a roller coaster strategy was reported to take by bar-headed geese, rising and falling with the relief of the terrain, to minimize energy expenditure during their trans-Himalayan flights [18]. Overall, these studies have greatly improved our understanding of the adaptive mechanism of bar-headed goose for high-altitude conditions.

Recent advances in genomics have opened a huge opportunity to study the genetic basis of high-altitude adaptation characteristics [19,20]. In our previous study, we reported the first *de novo* whole genome sequencing, assembly, gene prediction and annotation of the bar-headed goose [21]. Transcriptome is a useful tool to analyze

changes in gene expression, and has been widely used to explore high-altitude adaptations in a group of birds [22]. Therefore, in this study, we aimed to compare the transcriptome profiles of heart tissues between bar-headed geese reared in low- and high-altitude environment. Unlike previous studies that compared high-altitude bar-headed geese with the other lower-altitude birds, our current study focused on comparison between the low-altitude reared bar-headed geese, where these geese were transported to and there is no distribution of wild bar-headed geese, with the high-altitude reared bar-headed geese. The major advantage of this experimental setup is that the genetic backgrounds of the subjects being compared are the same. The results will enrich the transcriptomic resources of bar-headed geese and laid a foundation for the functional study of genes related to low altitude adaptation in this species.

MATERIAL AND METHODS

Ethics Statement

This study conformed to the guidelines for the care and use of experimental animals established by the Ministry of Science and Technology of the People's Republic of China (Approval number: 2006-398). The research protocol was reviewed and approved by the Ethical Committee of Qinghai University.

Animal Sample Collection

Eggs of wild bar-headed geese were collected and hatched at the Fei Yan specialized breeding and rearing farming cooperative (elevation 3.000 meter) in Huangzhong District, Xi'ning City, Qinghai Province, China. After hatch, three individuals were reared at this cooperative for three years until the experiment began and labeled as high-altitude group (HA group, n=3) (Fig. 1). Another three individuals were transported to and reared at the He Ming breeding Co., LTD. (elevation 30 meter) in Hengshui City, Hebei Province, China. The latter three individuals were also reared for three years until the experiment began and labeled as low-altitude group (LA group, n=3) (Fig. 1).

These bar-headed geese lived freely in both wild and artificially reared environments. These birds were healthy during experimental period. The heart tissues were sampled rapidly from each carcass, and immediately frozen in the liquid nitrogen. All heart samples were stored at -80°C until used.

RNA Extraction and Transcriptome Sequencing

For each sample, total RNA was isolated using Trizol reagent (Invitrogen, California, USA) following the manufacturer's instructions. Residual genomic DNA was digested by RNase-free DNase (Qiagen, Germany). The RNA concentration and overall quality were assessed using a Qubit 2.0 Fluorometer (Invitrogen, California, USA), and an Agilent 2100




Fig 1. The schematic diagram of our experiment

bioanalyser (Agilent Technologies, USA), respectively. The RNA integrity number threshold was set at 7.0 for the construction of library. mRNA was further purified using poly-T oligo-attached magnetic beads. Finally, six libraries with 350 bp insert sizes were constructed using the NEBNext Ultra™ RNA Library Prep Kit for Illumina (NEB, USA). The constructed libraries were then sequenced by the Illumina NovaSeq 6000 platform at Novogene Bioinformatics Technology Co. Ltd (Beijing, China). The whole process followed a standardized procedure and was monitored by Novogene's Quality Control System.

Data Accessibility

The raw sequencing data has been deposited in the National Center for Biotechnology Information (NCBI) sequence read archive (SRA) with the bioproject number PRJNA612653.

Bioinformatics Analyses

- Quality Control

To produce the high-quality clean reads, raw reads were assessed for quality using FastQC v.0.11.9 (<https://www.bioinformatics.babraham.ac.uk/projects/fastqc/>) and filtered for low-quality reads according to the following rules: 1) removing the reads with adapter contamination; 2) removing the reads with unknown nucleobases; 3) removing the low-quality reads (reads with a $Q_{\text{PHRED}} \leq 20$ base number accounting for more than 50% of the total read length). At the same time, Q20, Q30 and GC content of the clean reads were counted. The clean reads with high quality were used for further analysis.

- Genome Mapping

The bar-headed goose reference genome (DDBJ/ENA/

GenBank under the accession VDDG01000000) [21] was used as the reference genome for our transcriptome sequencing data. Genome mapping was conducted using HISAT2 v2.0.5 software [23].

- Quantification of Gene Expression Level

FeatureCounts v1.5.0-p3 [24] was used to calculate the number of mapped reads to each gene, which were then normalized for the gene length and library size. FPKM (Fragments Per Kilobase of transcript per Million fragments mapped) of each gene was counted based on the length of the gene and reads count mapped to this gene.

- Differential Expression Analysis

Genes with very low expression levels were filtered out of the dataset. Then, differentially expressed genes were analyzed using DESeq2 R package (1.16.1) [25] between HA and LA groups. The DEGs were identified using the thresholds of $|\log_2 \text{fold change}| \geq 1.0$ and $\text{P}_{\text{adj}} < 0.05$. Clustering analysis of the identified DEGs was implemented by the heatmaps in R software.

- GO Function and KEGG Pathway Enrichment Analysis

GO functional enrichment and KEGG pathway analysis of DEGs was conducted by the clusterProfiler R package [26], in which gene length bias was corrected. GO terms or KEGG pathways with $P < 0.05$ were significantly enriched.

- Real-time Quantitative PCR (RT-qPCR) Verification

To verify the repeatability of DEGs derived from transcriptome data, 16 DEGs were selected randomly for RT-qPCR validation. The primer sequences were shown in Table 1. RNAiso Pure RNA Isolation Kit (TaKaRa, Dalian, China) was used to extract RNA. A PrimeScript RT reagent

Table 1. Primers used in this study for real-time quantitative PCR (RT-qPCR)

| Gene | Regulation | Primer Sequences (5'-3') | Amplicon Size (bp) |
|---------|------------|----------------------------------|--------------------|
| DUSP1 | Up | Forward: CGATGGAGGAAGGGTGTGG | 152 |
| | | Reverse: TGAAGTTGGGGAGATGATGC | |
| NR4A1 | Up | Forward: CCCTCCAAGCCCAAGCA | 238 |
| | | Reverse: CGGCGAAGCCCTGAATC | |
| FOS | Up | Forward: TTCTATGCGTCGACTGGG | 113 |
| | | Reverse: AAGGTGGAGGTGTAGGTGCTG | |
| JUN | Up | Forward: GGAAAAGGAAGTTGGAAAGGAT | 155 |
| | | Reverse: CTGGCACCCGCTGTTGA | |
| HSP70 | Up | Forward: AGCGTAACACCACCATTCCC | 94 |
| | | Reverse: TCACCTCATAACCTGGACC | |
| FABP | Up | Forward: CCAAGCCCAATGTAACATCAGC | 237 |
| | | Reverse: CGACCAGTTCCCATCCAC | |
| APOA1 | Up | Forward: GACCCTCGCCTGCTCTT | 209 |
| | | Reverse: GCGTGTCCAGGTTGTCGG | |
| SCD | Up | Forward: TCAACCCACGGGAGAACC | 213 |
| | | Reverse: TCCGCATTTCCGAGCC | |
| ACACB | Down | Forward: TGAGGTGGGGATGTTGC | 96 |
| | | Reverse: TTGTGCGTGATGCTGTTGC | |
| TNC | Down | Forward: CCAAGGGGCACCAACAA | 103 |
| | | Reverse: CGGAAGCCGTCTGGAGTAGC | |
| COL4A1 | Down | Forward: ACAAGGCAATGAAAGAGCACA | 150 |
| | | Reverse: GGCGTTGACAGCCAGTAGG | |
| MYLK | Down | Forward: CGCCTACCAGCCCGATAA | 177 |
| | | Reverse: CAGGGTCAGGATAGCCTCAA | |
| FLNC | Down | Forward: CATCAAGAACGACAACGACACC | 108 |
| | | Reverse: AACGGGCTGCTGGGGAT | |
| GK | Down | Forward: ATGTAACCAATGCCAGTAGAACG | 185 |
| | | Reverse: GTCACCCAAGCACCCAGAA | |
| GPCPD1 | Down | Forward: TGGATGCTGGGTGGCTGAC | 193 |
| | | Reverse: TTCTGAGGAAGTGTAGGGGATGT | |
| SLC9A1 | Down | Forward: GCAGTCGCCAGAGTCCGT | 110 |
| | | Reverse: GCTCCTTCACCCGCATCA | |
| TUBULIN | - | Forward: ATCAGGTGGTCCCAAGAGC | 143 |
| | | Reverse: GGTGATGAGATGGCGGAGG | |

DUSP1: Dual Specificity Phosphatase 1; **NR4A1:** Nuclear Receptor Subfamily 4 Group A Member 1; **FOS:** Fos Proto-Oncogene, AP-1 Transcription Factor Subunit; **JUN:** Jun Proto-Oncogene, AP-1 Transcription Factor Subunit; **HSP70:** Heat Shock Protein 70; **FABP:** Fatty Acid Binding Protein; **APOA1:** Apolipoprotein A1; **SCD:** Stearoyl-CoA Desaturase; **ACACB:** Acetyl-CoA Carboxylase Beta; **TNC:** Tenascin C; **COL4A1:** Collagen Type IV Alpha 1 Chain; **MYLK:** Myosin Light Chain Kinase; **FLNC:** Filamin C; **GK:** Glycerol Kinase; **GPCPD1:** Glycerophosphocholine Phosphodiesterase 1; **SLC9A1:** Solute Carrier Family 9 Member A1

kit (TaKaRa, Dalian, China) was used to synthesize the single stranded cDNA. qPCR was conducted on Gene9600 RT-PCR detection system (Hangzhou Bioer Technology Co. Ltd., Hangzhou, China) and TB Green Fast qPCR Mix (TaKaRa, Dalian, China). The 6 samples were run in triplicate for the 16 genes. *TUBULIN* was used as an internal control in this study. The method of $2^{-\Delta\Delta CT}$ [27] was used to count the relative expression of the 16 genes.

RESULTS

Overview of Transcriptome Sequencing Data

Six sequencing libraries from the two distinct groups (high- and low-altitude) were constructed and sequenced using the Illumina HiSeq platform. The overall sequencing results were shown in *Table 2*. A total of 391,970,972 raw

Table 2. Summary of read statistics from the transcriptome sequencing

| Samples | Raw Reads | Clean Reads | Clean Bases (Gb) | Error Rate (%) | Q20 (%) | Q30 (%) | GC (%) |
|---------|------------|-------------|------------------|----------------|---------|---------|--------|
| HA_h1 | 59,914,010 | 58,248,934 | 8.74 | 0.03 | 96.34 | 90.93 | 50.84 |
| HA_h2 | 64,648,766 | 62,867,252 | 9.43 | 0.03 | 96.62 | 91.39 | 50.56 |
| HA_h3 | 67,105,524 | 65,327,764 | 9.80 | 0.03 | 96.05 | 90.13 | 50.42 |
| LA_h1 | 68,834,484 | 67,275,492 | 10.09 | 0.03 | 96.76 | 91.65 | 50.20 |
| LA_h2 | 66,169,684 | 64,443,694 | 9.67 | 0.03 | 96.29 | 90.65 | 50.56 |
| LA_h3 | 65,298,504 | 63,595,928 | 9.54 | 0.03 | 96.49 | 91.15 | 50.46 |

Table 3. Reads coverage mapping to the reference genome of bar-headed goose

| Samples | Total Reads | Total Map | Unique Map | Multiple Map | Exon | Intron | Intergenic |
|---------|-------------|--------------------|--------------------|----------------|-------------------------|-----------------------|-------------------------|
| HA_h1 | 58,248,934 | 36,857,118(63.28%) | 36,275,462(62.28%) | 581,656(1.0%) | 3,699,777,153(67.2575%) | 571,953,754(10.3974%) | 1,229,188,619(22.3451%) |
| HA_h2 | 62,867,252 | 40,369,734(64.21%) | 39,737,652(63.21%) | 632,082(1.01%) | 4,014,112,346(66.5661%) | 614,605,303(10.192%) | 1,401,548,436(23.2419%) |
| HA_h3 | 65,327,764 | 40,977,873(62.73%) | 40,401,956(61.85%) | 575,917(0.88%) | 4,066,706,910(66.433%) | 654,328,150(10.689%) | 1,400,487,751(22.8781%) |
| LA_h1 | 67,275,492 | 41,840,766(62.19%) | 41,197,140(61.24%) | 643,626(0.96%) | 4,245,606,572(67.9315%) | 648,989,314(10.3841%) | 1,355,233,098(21.6843%) |
| LA_h2 | 64,443,694 | 40,178,319(62.35%) | 39,525,864(61.33%) | 652,455(1.01%) | 4,097,059,115(68.2738%) | 607,198,221(10.1184%) | 1,296,665,508(21.6078%) |
| LA_h3 | 63,595,928 | 39,515,638(62.14%) | 38,940,133(61.23%) | 575,505(0.9%) | 3,805,155,967(64.4771%) | 696,974,108(11.81%) | 1,399,431,085(23.7129%) |

reads were generated and were deposited in the NCBI SRA database (accession number SRP252835). All error rates were less than 0.04%, and the quality of the base values were $Q20 \geq 96\%$ and $Q30 \geq 90\%$ in the six samples, suggesting that the transcriptome data were technically qualified. After filtering the adapter sequences, the ambiguous N nucleotides, and the low-quality reads, we got a total of 381,759,064 clean reads. These clean reads were then used for subsequent analysis. The percentage of clean reads among raw reads ranged from 97.22% to 97.74% in each sample. The guanine + cytosine (GC) values ranged from 50.20% to 50.84%.

The mapping results showed that more than 62% of the clean reads (total mapped reads) matched to the bar-headed goose reference genome (Table 3). Among the mapped reads, more than 61% of the clean reads were matched to a unique genomic location, and less than 1% of the clean reads were matched to multiple genomic locations (Table 3). For the high-altitude group and low-altitude group, 66.75% and 66.89% were mapping to the exon region, 10.43% and 10.77% belonged to the intron region, and 22.82% and 22.34% were in the intergenic sequence respectively (Table 3). These results suggested that the transcriptome data were good enough for further analyses.

DEGs Analysis and RT-qPCR Validation

DEGs between the low- and high-altitude bar-headed geese hearts were analyzed by the DESeq2 package, with the criteria of $|\log_2 \text{fold change}| \geq 1.0$ and $\text{P}_{adj} < 0.05$. A total of 76 DEGs were identified, of which 34 showed up-regulation, and 42 showed down-regulation in the low-altitude group compared with the high-altitude group (Fig. 2).

A heat map of the DEGs between the low- and high-altitude




Fig 2. Differentially expressed genes (DEGs) in low-altitude hearts (LA_h) versus high-altitude hearts (HA_h). Red indicates up regulated genes, green indicates down regulated genes, and blue indicates genes with unchanged expression

groups was shown with hierarchical clustering of the samples and genes in Fig. 3. Hierarchical clustering result clearly separated the high- and low-altitude groups, while samples within each group showed strong correlations (Fig. 4).

In order to verify the reliability of the candidate DEGs, we selected and quantified the expression levels for 16 DEGs (8 up-regulated DEGs and 8 down-regulated DEGs)

by using the RT-qPCR technique (Fig. 5). We successfully confirmed the differential expression for all the selected DEGs (Fig. 5). The results revealed the tendency of gene expression was in line with the transcriptome results,

suggesting the transcriptome sequencing results were reliable for following functional analysis.

Functional Analysis for DEGs

To further explore the possible biological functions and metabolic pathways of the identified DEGs, GO and KEGG enrichment analyses were performed.

The DEGs in the comparison group (LA_h vs. HA_h) were first annotated by the GO function database. The DEGs were significantly enriched in 1 terms of BP (biological process), 1 terms of CC (cellular component), and 11 terms of MF (molecular function) (Fig. 6). Under the molecular function category, the genes mainly enriched in the functions involved in “extracellular matrix structural constituent”, “GTPase activity”, “hydrolase activity”, and “nucleotide/nucleoside binding” (Fig. 6).

We then performed a KEGG analysis of the DEGs. In our study, 76 DEGs were mapped to 32 KEGG pathways. A total




Fig 3. Heat map of the 76 differentially expressed genes (DEGs) in the two groups. Colors represent the normalized gene expression values of DEGs




Fig 4. Hierarchical clustering (Pearson correlation) of transcriptome sequencing results of low-altitude hearts (LA_h, n=3) and high-altitude hearts (HA_h, n=3)




Fig 5. Validation of RNA-seq results by Real-time quantitative PCR (RT-qPCR). (A) validation of up-regulated differentially expressed genes (DEGs) by RT-qPCR, (B) validation of down-regulated differentially expressed genes (DEGs) by RT-qPCR. The RT-qPCR results are the means ± standard deviations (± SDs) of three replicates




Table 4. Differentially expressed genes for the six significant KEGG signal pathways

| KEGG Signal Pathway | Up_regulated Genes | Down_regulated Genes |
|------------------------------------------------------|--------------------|------------------------------------------|
| ECM-receptor interaction | - | TNC, COL4A1, COL4A2, THBS1, COL4A5 |
| Focal adhesion | - | TNC, COL4A1, MYLK, COL4A2, THBS1, COL4A5 |
| AGE-RAGE signaling pathway in diabetic complications | EGR1 | COL4A1, COL4A2, COL4A5 |
| Glycosaminoglycan degradation | - | HYAL2 |
| mTOR signaling pathway | FZD8, FZD2 | ULK2, ATP6V1C2 |
| Wnt signaling pathway | FZD8, FZD2, AXIN2 | - |

of 6 statistically significant pathways were identified when comparing LA_h group with HA_h group, including three in environmental information processing, one in cellular processes, one in human diseases, and one in metabolism categories, respectively (Table 4). Within the upregulated DEGs, the most enriched pathway term was “Wnt signaling pathway” (Table 4). For the downregulated DEGs, the dominant pathways were as follows: “focal adhesion”, “ECM-receptor interaction”, “Advanced glycation end-product (AGE) - role of receptor for advanced glycation end-product (RAGE) signaling pathway in diabetic complications”, and “glycosaminoglycan degradation” (Table 4).

DISCUSSION

The bar-headed goose provides an extraordinary opportunity to investigate the mechanism of high-altitude adaptation for their migratory flight at extreme altitude, and attracted many experts from various fields [5,9,10,12]. However, the adaptive mechanism of bar-headed geese migrating to low altitude environment has not been

reported. We newly sequenced a bar-headed goose genome to provide a useful resource for detection of genomic adaptive changes at the DNA sequence level associated with this species [21]. Transcriptome intermediate between DNA sequences and physiological traits can extend the genomic information by concentrate on gene expression and molecular pathways involved in different altitude adaptation. Here, for the first time, we performed transcriptome analysis in bar-headed geese reared in both high-altitude and low-altitude environment for one hypoxia-sensitive tissue, heart. In this study, bar-headed geese were transported to low-altitude areas for artificial rearing, and then comparative transcriptome analysis was conducted with high-altitude bar-headed geese. This kind of comparison, rather than comparing high-altitude bar-headed geese with the relative geese species living at low-altitude, will better explore the adaptability of bar-headed geese to low-altitude environment.

In this study, each sample obtained more than 8.74 GB of high-quality clean sequencing data, and the base quality

value (error rate, Q20, and Q30) indicating good quality and high reliability. Eight up-regulated DEGs and 8 down-regulated DEGs were chosen for RT-qPCR test to confirm the expression levels calculated by FPKM values. The similar changing trends were found between RT-qPCR validation and transcriptome sequencing. Overall, these results indicate that the results of our transcriptome data are reliable.

In this work, GO annotation results showed that most DEGs were found to be related to cell surface receptor signaling pathway, extracellular matrix structural constituent, GTPase and hydrolase activity, and nucleotide/nucleoside binding. This suggests that when bar-headed geese migrate from high altitude to low altitude, the interactions between membrane receptors and extracellular ligands of the heart were changed. The results were similar to those of Qi et al. in the study of yak heart transcriptome [28]. The effects of these changes on the health status of low-altitude bar-headed geese need to be studied and determined with the help of more morphological, physiological, and pathological data in the future. The annotation analysis of metabolic pathways of DEGs is helpful for further interpretation of gene function. KEGG database is the main public database on metabolic pathways, and six KEGG pathways were significantly enriched in this study. "Wnt signaling pathway" (*FZD8*, *FZD2*, *AXIN2*) was the only upregulated KEGG pathways in the hearts of low-altitude bar-headed geese. A large body of evidence shows that the Wnt signaling pathway plays an important role in various stages of heart development [29]. While this pathway is quiescent in adult hearts, and activation occurs upon pathological stress [30]. In the future, it is necessary to further study the correlation between the activation of this signaling pathway and heart disease of bar-headed geese reared at low altitude. "focal adhesion" and "ECM-receptor interaction" were found to be downregulated in the hearts of low-altitude bar-headed geese. Collagens that encode cell-ECM (including downregulated *COL4A1*, *COL4A2*, and *COL4A5*) are the major structural components of the basement membrane and are important during cardiac morphogenesis by promoting cardiomyocyte proliferation [31]. We also identified three DEGs in focal adhesion pathways (including *TNC*, *MYLK*, and *THBS1*) with significantly downregulated in the hearts of low-altitude bar-headed geese. Those genes usually provide a force-transmitting physical link between the EMC and cytoskeleton, and are helpful for the species to adapt to high altitude [28]. But we hypothesized that the expression of these genes was suppressed in low-altitude environment where oxygen was abundant.

Glycosaminoglycans are a class of anionic linear periodic polysaccharides, which play a key role in tissue homeostasis not only by providing mechanical load resistance but also signaling mediators that play an important physiological function in cell proliferation, differentiation, metastasis,

and recognition [32]. The glycosaminoglycan degradation pathway in the hearts of low-altitude bar-headed geese is involved in the regulation of the structure and function of glycosaminoglycans in the hearts. The down-regulated expression of *HYAL2* in this pathway was detected by transcriptome data. Hyaluronidase is an endogenous glycosidase that can degrade glycosidic bonds in hyaluronic acid polymers (one form of glycosaminoglycans). We hypothesized that high-altitude bar-headed geese need hyaluronidase and other hyaluronidase to maintain the permeability of extracellular matrix of heart, so as to facilitate oxygen exchange and transportation in anoxic environment. However, when high-altitude bar-headed geese migrate to low-altitude environment for growth, oxygen supply is sufficient, and hyaluronic acid in the extracellular matrix of heart does not need to be greatly degraded, so the glycosaminoglycan degradation pathway is inhibited. All these differentially expressed signal pathways and metabolic pathways in the hearts of bar-headed geese intercrossed with each other, and thus forming a huge and complex network. The specific regulatory mechanism and biological effects of these differentially expressed signaling pathways remain to be further studied. Our study has a limitation that also suggests direction for further research. We did not do biochemical, physiological, and histological tests, and how these observed differently expressed genes and pathways may translate into functional and phenotypic changes remains unclear. However, the transcriptome data obtained in this study provides us with a useful candidate pathway and gene resource for future functional studies of altitude adaptation.

In conclusion, this study lays the first foundations for comparative analysis of transcriptional and signaling pathway changes in bar-headed geese heart tissue under low-altitude environment. A total of 76 DEGs were identified, of which 34 showed up-regulation and 42 showed down-regulation in the low-altitude bar-headed geese compared with the high-altitude bar-headed geese. Nine KEGG pathways and 13 GO terms were significantly enriched in low-altitude bar-headed geese. Further studies are required to confirm the functions of these different pathways and genes reported here between low-altitude bar-headed geese and high-altitude bar-headed geese, and the extent to which these differences may contribute to the low-altitude adaptation.

ACKNOWLEDGMENTS

We thank Xiulan Liu (Fei Yan specialized breeding and rearing farming cooperative, China) and Professor Dongming Li (Hebei Normal University, China) for assistance with samples collection.

DECLARATION OF CONFLICTING INTERESTS

The authors declare that they have no conflicts of interest.

FUNDING

This research was funded by the National Natural Science Foundation of China (grant No. 31960277); the National Natural Science Foundation of China and Russian Foundation for Basic Research Cooperative Exchange Project (grant No. 32111530018, 21-54-53031); the Natural Science Foundation of Qinghai Province of China (grant No. 2018-ZJ-932Q); the Open Project of State Key Laboratory of Plateau Ecology and Agriculture, Qinghai University (grant No. 2017-ZZ-21). Dr. Wen Wang was supported by "1000Talent" programs of Qinghai Province.

AUTHOR CONTRIBUTIONS

WW conceived and designed the study. WW, and KS critically revised the manuscript. YL, FW, XLG, and LLZ performed the experiments, analyzed the data. All authors read and approved the final manuscript.

REFERENCES

- Takekawa JY, Heath SR, Douglas DC, Perry WM, Javed S, Newman SH, Suwal RN, Rahmani AR, Choudhury BC, Prosser DJ, Yan BP, Hou YS, Batbayar N, Natsagdorj T, Bishop CM, Butler PJ, Frappell PB, Milsom WK, Scott GR, Hawkes LA, Wikelski M: Geographic variation in bar-headed geese *Anser indicus*: Connectivity of wintering areas and breeding grounds across a broad front. *Wildfowl*, 59, 100-123, 2009.
- Zhang Y, Hao M, Takekawa JY, Lei F, Yan B, Prosser DJ, Douglas DC, Xing Z, Newman SH: Tracking the autumn migration of the bar-headed goose (*Anser indicus*) with satellite telemetry and relationship to environmental conditions. *Int J Zool*, 2011:323847, 2011. DOI: 10.1155/2011/323847
- Prins HHT, Wieren SE: Number, population structure and habitat use of bar-headed geese *Anser indicus* in Ladakh (India) during the brood-rearing period. *Acta Zoologica Sinica*, 50, 738-744, 2004.
- Liu D, Zhang G, Li F, Ma T, Lu J, Qian F: A revised species population estimate for the bar-headed goose (*Anser indicus*) *Avian Res*, 8:7, 2017. DOI: 10.1186/s40657-017-0064-7
- Hawkes LA, Balachandran S, Batbayar N, Butler PJ, Frappell PB, Milsom WK, Tseevenmyadag N, Newman SH, Scott GR, Sathiyaselvam P, Takekawa JY, Wikelski M, Bishop CM: The trans-Himalayan flights of bar-headed geese (*Anser indicus*). *Proc Natl Acad Sci USA*, 108 (23): 9516-9519, 2011. DOI: 10.1073/pnas.1017295108
- Hawkes LA, Balachandran S, Batbayar N, Butler PJ, Chua B, Douglas DC, Frappell PB, Hou Y, Milsom WK, Newman SH, Prosser DJ, Sathiyaselvam P, Scott GR, Takekawa JY, Natsagdorj T, Wikelski M, Witt MJ, Yan B, Bishop CM: The paradox of extreme high-altitude migration in bar-headed geese *Anser indicus*. *Proc R Soc B*, 280 (1750): 20122114, 2013. DOI: 10.1098/rspb.2012.2114
- Swan LW: Goose of the Himalayas. *Nat Hist*, 79, 68-75, 1970.
- Ward S, Bishop CM, Woakes AJ, Butler PJ: Heart rate and the rate of oxygen consumption of flying and walking barnacle geese (*Branta leucopsis*) and bar-headed geese (*Anser indicus*) *J Exp Biol*, 205 (21): 3347-3356, 2002.
- Butler PJ: High fliers: The physiology of bar-headed geese. *Comp Biochem Physiol A Mol Integr Physiol*, 156 (3): 325-329, 2010. DOI: 10.1016/j.cbpa.2010.01.016
- Scott GR, Hawkes LA, Frappell PB, Butler PJ, Bishop CM, Milsom WK: How bar-headed geese fly over the Himalayas. *Physiology*, 30 (2): 107-115, 2015. DOI: 10.1152/physiol.00050.2014
- Scott GR, Milsom WK: Control of breathing and adaptation to high altitude in the bar-headed goose. *Am J Physiol Regul Integr Comp Physiol*, 93 (1): R379-R391, 2007. DOI: 10.1152/ajpregu.00161.2007
- Scott GR: Elevated performance: The unique physiology of birds that fly at high altitudes. *J Exp Biol*, 214 (15): 2455-2462, 2011. DOI: 10.1242/jeb.052548
- Black CP, Tenney SM: Oxygen transport during progressive hypoxia in high-altitude and sea-level waterfowl. *Respir Physiol*, 39 (2): 217-239, 1980. DOI: 10.1016/0034-5687(80)90046-8
- McCracken KG, Barger CP, Sorenson MD: Phylogenetic and structural analysis of the HbA (alphaA/betaA) and HbD (alphaD/betaA) hemoglobin genes in two high-altitude waterfowl from the Himalayas and the Andes: bar-headed goose (*Anser indicus*) and Andean goose (*Chloephaga melanoptera*). *Mol Phylogenet Evol*, 56 (2): 649-658, 2010. DOI: 10.1016/j.ympev.2010.04.034
- Scott GR, Egginton S, Richards JG, Milsom WK: Evolution of muscle phenotype for extreme high altitude flight in the bar-headed goose. *Proc Biol Sci*, 276 (1673): 3645-3653, 2009. DOI: 10.1098/rspb.2009.0947
- Jendroszek A, Malte H, Overgaard CB, Beedholm K, Natarajan C, Weber RE, Storz JF, Fago A: Allosteric mechanisms underlying the adaptive increase in hemoglobin-oxygen affinity of the bar-headed goose. *J Exp Biol*, 221 (Pt 18): jeb185470, 2018. DOI: 10.1242/jeb.185470
- Natarajan C, Jendroszek A, Kumar A, Weber RE, Tame JRH, Fago A, Storz JF: Molecular basis of hemoglobin adaptation in the high-flying bar-headed goose. *PLOS Genet*, 14(4):e1007331, 2018. DOI: 10.1371/journal.pgen.1007331
- Bishop CM, Spivey RJ, Hawkes LA, Batbayar N, Chua B, Frappell PB, Milsom WK, Natsagdorj T, Newman SH, Scott GR, Takekawa JY, Wikelski M, Butler PJ: The roller coaster flight strategy of bar-headed geese conserves energy during Himalayan migrations. *Science*, 347 (6219): 250-254, 2015. DOI: 10.1126/science.1258732
- Chevron ZA, Brumfield RT: Genomic insights into adaptation to high-altitude environments. *Heredity*, 108 (4): 354-361, 2012. DOI: 10.1038/hdy.2011.85
- Zhang G: Bird sequencing project takes off. *Nature*, 522:34, 2015. DOI: 10.1038/522034d
- Wang W, Wang F, Hao R, Wang A, Sharshov K, Druzyaka A, Lancuo Z, Shi Y, Feng S: First de novo whole genome sequencing and assembly of the bar-headed goose. *PeerJ*, 8:e8914, 2020. DOI: 10.7717/peerj.8914
- Hao Y, Xiong Y, Cheng Y, Song G, Lei F: Comparative transcriptomics of 3 high-altitude passerine birds and their low-altitude relatives. *Proc Natl Acad Sci USA*, 116 (24): 11851-11856, 2019. DOI: 10.1073/pnas.1819657116
- Kim D, Langmead B, Salzberg SL: HISAT: A fast spliced aligner with low memory requirements. *Nat Methods*, 12 (4): 357-360, 2015. DOI: 10.1038/nmeth.3317
- Liao Y, Smyth GK, Shi W: Feature counts: An efficient general purpose program for assigning sequence reads to genomic features. *Bioinformatics*, 30 (7): 923-930, 2014. DOI: 10.1093/bioinformatics/btt656
- Love MI, Huber W, Anders S: Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome Biol*, 15 (12): 550, 2014. DOI: 10.1186/s13059-014-0550-8
- Yu G, Wang LG, Han Y, He QY: ClusterProfiler: An R package for comparing biological themes among gene clusters. *OMICS*, 16 (5): 284-287, 2012. DOI: 10.1089/omi.2011.0118
- Livak KJ, Schmittgen TD: Analysis of relative gene expression data using real-time quantitative PCR and the 2^{-ΔΔCT} method. *Methods*, 25 (4): 402-408, 2001. DOI: 10.1006/meth.2001.1262
- Qi X, Zhang Q, He Y, Yang L, Zhang X, Shi P, Yang L, Liu Z, Zhang F, Liu F, Liu S, Wu T, Cui C, Ouzhuluobu, Bai C, Baimakangzhuo, Han J, Zhao S, Liang C, Su B: The transcriptomic landscape of yaks reveals molecular pathways for high altitude adaptation. *Genome Biol Evol*, 11 (1): 72-85, 2019. DOI: 10.1093/gbe/evy264
- Dohn TE, Waxman JS: Distinct phases of Wnt/β-catenin signaling direct cardiomyocyte formation in zebrafish. *Dev Biol*, 361 (2): 364-376, 2012. DOI: 10.1016/j.ydbio.2011.10.032
- Deb A: Cell-cell interaction in the heart via Wnt/β-catenin pathway after cardiac injury. *Cardiovasc Res*, 102 (2): 214-223, 2014. DOI: 10.1093/cvr/cvu054
- Williams C, Sullivan K, Black LD: Partially digested adult cardiac extracellular matrix promotes cardiomyocyte proliferation *in vitro*. *Adv Healthc Mater*, 4 (10): 1545-1554, 2015. DOI: 10.1002/adhm.201500035
- Daquinag AC, Gao Z, Fussell C, Sun K, Kolonin MG: Glycosaminoglycan modification of decorin depends on MMP14 activity and regulates collagen assembly. *Cells*, 9(12):2646, 2020. DOI: 10.3390/cells9122646