

Evaluation for Some Bacterial and Viral Abortions of Dairy Cattle Farms in Burdur District of Turkey

Dilek OZTURK * 
Sibel HASIRCIOGLU **

Mehmet KALE **
Hulya TURUTOGLU *

Faruk PEHLIVANOGLU *

* Mehmet Akif Ersoy University, Faculty of Veterinary Medicine, Department of Microbiology, TR-15100 Burdur - TURKEY

** Mehmet Akif Ersoy University, Faculty of Veterinary Medicine, Department of Virology, TR-15100 Burdur - TURKEY

Makale Kodu (Article Code): KVFD-2011-5381

Summary

In this study, the presence of antibodies to *Brucella abortus*, *Chlamydophila abortus*, *Coxiella burnetii*, Bovine herpesvirus-1 (BHV-1), Bovine viral diarrhoea virus (BVDV) and Bovine herpesvirus-4 (BHV-4) were investigated in dairy cattle herds with abortion history in Burdur province, Southwest region of Turkey. The blood samples were collected from 932 dairy cattle ≥ 2 years of age from 10 herds not vaccinated against for mentioned infections and all of the serum samples were tested for *B. abortus* by ELISA. Seronegative herds for *B. abortus* were investigated for *C. burnetii* and *C. abortus*. While seropositivity for *B. abortus* and *C. burnetii* were detected 25.3% (236/932) and 10.2% (19/186), antibodies against *B. abortus* was not detected. Seronegative herds for *B. abortus*, *C. burnetii* and *C. abortus* were analysed for antibodies to BHV-1, BVDV and BHV-4 and seropositivity were found as 43.5% (40/92), 81.5% (75/92), 42.4% (39/92), respectively. BVDV antigen was determined in 2.2% (2/92) of the samples. As a result, we determined that *C. abortus* is not an important agent causing abortion in dairy herds in Burdur, but the seropositivity of *B. abortus*, *C. burnetii*, BHV-1, BVDV and BHV-4 are high. Thus, the herds should be tested for these infections regularly and control measures must be taken.

Keywords: Abortion, ELISA, Dairy cattle, *Brucella abortus*, *Chlamydophila abortus*, *Coxiella burnetii*, Bovine herpesvirus-1, Bovine viral diarrhoea virus, Bovine herpesvirus-4

Burdur Yöresinde Süt Sığırı İşletmelerinin Bazı Bakteriyel ve Viral Abortlar Yönünden Değerlendirilmesi

Özet

Bu çalışmada, Türkiye'nin güneybatı bölgesinde yer alan Burdur ili'nde yavru atma problemleri süt sığırlarında *Brucella abortus*, *Chlamydophila abortus*, *Coxiella burnetii*, Bovine herpesvirus-1 (BHV-1), Bovine viral diarrhoea virus (BVDV) ve Bovine herpesvirus-4 (BHV-4)'e karşı antikorların varlığını araştırıldı. Kan örnekleri araştırılan enfeksiyonlara karşı aşılanmamış 10 sürüden 2 yaş ve üzeri 932 hayvandan toplandı. Serumlar önce *B. abortus* yönünden ELISA ile test edildi. *B. abortus* negatif bulunan serum örnekleri *C. burnetii* ve *C. abortus*'a karşı antikorların varlığını yönünden araştırıldı. *B. abortus* ve *C. burnetii* için pozitiflik sırasıyla %25.3 (236/932) ve %10.2 (19/186) belirlenirken, *C. abortus* antikorları tespit edilemedi. *B. abortus*, *C. burnetii* ve *C. abortus* negatif bulunan sürülerde BHV-1, BVDV ve BHV-4'e karşı antikorların varlığını araştırıldı. Seropozitiflik bu hastalıklar için sırasıyla %43.5 (40/92), %81.5 (75/92), %42.4 (39/92) olarak belirlendi. BVDV antijen ise örneklerin %2.2'sinde tespit edildi. Sonuç olarak, *C. abortus*'un Burdur'da süt sığırı işletmelerinde aborta neden olan önemli bir ajan olmadığı, *B. abortus*, *C. burnetii*, BHV-1, BVDV ve BHV-4'ün seropozitifliğinin yüksek olduğu belirlendi. Bu nedenle, sürülerin bu enfeksiyonlar yönünden düzenli olarak test edilmesi ve gerekli kontrol önlemlerinin alınması gerektiği sonucuna varıldı.

Anahtar sözcükler: Abort, ELISA, Süt sığırı, *Brucella abortus*, *Chlamydophila abortus*, *Coxiella burnetii*, Bovine herpesvirus-1, Bovine viral diarrhoea virus, Bovine herpesvirus-4

INTRODUCTION

Bovine abortion is a significant cause of economic losses in dairy cattle herds¹. Although the cause of abortion depends on several factors including idiopathic, metabolic

or hormonal abnormalities, nutritional deficiencies, trauma and infections, the infectious agents including bacterial, viral, fungal and protozoan agents are the most important



İletişim (Correspondence)



+90 248 2132062



sedilek@yahoo.com

factors associated with abortions among dairy herds worldwide ^{2,3}. The infections causing abortion may vary, presumably due to climate, production types, management practices and geographic factors in different regions ⁴. Effective preventive measures for abortions require not only prompt and accurate diagnosis, but also understanding of the multicausal factors involved ⁵.

The aim of this study was to investigate *Brucella abortus*, *Chlamydomphila abortus*, *Coxiella burnetii*, Bovine herpesvirus-1, Bovine viral diarrhoea virus and Bovine herpesvirus-4 infections of dairy cattle herds with abortion history.

MATERIAL and METHODS

Samples

This study was conducted in 10 dairy cattle herds with abortion history in Burdur province, Southwest region of Turkey. Blood samples were collected from 932 dairy cattle ≥ 2 years of age. Blood samples were taken into tubes without anticoagulant. After clotting, the tubes were centrifuged, and the serum samples were separated and kept at -20°C until used. All of the serum samples were tested for *B. abortus* by ELISA. The seronegative samples for *B. abortus* were investigated for *C. abortus* and *C. burnetii* by ELISA. After, seronegative herds for *B. abortus*, *C. burnetii* and *C. abortus* were analysed for antibodies to BHV-1, BVDV and BHV-4. Leucocyte were obtained from blood samples, taken into tubes with EDTA, by centrifugation at 2.500 rpm/25 min. and stored in deep freezer at -20°C till used. All of animals were not vaccinated against BHV-1, BVDV, BHV-4 and *B. abortus* before.

Antibody Detections in Sera by ELISA

Presence of antibodies against *B. abortus*, *C. burnetii* and *C. abortus* (Institute Pourquier, France), BHV-1, BVDV and BHV-4 (Bio-X Diagnostics, Belgium) were investigated by ELISA. The tests were carried out according to the kit procedure.

BVDV Antigen - ELISA

The kit of BVD/MD Antigen Mix Screening ELISA (Institute Pourquier, France) was used to determine the existence of antigen in the leucocyte samples. The test was carried out according to the kit procedure.

RESULTS

The seropositivity for *B. abortus* was found in 25.3% (236/932) of the sera. While seropositivity for *C. burnetii* was found in 10.2% (19/186) of the samples, *C. abortus* was not detected in any of the sera (0/225). In seronegative herds for *B. abortus*, *C. burnetii* and *C. abortus*, seropositivity for BHV-1, BVDV and BHV-4 were found as 43.5% (40/92), 81.5% (75/92), and 42.4% (39/92), respectively (Table 1).

Antibodies rates of BHV-1 + BVDV + BHV-4, BHV-1 + BHV-4, BHV-1 + BVDV and BVDV + BHV-4 were 21.7% (20/92), 21.7% (20/92), 40.2% (37/92), 37% (34/92), respectively. Also, the rates at which cattle were detected for both BHV-1 + BVDV (antigen) + BHV-4 and BHV-1 + BVDV (antigen) were 1.1% (1/92) (Fig. 1).

BVDV antigen was only detected as 2.2% (2/92) in the leucocyte samples in aborted cattle.

Table 1. The rates of seropositivity for *B. abortus*, *C. burnetii*, *C. abortus*, BHV-1, BVDV, BHV-4 in dairy cattle herds with abortion history


Tablo 1. Abort hikayesi bulunan süt sığırı işletmelerinde *B. abortus*, *C. burnetii*, *C. abortus*, BHV-1, BVDV, BHV-4'ün seropozitiflik oranları

Bacterial and Viral Agents	Number of Serum Samples	Positive	%
<i>B. abortus</i>	932	236	25.3
<i>C. burnetii</i>	186	19	10.2
<i>C. abortus</i>	225	0	0
BHV-1	92	40	43.5
BVDV	92	75	81.5
BHV-4	92	39	42.4

Fig 1. Multiple infection rates of BHV-1, BVDV and BHV-4 in dairy cattle herds with abortion history

(Ab: Antibody, Ag: Antigen a: BHV-1 (Ab) + BVDV (Ab) + BHV-4 (Ab), b: BHV-1 (Ab) + BHV-4 (Ab), c: BHV-1 (Ab) + BVDV (Ab), d: BVDV (Ab) + BHV-4 (Ab), e: BHV-1 (Ab) + BVDV (Ab-/Ag+) + BHV-4 (Ab), f: BHV-1 (Ab) + BVDV (Ab-/Ag+)).

Şekil 1. Abort hikayesi bulunan süt sığırı sürülerinde BHV-1, BVDV and BHV-4 için çoklu enfeksiyon oranları



DISCUSSION

Abortion has caused serious economic losses worldwide. Regional differences in the prevalence of abortion-causing diseases can be a result of a variety of reasons such as methods of herd health management, climate, types of cattle operations, feed, vaccination programmes, geographic factors, number of pregnancies and sanitation^{1,4}. Brucellosis is also very common and an economically important zoonotic disease in Turkey^{6,7} and worldwide^{8,9}. In this study, the seropositivity for *B. abortus* was found to be 25.3% in dairy cattle herds with abortion history in Burdur province. The prevalence studies for *B. abortus* have been conducted in different parts of Turkey by researchers^{6,7,10}. In these studies⁶⁻⁹, prevalence have changed from 1.4% to 18% and quite low than this study. There is one study related with the seropositivity of brucellosis in dairy cattle herds with abortions problems in Burdur which reported the rate of brucellosis as 10.1% by Rose Bengal plate test (RBPT), and 6.8% by serum agglutination test (SAT) during 2007-2008¹¹. This study results was rather lower than present study. The reason of this may be due to higher sensitivity and specificity of ELISA compared to RBPT and SAT. Seroprevalance of brucellosis was reported in between 3.3-22.2% in other countries^{8,9}. We thought that these differences may originate from factors such as use of different detection methods, climate, vaccination programmes, sampling method of animals, intensification of animal movements due to the high potential of the region and lack of herd screening for *B. abortus*. This study supports researchers who state that prevalence of infections causing abortion may be affected by the mentioned factors^{1,4,12}.

Abortions caused by *Chlamydomphila* spp. in cattle are rather widespread worldwide and are generally associated with *C. abortus* infections. Because *Chlamydomphila* infections are not considered as a major threat to the cattle industry in Turkey, there are less studies of cattle than sheep or goats^{13,14}. In this study, the antibodies for *C. abortus* can not be detected in blood serum samples of cattle. In the South East part of Turkey, seroprevalance of *C. abortus* was determined to be 8.3% in aborted dairy cattle¹⁵. The seroprevalance for *Chlamydomphila* infection was also reported in between 0.4-57% in other countries by ELISA¹⁶⁻¹⁸. *Chlamydomphila* spp. was not responsible from abortion cases of cattle in this region, because positive serum samples were not detected.

In this study, the prevalence of *C. burnetii* was 10.2%, while studies conducted in different parts of Turkey have results ranging from 4.3% to 44.5% in cattle herds with abortion problems¹⁹⁻²¹. Özyer et al.¹⁹ reported that the seropositivity of *C. burnetii* was 44.5% in cattle with abortion history by complement fixation test (CFT). The seropositivity of *C. burnetii* was 22.6% in Erzurum²⁰ and 16.3% in 16 locations of Eastern Turkey²¹ by ELISA. In

Aydin, Kirkan et al.²² were found that *C. burnetii* was 4.3% in cattle by polymerase chain reaction (PCR). The prevalence of *C. burnetii* infection was found between 5% and 44.9% in other countries²³⁻²⁵. The seropositivity of *C. burnetii* was found in 39% of cattle by indirect fluorescence in Zimbabwe²³. Serbezov et al.²⁴ reported that the occurrence of *C. burnetii* was 5% to 31% in cattle by CFT in Bulgaria. The seropositivity was determined 44.9% in dairy cattle of Northern Italy by ELISA²⁵. These differences may be associated with the differences of sensitivity and specificity of diagnostic methods used for diagnosis of *C. burnetii* infections.

In the present study, the seropositivity of BHV-1, BVDV and BHV-4 was found to be 43.5%, 81.5% and 42.4%, respectively. The presence of BHV-1 antibodies detected in 25-60%^{4,26,27} and for BVDV were detected 75-80%^{28,29}, for BHV-4 were found 36-88%³⁰⁻³², in aborted cattle. However, there are also contradictory studies³³⁻³⁵. But, Kale et al.³⁶ investigated presence of BHV-1, BVDV and BHV-4 infections in 204 dairy cows 3 to 5 years of age that aborted at 5-7 months of gestation in same area³⁶. They found lower rates of BHV-1 (21.6%), BVDV (25.5%) and BHV-4 (29.4%) seropositivity than the current study. As a result, there has been an increase in the infections of BHV-1, BVDV and BHV-4 in the region. In Kars, Yildirim et al.³⁷ reports that the seropositivity of BHV-1 and BHV-4 were found 61.4%, 29.3% by ELISA and the presence of BVDV specific antibody was found 52.9% by virus neutralization test in aborted cattle. The seropositivity of BHV-1 and BVDV are higher than this study. Often, no obvious clinical signs are seen in herds with fetal losses due to BVD virus and abortions occurring days or several weeks following maternal infection. In the study, seroprevalance of BVDV antigen was detected as 2.2% in aborted cattle. Research has shown <10% of abortion is related to BVDV^{38,39}. The seroprevalance of BHV-1, BVDV and BHV-4 had higher levels than that of other studies which viral and bacterial mix in the study⁴⁰. However, there are also higher seroprevalance results⁴¹ than seen in the study. Results of this study indicate that BHV-1, BVDV and BHV-4 are a significant viral cause of abortion in the area. Bovine abortion can be caused by several pathogens and evidence of multiple agent involvement has been indicated⁴². Furthermore, in the research, while seroprevalance of BHV-1 + BVDV have the highest and 40.2%, BHV-1 + BVDV (negative antibody/positive antigen) + BHV-4 and BHV-1 + BVDV (negative antibody/positive antigen) was the lowest and 1.1%. In context, it is important to point out that in mixed cases it is extremely difficult to establish the actual agent responsible for the abortion²⁷.

In conclusion, we determined that seroprevalance of *B. abortus*, *C. burnetii*, BHV-1, BVDV and BHV-4 in dairy cattle farms in Burdur were higher than those in other parts of Turkey, but *C. abortus* is not a major agent causing abortion in cattle population in the region. Thus, the herd surveys should be performed regularly for such infections in the region.

REFERENCES

1. **Anderson ML, Blanchard PC, Barr BC, Hoffman RL:** A survey of causes of bovine abortion occurring in the San Joaquin Valley, California. *J Vet Diagn Invest*, 2, 283-287, 1990.
2. **Jamaluddin AA, Case JT, Hird DW, Blanchard PC, Peuroi JR, Anderson ML:** Dairy cattle abortion in California: Evaluation of diagnostic laboratory data. *J Vet Diagn Invest*, 8, 210-218, 1996.
3. **Corbellini LG, Pescador CA, Frantz F, Wunder E, Steffen D, Smith DR, Driemeier D:** Diagnostic survey of bovine abortion with special reference to *Neospora caninum* infection: Importance, repeated abortion and concurrent infection in aborted fetuses in Southern Brazil. *Vet J*, 172, 114-120, 2006.
4. **Anderson ML:** Diagnosis of infectious causes of bovine abortion. *American Association of Bovine Practitioners. Conference, The AABP Proceedings*, Sep. 23-Sep. 25, 37, 90-96. USA, 2004.
5. **Khodakaram-Tafti A, Ikede BO:** A retrospective study of sporadic bovine abortions, stillbirths, and neonatal abnormalities in Atlantic Canada, from 1990 to 2001. *Can Vet J*, 46, 635-637, 2005.
6. **Denizli AN, Gokcen S, Ergun A:** Ege bölgesi sığırlarında brucella enfeksiyonunun ELISA testi ile insidensinin saptanması. *Hayvan Aşılari Kontrol Merkezi Müdürlüğü*, 16, 48-57, 1992.
7. **Iyisan S, Akmaz O, Duzgun SG, Ersoy Y, Eskiüzümlü S, Guler L, Gunduz K, Isik N, Icyerioglu AK, Kalender H, Karaman Z, Kucukayan U, Ozcan C, Seyitoglu S, Tuna I, Tunca T, Ustunakin K, Yurtalan S:** Türkiye'de sığır ve koyunlarda Brucellosis'in epidemiyolojisi. *Pendik Vet Mikrobiyol Derg*, 31, 21-75, 2000.
8. **Trangadia B, Rana SK, Mukherjee F, Srinivasan VA:** Prevalence of brucellosis and infectious bovine rhinotracheitis in organized dairy farms in India. *Trop Anim Health Prod*, 42, 203-207, 2010.
9. **Ibrahim N, Belihu K, Lobago F, Bekana M:** Sero-prevalence of bovine brucellosis and its risk factors in Jimma zone of Oromia Region, South-western Ethiopia. *Trop Anim Health Prod*, 42, 35-40, 2010.
10. **Demirozu K, Celik M, Iyisan S:** Kars ilinde brusellosisin sero-epidemiyolojisi. *Pendik Vet Mikrobiyol Derg*, 24, 123-132, 1993.
11. **Pehlivanoglu F, Ozturk D, Gunlu S, Guldali Y, Turutoglu H:** Prevalence of Brucellosis in dairy herds with abortion problems. *Kafkas Univ Vet Fak Derg*, 17 (4): 615-620, 2011.
12. **Peter AT:** Abortions in dairy cows: New insights and economic impact. *Adv Dairy Sci Tech*, 12, 233-244, 2000.
13. **Turutoglu H, Iyisan AS, Duru A, Altinel C:** Koyunlarda *Chlamydia psittaci* enfeksiyonunun mikrokomplement fikzasyon testi ile saptanması. *Pendik Vet Mikrobiyol Derg* 26, 67-78, 1995.
14. **Ozturk M, Nadas UG, Turutoglu H:** İstanbul bölgesindeki koyunlarda *Chlamydia psittaci* enfeksiyonunun prevalansının saptanması. *Pendik Vet Mikrobiyol Derg*, 29, 72-80, 1998.
15. **Gokce HI, Kacar C, Genc O, Sozmen M:** Seroprevalance of *Chlamydia abortus* in aborting ewes and dairy cattle in the North east part of Turkey. *B Vet I Pulawy*, 51, 9-13, 2007.
16. **Godin AC, Bjorkman C, Englund S, Johansson KE, Niskanen R, Alenius S:** Investigation of *Chlamydia* spp. in dairy cows with reproductive disorders. *Acta Vet Scan*, 50, 39, 2008.
17. **Vlahovic K, Dovc A, Zupancic Z, Pavlaki M, Jercici J, Marina P, Jure J:** Comparison of serological procedures for diagnosis of infection with *Chlamydia* spp. in bovines. *Vet Arhiv*, 71, 67-379, 2001.
18. **Greco G, Corrente M, Buonavoglia D, Campanile G, Di Palo R, Martella V, Bellacicco AL, Abramo MD, Buonavoglia C:** Epizootic abortion related to infections by *Chlamydia abortus* and *Chlamydia pecorum* in water buffalo (*Bubalus bubalis*). *Therogenology*, 69, 1061-1069, 2008.
19. **Ozyer M, Mirioglu M, Koksall F:** Çukurova bölgesinde yaşayan insan ve hayvanlarda Q-Fever enfeksiyonu insidansının komplement fikzasyon testi ile karşılaştırılması. *Pendik Hayvan Hast Mer Aras Enst Derg*, 21, 28-39, 1990.
20. **Seyitoglu S, Ozkurt Z, Dinler U, Okumus B:** The seroprevalance of Coxiellosis in farmers and cattle in Erzurum district in Turkey. *Turk J Vet Anim Sci*, 30, 71-75, 2006.
21. **Ceylan E, Berktaş M, Keles I, Agaoglu Z:** Seroprevalance of Q Fever in cattle and sheep in the East of Turkey. *Asian J Anim Adv*, 4, 114-121, 2009.
22. **Kirkan S, Kaya O, Tekbiyik S, Parin U:** Detection of *Coxiella burnetii* in cattle by PCR. *Turk J Vet Anim Sci*, 32, 215-220, 2008.
23. **Kelly PJ, Matthewman LA, Mason, PR, Raoult D:** Q fever in Zimbabwe. A review of the disease and the results of a serosurvey of humans, cattle, goats and dogs. *S Afr Med J*, 83, 21-25, 1993.
24. **Serbezov V, Kazar J, Novkirishki V, Gatcheva N, Kovacova E and Voynova V:** Q fever in Bulgaria and Slovakia. *Emerg Infect Dis*, 5, 388-394, 1999.
25. **Cabassi CS, Taddei S, Donofrio G, Ghidini F, Piancastelli C, Flammini CF, Cavirani S:** Association between *Coxiella burnetii* seropositivity and abortion in dairy cattle of Northern Italy. *New Microbiol*, 29, 211-214, 2006.
26. **Medici KC, Alfieri AA, Alfieri AF:** Prevalence of neutralizing antibodies against bovine herpesvirus Type 1, due natural infection, in herds with reproductive problems. *Ciência Rural*, 30, 347-350, 2000.
27. **Dechicha A, Gharbi S, Kebbal S, Chatagnon G, Tainturier D, Ouzrout R, Guetarni D:** Serological survey of etiological agents associated with abortion in two Algerian dairy cattle breeding farms. *J Vet Med Anim Health*, 2, 1-5, 2010.
28. **Rüfenacht J, Schaller P, Audige L, Knutti B, Kupfer U:** The effect of infection with bovine viral diarrhoea virus on the fertility of swiss dairy cattle. *Therogenology*, 56, 199-210, 2001.
29. **Hall CA, Reichel MP, Ellis JT:** Neospora abortions in dairy cattle: Diagnosis, mode of transmission and control. *Vet Parasitol*, 128, 231-241, 2005.
30. **Naeem K, Goyal SM, Werdin RE:** Prevalence of bovid herpesvirus and its antibody in cattle in Minnesota. *Am J Vet Res*, 50, 1931-1935, 1989.
31. **Frazier KS, Baldwin CA, Pence M, West J, Bernard J, Liggett A, Miller D, Hines II ME:** Seroprevalence and comparison of isolates of endometriotropic Bovine Herpesvirus-4. *J Vet Diagn Invest*, 14, 457-462, 2002.
32. **Kale M, Ata A, Kocamüftüoğlu M, Hasircioğlu S:** Bovine herpesvirus type 4 (BHV-4) infection in relation to fertility in repeat breeder dairy cows. *Acta Vet Beo*, 61, 13-19, 2011.
33. **Mineo TWP, Alenius S, Naslund K, Montassier HJ, Bjorkman C:** Distribution of antibodies against *Neospora caninum*, BVDV and BHV-1 among cows in Brazilian dairy herds with reproductive disorders. *Rev Bras Parasitol Vet*, 15, 188-192, 2006.
34. **Bilge Dagalp S, Demir AB, Gungor E, Alkan F:** The seroprevalence of Bovine Herpes Virus Type 4 (BHV4) infection in dairy herds in Turkey and possible interaction with reproductive disorders. *Rev Med Vet-Toulouse*, 158, 201-205, 2007.
35. **Dhami JS, Dwivedi PN, Ramneek DD, Maiti NK, Oberoi MS:** Diagnosis of bovine herpesvirus-1 (BHV-1) infection in aborting cattle and buffaloes by virus isolation and polymerase chain reaction (PCR). *Indian J Anim Sci*, 78, 1235-1237, 2008.
36. **Kale M, Hasircioğlu S, Acar A:** Gebeliginin 5-7. ayları arasında abort yapmış süt sığırlarında Bovine Herpesvirus-1, Bovine Viral Diarrhoea Virus ve Bovine Herpesvirus-4 enfeksiyonlarının araştırılması. *VIII. Ulusal Veteriner Mikrobiyoloji Kongresi (Uluslar Arası Katılımlı)*, 218-219, 07-09 Ekim, Kampus - Van, Türkiye, 2008.
37. **Yıldırım Y, Yılmaz V, Kalaycıoğlu AT, Dağalp SB, Farajı Majarashın AR, Çelebi Ö, Akça D:** An investigation of a possible involvement of BVDV, BHV-1 and BHV-4 infections in abortion of dairy cattle in Kars district of Turkey. *Kafkas Univ Vet Fak Derg*, 17, 879-883, 2011.
38. **Meyling A, Houe H, Jensen AM:** Epidemiology of bovine virus diarrhoea virus. *Rev Sci Tech Off Int Epiz*, 9, 75-93, 1990.
39. **Heuer C, Healy A, Zerbini C:** Economic effects of exposure to bovine viral diarrhoea virus on dairy herds in New Zealand. *J Dairy Sci*, 90, 5428-5438, 2007.
40. **Norton JH, Tranter WP, Campbell RSF:** A farming systems study of abortion in dairy cattle on the Atherton Tableland: 2. The pattern of infectious diseases. *Aust Vet J*, 66, 163-167, 2008.
41. **Escamilla HP, Juan Martinez MJ, Mario Medina C, Elizabeth Morales S:** Frequency and causes of infectious abortion in a dairy herd in Queretaro, Mexico. *Can J Vet Res*, 71, 314-317, 2007.
42. **Caldow GL:** Bovine abortion outbreak associated with *Neospora* and other infectious agents. *Vet Rec*, 142, 118-119, 1998.