

# Effect of Energy Sources and Levels on Caecal Microbial Population, Jejunal Morphology, Gene Expression of Jejunal Transporters (SGLT1, FABP) and Performance of Broilers Under Heat Stress

Asghar GHAHREMANI <sup>1</sup> Ali Asghar SADEGHI <sup>1</sup>✍ Saeed HESARAKI <sup>2</sup>  
Mohammad CHAMANI <sup>1</sup> Parvin SHAWRANG <sup>3</sup>

<sup>1</sup> Department of Animal Science, Science and Research Branch, Islamic Azad University, Tehran, IRAN

<sup>2</sup> Department of Veterinary Medicine, Science and Research Branch, Islamic Azad University, Tehran, IRAN

<sup>3</sup> Nuclear Agriculture Research School, Nuclear Science and Technology Research Institute, Atomic Energy Organization of Iran, Karaj, IRAN

Article Code: KVFD-2016-16899 Received: 09.10.2016 Accepted: 10.12.2016 Published Online: 12.12.2016

## Citation of This Article

Ghahremani A, Sadeghi AA, Hesarakı S, Chamani M, Shawrang P: Effect of energy sources and levels on caecal microbial population, jejunal morphology, gene expression of jejunal transporters (SGLT1, FABP) and performance of broilers under heat stress. *Kafkas Univ Vet Fak Derg*, 23 (3): 415-422, 2017. DOI: 10.9775/kvfd.2016.16899

## Abstract

The present study was conducted to evaluate the effects of energy sources and levels on microbial population, jejunal morphology, gene expression of glucose transporter (SGLT1), fatty acid binding protein (FABP) and performance in broilers under heat stress. In a completely randomized design, 600 one-day-old Cobb broiler chickens were assigned to five dietary treatments and four replicates. Chicks were fed diet based on corn as main energy source and energy level based on Cobb standard considered as control (C), corn based diet with 3% lesser energy than control (T1), corn based diet with 6% lesser energy than control (T2), corn and soybean oil based diet according to Cobb standard (T3), corn and soybean oil based diet with 3% upper energy than control (T4). Temperature was increased to 34°C for 8 hours daily from day 21 to 41 to induce heat stress. Chickens in T3 and T4 had higher *Lactobacillus* population and lower *Escherichia coli* population than C group (P<0.05). Chickens in T1 and T2 had shorter jejunal villi, deeper crypts, and lower villus height: crypt depth than those fed C, T3 and T4 diets (P<0.05). There were no significant differences among treatments for gene expression of both nutrient transporters (P>0.05). Chickens in T3 and T4 had higher weight gain compared to C, T1 and T2 (P<0.05). Feed intake in T3 was lower than C, consequently, feed conversion ratio of chicks fed T3 was better than C group (P<0.05). In conclusion, replacement a part of dietary energy source with soybean oil might improve intestinal parameters and performance of broilers under heat stress.

**Keywords:** Energy, Lipid, Intestinal morphology, Transporter, Broiler

## Isı Stresi Altında Enerji Kaynakları ve Seviyelerinin Sekum Mikrobiyal Popülasyonuna, Jejunum Morfolojisine, Jejunal Transporterlerin (SGLT1, FABP) Gen Ekspresyonuna ve Broiler Performansına Etkileri

## Özet

Bu çalışma; enerji kaynakları ve seviyelerinin ısı stresi altında mikrobiyal popülasyona, jejunum morfolojisine, glukoz transporteri (SGLT1) ve yağ asitleri bağlayıcı protein (FABP) gen ekspresyonlarına ve broiler performansına etkilerini araştırmak amacıyla yürütülmüştür. Rastgele örneklemeyle, 600 adet 1 günlük Cobb broiler civiv, 4 tekrar olmak üzere 5 farklı beslenme uygulanmasına alındı. Civivler ana enerji kaynağı olarak mısıra dayalı diyetle beslendi. Kontrol grubuna (C) bazal seviyede Cobb standardına göre yem verilirken, T1 grubuna kontrole göre %3 daha düşük enerjili yem, T2 grubuna kontrole göre %6 daha düşük enerjili yem, T3 grubuna Cobb standardına göre mısır ve soya fasulyesi yağı tabanlı diyet ve T4 grubuna kontrole göre %3 daha fazla enerjili mısır ve soya fasulyesi yağı tabanlı diyet uygulandı. Isı stresi oluşturmak amacıyla 21. günden 41. güne kadar günde 8 saat boyunca sıcaklık 34°C'ye çıkarıldı. T3 ve T4 grubundaki civivlerde kontrol grubuna oranla *Lactobacillus* popülasyonu daha yüksek iken *Escherichia coli* popülasyonu daha düşüktü (P<0.05). T1 ve T2 grubundaki civivlerde C, T3 ve T4 grubundakilere kıyasla daha kısa jejunal villi, daha derin kriptler ve daha düşük villus yüksekliği:derinliği belirlendi (P<0.05). Her iki besin transporteri için de gen ekspresyonlarında farklı uygulamalar için herhangi bir fark gözlemlenmedi (P>0.05). T3 ve T4 grubundaki civivlerde C, T1 ve T2 gruplarına kıyasla daha yüksek kilo kazanımı belirlendi (P<0.05). Kontrol grubuna oranla T3 grubundaki civivlerin yem tüketimi daha düşük ve dolayısıyla yem konversiyon oranı daha iyiydi (P<0.05). Sonuç olarak, diyetteki enerji kaynağının bir bölümünün soya fasulyesi yağı ile değiştirilmesinin bağırsak parametrelerini ve ısı stresi altındaki broilerin performansını iyileştirebileceği belirlendi.

**Anahtar sözcükler:** Enerji, Yağ, Bağırsak morfolojisi, Transporter, Broiler



## İletişim (Correspondence)



+98 919 5579663



a.sadeghi@srbiau.ac.ir

## INTRODUCTION

Heat stress is considered as a major problem in poultry production on subtropical and tropical regions, where major broiler farms exist<sup>[1]</sup>. Heat stress occurs when a negative balance between the amount of heat energy produced by the animal and the amount of energy flowing from the broiler body to environment exists. This condition affects negatively the health and performance of broilers<sup>[2]</sup>. In terms of changes in metabolic pathways<sup>[3]</sup>, microbial population in the small intestine<sup>[4]</sup>, hormonal levels<sup>[5]</sup>, low feed consumption, body weight gain, high feed conversion ratio (FCR)<sup>[6]</sup> and damage to small intestine structure<sup>[7,8]</sup>. These changes not only could affect nutrient absorption, but also the merit of substrates for metabolism<sup>[9,10]</sup>.

Intestinal bacterial flora are considered important for priming and maintaining an active immune system<sup>[11]</sup>. The microbiota composition can be affected by environmental factors, genetics and substrate availability within the gut<sup>[12]</sup>. Development of the intestinal villi in the early chicken's life could increase efficiency of nutrient utilization and enhance the growth performance. Furthermore, an increase in villi height may increase the intestinal surface area and nutrient absorption<sup>[13]</sup>. Mitchell and Carlisle<sup>[14]</sup> reported that chronic heat stress decreased small intestinal villus height and wet and dried small intestinal weight of birds.

The process of absorption of carbohydrates into the enterocytes of the small intestine is mediated by sugar transporters, such as sodium-glucose transporter 1 (SGLT-1)<sup>[15]</sup>, which diffuses monosaccharides into the extracellular fluid and then into the blood<sup>[16]</sup>. Triglycerides (TGs) are broken into glycerol and free fatty acids and then absorbed in the intestine and transported across the apical membrane of the enterocytes. Fatty-acid-binding proteins (FABPs) are intracellular lipid chaperones that transport lipids to a specific component in the cell<sup>[17]</sup>. Nutrient transporters in the small intestine are responsible for dietary nutrient assimilation; therefore, heat stress-related changes in the expression of these transporters affect the availability of nutrients and energy to the animal for growth and development. However, the effects of heat stress on the expression of nutrient transporters in the small intestine of broiler chickens are unclear. Some studies showed that starvation stress caused the increased expression of sodium glucose co-transporter 1 (SGLT1) mRNA in the small intestine of chickens and rats<sup>[18,19]</sup>.

Wang et al.<sup>[20]</sup> showed that high apparent metabolizable energy in the diets fed to broilers improved their feed conversion ratio, emphasizing the potential role of nutrient density as an important factor that may affect animal intestine development. Chickens fed a higher nutrient density diet grow faster throughout all growing phases<sup>[21]</sup>. There are evidences that a higher dietary fat content contributes to improved heat tolerance in broiler chickens<sup>[22]</sup>. Zulkifli et al.<sup>[23]</sup> reported that providing diets containing

high levels of palm oil enhanced growth performance and survivability of heat-stressed broiler chickens.

To our knowledge, there was no report concerning the effect of energy sources and levels on intestinal transporters gene expression, morphology and microbiology, especially in the context of heat stress condition. It was hypothesized that in the heat stress condition, lipid addition with higher energy level to diet could improve the small intestine morphology, gene expression of transporter and beneficiary bacteria population compared to diet containing main energy source from carbohydrate. Therefore, the main objective was to evaluate the effects of energy sources and levels on microbial population, small intestine morphology, gene expression of glucose transporter (SGLT1), fatty acid binding protein (FABP) and performance in broiler chickens under heat stress conditions.

## MATERIAL And METHODS

### Chickens Management

All animal procedures were approved by the Animal Care Committee of the Animal Sciences Research Institute of Iran. The use of broilers in this study was approved by the Animal Care Committee (Protocol 17-16-5-10938; 90-11-15).

A total of 600 one-day-old Cobb 500 male broiler chicks with an average weight of  $39 \pm 0.50$ g was obtained from a local hatchery and randomly allocated to 20 floor pens (200 cm × 180 cm) covered with pine shaving. Chicks were randomly assigned to five dietary treatments with four replicates and 30 chicks per each. Chicks were raised under environmentally controlled conditions and lighting program based on Cobb 500 broiler guides (Cobb Broiler Management Guide, 2010), except temperature. Feed intake and live body weight were recorded in the beginning and at the end of experiment, and the feed conversion ratio was then calculated. Dead chicks were collected daily and weighed at the time of carcass removal; carcass weights were included in the feed conversion ratio calculations.

### Experimental Design

Dietary treatments were: control group (C) which broilers fed diet with main energy from corn and energy level; T1: broilers fed diet with main energy from corn and 3% lesser energy; T2: broilers fed diet with main energy from corn and 6% lesser energy; T3: broilers fed diet with main energy from corn and soy oil and energy level T4: broilers fed diet with main energy from corn and soy oil and 3% upper energy. The experimental diets were formulated based on Cobb standard (Cobb instruction manual, 2012) (Table 1). Washed sand as filler was used to balance for dietary metabolizable energy levels. Chicks were fed a starter diet from day 1 to 10, grower diet from day 11 to 28, followed by a finisher diet from day 29 to 42 of age. Feed

**Table 1.** Composition (measured in %) of the experimental diets for broiler chickens

Ingredients	Starter (0-10 days old)					Grower (11-28 days old)					Finisher (29-42 days old)				
	C	T1	T2	T3	T4	C	T1	T2	T3	T4	C	T1	T2	T3	T4
Corn	63.35	63.33	62.13	58.24	54.26	69.22	68.8	67.7	65.31	59.31	70.18	71.47	71.5	65.07	62.4
Soybean meal	22.57	28.86	31.48	31.90	32.21	18	23.43	26.2	24	28.45	19.3	20	24.2	25.8	25
Soybean oil	-----	-----	-----	2.5	4.14	-----	-----	-----	2	5	-----	-----	-----	3.5	5
Corn gluten meal	9.17	3	-----	2.7	4.7	8.2	3.24	-----	4.2	2.9	6.23	4.2	-----	1.5	3.5
Di-calcium phosphate	2.07	2.06	2.06	2.05	2.05	1.9	1.9	1.9	1.9	1.9	1.7	1.7	1.7	1.7	1.7
Calcium carbonate <sup>1</sup>	1.06	1.03	1.01	1.01	1.01	1.05	1.05	1.05	1.05	1.05	0.92	0.92	0.92	0.90	0.90
NaCl	0.38	0.38	0.38	0.38	0.38	0.37	0.37	0.37	0.37	0.37	0.32	0.32	0.32	0.32	0.32
DL - Methionine	0.27	0.33	0.38	0.33	0.30	0.22	0.27	0.30	0.25	0.25	0.18	0.22	0.27	0.22	0.20
L - Lys HCl	0.53	0.40	0.35	0.31	0.36	0.44	0.34	0.28	0.32	0.22	0.36	0.36	0.27	0.21	0.22
L - Threonine	0.10	0.11	0.13	0.08	0.09	0.10	0.10	0.12	0.10	0.05	0.09	0.09	0.10	0.06	0.04
Vitamin & Mineral Permixon <sup>2</sup>	0.50	0.50	0.50	0.50	0.50	0.50	0.50	0.50	0.50	0.50	0.50	0.50	0.50	0.50	0.50
Choline chloride	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	0.22	0.22	0.22	0.22	0.22
Filler <sup>3</sup>	-----	-----	1.58	-----	-----	-----	-----	1.58	-----	-----	-----	-----	-----	-----	-----
Total	100	100	100	100	100	100	100	100	100	100	100	100	100	100	100
<b>Analyzed Nutrient content</b>															
ME (kcal/kg)	3035	2934	2853	3035	3120	3108	3014	2921	3108	3201	3185	3085	2990	3185	3275
Digestible Methionine%	0.59	0.59	0.59	0.59	0.59	0.53	0.53	0.53	0.53	0.53	0.48	0.48	0.48	0.48	0.48
Digestible Lysine%	1.18	1.18	1.18	1.18	1.18	1.05	1.05	1.05	1.05	1.05	0.95	0.95	0.95	0.95	0.95
Digestible Threonine%	0.77	0.77	0.77	0.77	0.77	0.69	0.69	0.69	0.69	0.69	0.65	0.65	0.65	0.65	0.65
Calcium%	0.90	0.90	0.90	0.90	0.90	0.84	0.84	0.84	0.84	0.84	0.76	0.76	0.76	0.76	0.76
Available phosphorus%	0.45	0.45	0.45	0.45	0.45	0.42	0.42	0.42	0.42	0.42	0.38	0.38	0.38	0.38	0.38
Na%	0.17	0.17	0.17	0.17	0.17	0.17	0.17	0.17	0.17	0.17	0.16	0.16	0.16	0.16	0.16

<sup>1</sup> per kg contains: Ca, 23% and P, 18.5%; <sup>2</sup> Supplied by Razak Co., Tehran, Iran, and provided per kilogram of diet: vitamin A, 360,000 IU; vitamin D3, 800,000 IU; vitamin E, 7,200 IU; vitamin K3, 800 mg; vitamin B1, 720 mg; vitamin B9, 400 mg; vitamin H2, 40 mg; vitamin B2, 2,640 mg; vitamin B3, 4,000 mg; vitamin B5, 12,000 mg; vitamin B6, 1,200 mg; vitamin B12, 6 mg; Choline chloride, 200,000 mg; Manganese, 40,000 mg; Iron, 20,000 mg; Zinc, 40,000 mg; copper, 4,000 mg; Iodine, 400 mg; <sup>3</sup> Inert filler used to complete diet formulations to 100%

**Table 2.** Primers sequences used in RT-PCR

Primers	Sequence
SGLT1 F	5-GATGTGCGGATACCTGAAGC-3
SGLT1 R	5-AGGGATGCCAACATGACTGA-3
FABP F	5-AGAAAGTTAGGAGGAGCCACG-3
FABP R	5-TCGGTCCACGGATTCAGC-3
β-actin F	5-CCACCGCAAATGCTTCTAAAC-3
β-actinR	5-AAGACTGCTGCTGACACCTTC-3

and water were provided for *ad libitum* intake. From day 21 to 41, all the chickens were exposed to 34±1°C and 60-70% relative humidity for 8 hours per day from 08:00 to 16:00, and then raised at 24±1°C. Feed and water were provided throughout the heat challenge period.

### Sample Collection

On day 28 of age, five chicks from each pen were weighed and euthanized by cervical dislocation. After

excising jejunum, as described by Uni et al.<sup>[24]</sup> segments were washed with cold phosphate buffer saline, sectioned and immediately frozen in liquid nitrogen, and stored at -80°C. 2 cm sections were selected for histo-morphology examination. The sections were flushed with phosphate buffer solution and then fixed in buffered formalin solution (10%). The entry of ceca of five chicks from each pen was sealed, removed and placed in ice and the contents used immediately for microbial assays.

### Total RNA Extraction and Reverse Transcription

The frozen jejunum was crushed in a sterile mortar, and the powder was applied for total RNA extraction using a suitable kit (Bioneer Co., Seoul, South Korea). The integrity of the RNA was verified by optical density (OD) absorption ratio 1.97>OD260 nm/OD280 nm>1.9. Ribonucleic acid integrity was determined by gel electrophoresis on a 1% agarose gel. Extracted RNA was stored at -80°C. Then, cDNA for each transporter gene was synthesized based on reverse transcription technique using kit (Bioneer Co., Seoul, South Korea) and stored at -20°C.

### Quantitative Real-Time PCR

The relative abundance of SGLT1 and FABP mRNA was determined by quantitative real-time PCR. Quantitative real-time PCR was conducted using a Real-time PCR systems (Applied Biosystems). Each reaction contained the followings: 2  $\mu$ L of cDNA, 10  $\mu$ L of 2X SYBR Green Master Mix (Applied Biosystems), 0.4  $\mu$ L each of the forward primer (4  $\mu$ M) and reverse primer (4  $\mu$ M), and 7.2  $\mu$ L of nuclease-free water. Primers were designed by using Primer software (primer-BLAST) at website ([www.ncbi.nlm.nih.gov](http://www.ncbi.nlm.nih.gov)). PCR reactions were performed with primers designed and synthesized for both transporters and the house-keeping gene  $\beta$ -actin (Table 2). Amplification for transporter in the jejunum (SGLT1 and FABP) was performed for 45 cycles, which consisted of an initial activations step (95°C, 5 min), denaturation cycle (95°C, 30s) and annealing at 54°C for 30 s, extension at 72°C for 1 min, and final extension at 72°C for 5 min. The PCR products were electrophoresed on a 1% agarose gel. Average gene expression relative to the endogenous control for each sample was calculated using the  $2^{-\Delta\Delta CT}$  method [25].

### Intestinal Microbial Populations

Immediately, samples of ileum and ceca were collected into glass containers under CO<sub>2</sub> sealed and put on ice until enumeration of microbial populations. Ten grams of mixed contents were blended under CO<sub>2</sub> in 90 mL of anaerobic dilution solution. Further serial dilutions were made in anaerobic dilution solution for anaerobic bacterial enumeration. The initial dilution in anaerobic dilution solution was also used as a source for serial dilutions in phosphate buffered saline for enumeration of aerobic bacterial populations. Triplicate plates were then inoculated with 0.1mL samples and incubated at 37°C aerobically or anaerobically as appropriate. Three dilutions were plated for each medium. Bacteria were enumerated on MRS agar (Merck, Germany) for *Lactobacillus* and MacConkey's (Merck, Germany) for *Escherichia coli*. Colony forming units (cfu) were defined as being distinct colonies measuring at least 1 mm in diameter [26].

### Intestinal Morphological Analysis

Formalin fixed tissue sections were processed by dehydration through a series of graded alcohol solutions (50, 70, 80, 90, 95 and 100%), cleared with xylene, and embedded in paraffin. For each segment, a 5- $\mu$ m cross-section was sectioned using a microtome and placed on a glass slide. The slides were stained using routine procedures for Mayer's hematoxylin and eosin. Villus height, villus length and crypt depth were measured using a light microscope (Olympus attached camera) using the method presented by Wang et al. [27]. Double-stained of samples with Periodic Acid-Schiff and hematoxylin was done to determine the goblet cell count according to the method of Wang et al. [27]. The goblet cells were counted in

the scale of 300  $\mu$ m of epithelium length.

### Statistical Analysis

Statistical analyses were carried out using ANOVA of SAS for Windows version 9.1 (SAS Institute Inc., Cary, NC) [28]. The Kolmogorov-Smirnov test was used to test the normality of the data before ANOVA was performed. Tukey test was used to compare the means. Statistical differences were declared at  $P < 0.05$ .

## RESULTS

### Effect on Performance

The effect of energy sources and levels on daily body weight gain (WG), daily feed intake (FI) and feed conversion ratio (FCR) is presented in Table 3. Chickens fed T1 and T2 diets as compared to C group had lower WG, higher FI and higher FCR ( $P < 0.05$ ). Chickens in T3 and T4 that received soybean oil and energy level equal to or greater than guideline of Cobb had higher WG compare to those received equal to or lower energy with main energy source of corn grain ( $P < 0.05$ ). Chicks in C and T3 group received the same amount of energy with different sources, had difference in FI and FCR, but had no significant differences in weight gain ( $P > 0.05$ ). Feed intake in T3 was lower than C, consequently, FCR of chicks fed T3 was better than C group. Although chicks in T3 and T4 groups received diets containing soybean oil, chicks in T4 received 3% higher energy than T3. Chicks in T3 also had lower FI than, same FCR and weight gain as group T4. The highest BW was for T4 and the lowest for T2 group. No difference on BW was observed between C and T3 groups consuming the same energy level with different energy source ( $P > 0.05$ ).

### Effect on Small Intestine Morphology

Dietary treatments significantly influenced jejunal villus height, crypt depth and villus height: crypt depth ratio (Table 4), but had no effect ( $P > 0.05$ ) on villus width and goblet cell count. The smallest villus height, deeper crypt and lowest their ratio was related to chicks in T2 group, and in contrast respectively the largest, shallowest and highest of these parameters was related to T4 group. Chickens received diets with lower energy level had poorer jejunal morphological parameters than those fed with diet in sufficient or over energy level. There were no differences for villus height and crypt depth between T3 and T4, but their ratio was higher ( $P < 0.05$ ) in T4 than T3. Chicks in C and T3 group received the same amount of energy with different sources, had difference in villus height and villus height: crypt depth ratio, but difference for crypt depth was not significant ( $P > 0.05$ ).

### Effect on Microbial Population

Effect of energy sources and levels on microbial population of broiler chicks is shown in Table 5. There

were no significant differences among C, T1 and T2 for *Lactobacillus* population ( $P>0.05$ ), whereas addition of soybean oil to diet in T3 and T4 groups increased ( $P<0.05$ ) *Lactobacillus* population. The highest population of *Escherichia coli* was seen in C group and the lowest in chickens received soybean oil. Chickens in T2 group that received the lowest energy level had less *Escherichia coli* population than T1 and C group ( $P<0.05$ ). Chicks in T3 group that received soybean oil in comparison with C

group that received the same amount of energy main from corn grain had higher ( $P<0.05$ ) *Lactobacillus* population and lower *Escherichia coli* population.

#### Effect on SGLT1 and I-FABP Gene Expression

Effect of dietary treatment on jejunal gene expression of SGLT1 and FABP in broiler chickens is shown in Table 6. There were no significant differences ( $P>0.05$ ) among treatments for gene expression of both nutrient transporters.

**Table 3.** Effect of energy sources and levels on performance parameters of broiler chickens at total period<sup>1</sup>

Treatments	Daily Weight Gain (g/bird)	Daily Feed Intake (g/bird)	Feed Conversion Ratio	BW (g)
C	51.42 <sup>b</sup>	93.07 <sup>c</sup>	1.81 <sup>c</sup>	2160 <sup>b</sup>
T1	49.83 <sup>bc</sup>	96.67 <sup>b</sup>	1.94 <sup>b</sup>	2093 <sup>c</sup>
T2	48.33 <sup>c</sup>	99.55 <sup>a</sup>	2.06 <sup>a</sup>	2030 <sup>d</sup>
T3	52.09 <sup>ab</sup>	89.59 <sup>d</sup>	1.72 <sup>d</sup>	2188 <sup>b</sup>
T4	53.80 <sup>a</sup>	93.61 <sup>c</sup>	1.74 <sup>d</sup>	2260 <sup>a</sup>
SEM	0.490	0.579	0.029	18.690
P-value	0.002	0.004	0.001	0.001

<sup>a,b,c</sup> Means within a column with different superscripts are significantly different ( $P<0.05$ );<sup>1</sup> Data are means of 20 pens of 30 broilers each

**Table 4.** Effect of energy sources and levels on jejunal morphological parameters of broiler chickens at d 28 of age<sup>1</sup>

Treatments	Villus Height <sup>2</sup> (mm)	Villus Width <sup>2</sup> (mm)	Crypt Depth <sup>2</sup> (mm)	Height:Crypt Depth <sup>1</sup>	Goblet Cell (count/mm) <sup>2</sup>
C	1.01 <sup>b</sup>	0.74	0.40 <sup>ab</sup>	2.52 <sup>c</sup>	140
T1	0.90 <sup>bc</sup>	0.75	0.43 <sup>ab</sup>	2.09 <sup>d</sup>	138
T2	0.70 <sup>c</sup>	0.75	0.50 <sup>a</sup>	1.4 <sup>e</sup>	137
T3	1.13 <sup>a</sup>	0.74	0.38 <sup>b</sup>	2.97 <sup>b</sup>	147
T4	1.19 <sup>a</sup>	0.72	0.36 <sup>b</sup>	3.30 <sup>a</sup>	146
SEM	0.23	0.651	0.09	0.08	0.606
P-value	0.001	0.07	0.001	0.04	0.08

<sup>a,b,c</sup> Means within a column with different superscripts are significantly different ( $P<0.05$ );<sup>1</sup> Data are means of 100 birds (5 birds from each pen);<sup>2</sup> Data were obtained from transmission electron microscopy

**Table 5.** Effect of energy sources and levels on viable counts of *Lactobacilli* and *Escherichia coli* ( $\log_{10}$  CFU/g of digesta) in cecal digesta of broiler chickens at d 28 of age<sup>1</sup>

Microflora	Experimental Treatments					SEM	P-value
	C	T1	T2	T3	T4		
<i>Lactobacillus</i>	3.68 <sup>c</sup>	3.72 <sup>c</sup>	3.50 <sup>c</sup>	4.67 <sup>b</sup>	5.83 <sup>a</sup>	0.539	0.008
<i>Escherichia coli</i>	5.00 <sup>a</sup>	4.82 <sup>a</sup>	2.20 <sup>b</sup>	1.30 <sup>c</sup>	1.10 <sup>c</sup>	0.500	0.008

<sup>a,b,c</sup> Means within a row with different superscripts are significantly different ( $P<0.05$ );<sup>1</sup> Data are means of 100 birds (5 birds from each pen)

**Table 6.** Effects of energy sources and levels on SGLT1 and FABP gene expression in jejunal of broiler chickens at 28 d of age

Relative Gene Expression <sup>1</sup>	Experimental Treatments					SEM	P-value
	C	T1	T2	T3	T4		
SGLT1	1.000	1.074	1.264	1.255	1.187	0.22	0.06
FABP	1.000	1.112	1.127	1.181	1.413	0.33	0.06

Means within a row without superscripts are not different ( $P>0.05$ );<sup>1</sup> Relative gene expression ( $2^{-\Delta\Delta C_T}$ )  $\pm$  SEM was calculated using the  $\Delta\Delta C_T$  method

## DISCUSSION

Chicken health and performance are greatly influenced by diet quality and nutrient availability [29]. These traits are more dependent on the intestine health to assimilate nutrients efficiently as the bird shifts from the lipid-rich yolk as the main source of energy to a carbohydrate-based diet [30]. To improve the health and performance, addition of lipids to diet in the early life of chicks and later was recommended by many researchers [31-33]. In stress condition, plasma corticosterone level tends to increase and this hormone has negative effect on intestinal morphology and consequently performance [34]. A recent study revealed that birds exposed to chronic heat stress had poor growth performance and increased corticosterone concentrations [6]. Deteriorated performance resulted from heat stress in broilers can be attributed to a greater expenditure of energy for physiological adaptation to the stressful situation instead of growth enhancement [35]. Alternatively, it is believed that less weight gain in the heat stress groups is due to a smaller appetite and lower feed intake, as it maybe seemingly a defense mechanism to help reduce heat production.

The main objective of this study was to evaluate the effect of energy sources and levels on microbial population, SGLT1 and FABP gene expression and small intestinal parameters in broiler chickens under heat stress. In the literature, study about this subject under heat stress was not found; hence comparison of our findings with others was done on non-stressed condition.

The results of this study indicate that the source and level of metabolizable energy in broilers under heat stress had a statistically significant influence on intestinal morphology and microbiology and consequently performance ( $P < 0.05$ ), but it had no effect on gene expression of nutrient transporters of glucose and fatty acid binding protein. Maintenance of normal microarchitecture of the small intestine is very important for proper growth and development. Quite a few studies have reported that stress hampered the development of intestinal morphology and function [14]. Sohail et al. [6] observed that heat stress decreased villus height and width, crypt depth, and villus surface area. Stressors such as fasting and nutrient deficiency or corticosterone injections have noxious effects on the intestinal microarchitecture, resulting in reduction in the absorptive surface area [36,37]. Feeding ration with lower energy level to chickens resulted in short villus height and poor performance. The villi play a crucial role in the digestion and absorption processes of the small intestine, as is the first to make contact with nutrients in the lumen [38]. A shortening of the villus height may lead to poor nutrient absorption along with lower performance [26].

In contrast, substitution of a part of corn energy with soybean oil in the present study improved the intestinal

parameters. The positive effect of lipids on intestinal health status and growth performance of broilers is well documented [39]. The extra caloric effect of added soybean oil resulted in an improved body weight gain [40]. Inclusion of soybean oil and delivery of higher energy than Cobb standard (T4) resulted in the highest villus height, the shallowest crypt depth, lowest *E. coli* and highest lactic acid bacteria population and finally higher body weight and the highest feed efficiency. It was reported that dietary soybean oil supplementation significantly increased villus height of the jejunum [41].

In this experiment, addition of soybean oil in diet T3 and T4 enhanced the growth of *Lactobacillus*, but inhibited that of *E. coli* in the small intestine. Freitas et al. [42] reported that addition of soybean oil could maximize the growth of the bacterial population. Intestinal microflora plays an important role in digestive health. Microbial population is dependent on food rations as the ultimate source of the organic substrate metabolism [43]. It is known that dietary fats cause changes in the intestinal microflora composition with a direct effect on digestion and absorption of nutrients by the birds. Innis et al. [44] reported that canola oil supplementation increased lactic acid bacteria population in the gut. The effect of fat on the microbial flora of the gastrointestinal tract may be due to its effect on digestion, viscosity, pH level and the transport of nutrients in the gastrointestinal tract [45].

No differences were found among treatments for gene expression of SGLT1 and FABP in this study. It was demonstrated that regulation of nutrient transporters by dietary substrate appears to occur by increasing mRNA stability or by increasing gene transcription rate [46]. It seems that metabolizable substrates are not necessary for its regulation. Sun et al. [47] reported that the expression levels of FABP were significantly decreased by heat exposure, but heat stress had no significant effect on gene expression of SGLT1. In Shepherd et al. [48] mentioned study has suggested that environmental stress decreases GLUT-2 expression at the brush border membrane level but does not alter SGLT-1 expression. Moreover, corticosterone-induced stress does not alter the expression of SGLT-1 in the jejunum of broiler chickens [49].

Moreover, an increased body temperature during heat stress conditions can lead to increased maintenance energy requirements [50] to keep body temperature around a normal level. Different levels of energy significantly affected body weight, weight gain, feed intake and feed conversion ratio [51]. The final live weight was significantly highest in broiler chickens fed dietary treatment with normal energy and was lowest in those fed dietary treatment with low energy [52]. Increased live weight was mostly due to higher metabolizable energy consumption in the same unit of diets by chickens, similarly supplementation of oil caused a positive trend in cumulative live weight gain (g/bird) of broilers at different ages [53]. Fats and vegetable oils

has been frequently included in broiler diets to increase the energy density of the diet, to improve efficiency and to increase nutrient digestibility in broilers<sup>[54]</sup>. Lipid supplementation increases the energetic efficiency in two ways. It increases density of energy and it has a lower heat increment or greater net energy. Increased energy and nutrient density of the diet and replacing carbohydrate calories reduced feed intake almost proportionally but increased live weight gains in hot weather. Nitsan et al.<sup>[40]</sup> stated that feed conversion ratio were significantly improved with addition of 3% soybean oil in the diet.

The present study has demonstrated that the levels and sources of dietary energy have significant effects on intestinal populations of *Lactobacillus* and *E. coli* in broiler chickens. It can be concluded that providing high levels of dietary energy with oil in broiler nutrition, increases beneficial intestinal flora and reduction of noxious microbes and simultaneously cause an increase intestinal villus height and decrease crypt depth and intestinal absorption that finally can ensure the intestinal health of broilers. The results showed that the higher energy level than nutritional needs based on Cobb broiler chickens requirements as specified in the manual was affective on microbial population and morphological parameters that can cause increasing weight gain and final body weight under heat stress.

## ACKNOWLEDGMENTS

The authors are grateful to the Islamic Azad University for research funding support and to Animal Science Institute for supplying the basal diet and allowing use of poultry unit. We also thank all staffs in the poultry unit, for the assistance in the care and feeding of the chicks used in this research.

## REFERENCES

- Windhorst HW:** Changes in poultry production and trade worldwide. *World's Poult Sci J*, 62, 585-602, 2006. DOI: 10.1079/WPS2006114
- Lin H, Jaio HC, Buysse J, Decuyper E:** Strategies for preventing heat stress in poultry. *World's Poult Sci J*, 62, 71-85, 2006. DOI: 10.1079/WPS200585
- Schlesinger MJ, Ryan C, Chi MM, Carter JG, Pusateri ME, Lowry OH:** Metabolite changes associated with heat shocked avian fibroblast mitochondria. *Cell Stress Chaperones*, 2, 25-30, 1997.
- Lan PT, Sakamoto M, Benno Y:** Effects of two probiotic *Lactobacillus* strains on jejunal and cecal microbiota of broiler chicken under acute heat stress condition as revealed by molecular analysis of 16S rRNA genes. *Microbiol Immunol*, 48, 917-929, 2004. DOI: 10.1111/j.1348-0421.2004.tb03620.x
- Tao X, Zhang ZY, Dong H, Zhang H, Xin H:** Responses of thyroid hormones of market-size broilers to thermoneutral constant and warm cyclic temperatures. *Poult Sci*, 85, 1520-1528, 2006. DOI: 10.1093/ps/85.9.1520
- Sohail MU, Hume ME, Byrd JA, Nisbet DJ, Ijaz A, Sohail A, Shabbir MZ, Rehman H:** Effect of supplementation of prebiotic mannan-oligosaccharides and probiotic mixture on growth performance of broilers subjected to chronic heat stress. *Poult Sci*, 91, 2235-2240, 2012. DOI: 10.3382/ps.2012-02182
- Yu J, Yin P, Liu F, Cheng G, Guo K, Lu A, Zhu X, Luan W, Xu J:** Effect of heat stress on the porcine small intestine: a morphological and gene expression study. *Comp Biochem Physiol*, 156, 119-28, 2010. DOI: 10.1016/j.cbpa.2010.01.008
- Quinteiro-Filho WM, Ribeiro A, Ferraz-de-Paula V, Pinheiro ML, Sakai M, Sa LRM, Ferreira AJ, Palermo-Neto J:** Heat stress impairs performance parameters, induces intestinal injury, and decreases macrophage activity in broiler chickens. *Poult Sci*, 89, 1905-1914, 2010. DOI: 10.3382/ps.2010-00812
- Menten JFM, Barbosa Filho JAD, Silva MAN, Silva IJO, Racanicci AMC, Coelho AAD, Savino VJM:** Physiological responses of broiler chickens to pre-slaughter heat stress. *World's Poult Sci J*, 62, 254-257, 2006.
- Aksit M, Yalcin S, Ozkan S, Metin K, Ozdemir D:** Effects of temperature during rearing and crating on stress parameters and meat quality of broilers. *Poult Sci*, 85, 1867-1874, 2006. DOI: 10.1093/ps/85.11.1867
- Kohl KD:** Diversity and function of the avian gut microbiota. *J Comp Physiol B*, 182,591-602, 2012. DOI: 10.1007/s00360-012-0645-z
- Apajalahti JHA, Kettunen A, Bedford MR, Holben WE:** Percent G+C profiling accurately reveals diet-related differences in the gastrointestinal microbial community of broiler chickens. *Appl Environ Microbiol*, 67, 5656-5667, 2001. DOI: 10.1128/AEM.67.12.5656-5667.2001
- Soltan M:** Influence of dietary glutamine supplementation on growth performance, small intestinal morphology, immune response and some blood parameters of broiler chickens. *Int J Poult Sci*, 8, 60-68, 2009. DOI: 10.3923/ijps.2009.60.68
- Mitchell MA, Carlisle AJ:** The effect of chronic exposure to elevated environmental temperature on intestinal morphology and nutrient absorption in the domestic fowl (*Gallus domesticus*). *Comp Biochem Physiol*, 101, 137-142, 1992. DOI: 10.1016/0300-9629(92)90641-3
- Thorens B:** Glucose transporters in the regulation of intestinal, renal and liver glucose fluxes. *Am J Physiol*, 270, 541-553, 1996.
- Wright EM, Turk E:** The sodium/glucose co-transporter family SLC5. *Eur J Physiol*, 447, 510-518, 2004. DOI: 10.1007/s00424-003-1063-6
- Furuhashi M, Hotamisligil GS:** Fatty acid-binding proteins: role in metabolic diseases and potential as drug targets. *Nat Rev Drug Discov*, 7, 489-503, 2008. DOI: 10.1038/nrd2589
- Gal-Garber OS, Mabjeesh J, Sklan D, Uni Z:** Partial sequence and expression of the gene for and activity of the sodium glucose transporter in the small intestine of fed, starved and refeed chickens. *J Nutr*, 130, 2174-2179, 2000.
- Naruhashi K, Sai Y, Tamai I, Suzuki N, Tsuji A:** PepT1 mRNA expression is induced by starvation and its level correlates with absorptive transport of cefadroxil longitudinally in the rat intestine. *Pharm Res*, 19, 1417-1423, 2002. DOI: 10.1023/A:1020436028194
- Wang X, Peebles ED, Zhai W:** Effects of protein source and nutrient density in the diets of male broilers from 8 to 21 days of age on their subsequent growth, blood constituents, and carcass compositions. *Poult Sci*, 93, 1463-1474, 2014. DOI: 10.3382/ps.2013-03838
- Zai W, Peebles ED, Zumwalt CD, Mejia L, Corozo A:** Effects of dietary amino acid density regimens on growth performance and meat yield of Cobb×Cobb 700 broilers. *J Appl Poult Res*, 22, 447-460, 2013. DOI: 10.3382/japr.2012-00658
- Daghir NJ:** Broiler feeding and management in hot climates. In, Daghir NJ (Ed): Poultry Production in Hot Climate. 185-218, CAB Intl, Wallingford, UK, 1995.
- Zulkifli I, Liew PK, Israf DA, Omar AR, Hair-Bejo M:** Effects of early age feed restriction and thermal conditioning on heterophil/lymphocyte ratio, heat shock protein 70 and body temperature of male broiler chickens subjected to acute heat stress. *J Thermal Biol*, 28, 217-222, 2003. DOI: 10.1016/s0306-4565(02)00058-x
- Uni Z, Noy Y, Sklan D:** Post-hatch development of smallintestinal function in the poultry. *Poult Sci*, 78, 215-222, 1999. DOI: 10.1093/ps/78.2.215
- Livak KJ, Schmittgen TD:** Analysis of relative gene expression data using real-time quantitative PCR and the 2<sup>-ΔΔC<sub>T</sub></sup> method. *Methods*, 25, 402-408, 2001. DOI: 10.1006/meth.2001.1262

- 26. Xu ZR, Hu CH, Xia MS, Xia X, Zhan A, Wang MQ:** Effects of dietary fructo-oligosaccharide on digestive enzyme activities, intestinal microflora and morphology of male broilers. *Poult Sci*, 82, 1030-1036, 2003. DOI: 10.1093/ps/82.6.1030
- 27. Wang X, Peebles ED, Morgan TW, Harkess RL, Zhai W:** Protein source and nutrient density in the diets of male broilers from 8 to 21 d of age: Effects on small intestine morphology. *Poult Sci*, 94, 61-67, 2015. DOI: 10.3382/ps/peu019
- 28. SAS Institute Inc:** SAS/STAT 9.13 User's Guide. *SAS Inst Inc*, Cary, NC, 2004.
- 29. Geyra A, Uni Z, Sklan D:** Enterocyte dynamics and mucosal development in the post-hatch chick. *Poult Sci*, 80, 776-782, 2001. DOI: 10.1093/ps/80.6.776
- 30. Sklan D, Geyra A, Tako E, Gal-Gerbert O, Uni Z:** Ontogeny of brush border carbohydrate digestion and uptake in the chick. *Br J Nutr*, 89, 747-753, 2003. DOI: 10.1079/BJN2003853
- 31. Yaqoob P:** Fatty acids and the immune system: from basic science to clinical applications. *Proceed Nutr Soc*, 63, 89-104, 2004. DOI: 10.1079/PNS2003328
- 32. Sadeghi A, Mirmohseni A, Shawrang P, Aminafshar M:** The effect of soy oil addition to the diet of broiler chicks on the immune response. *Turk J Vet Anim Sci*, 37, 264-270, 2013. DOI: 10.3906/vet-1109-24
- 33. Cherian G:** Nutrition and metabolism in poultry: Role of lipids in early diet. *J Anim Sci Biotechnol*, 6, 28-35, 2015. DOI: 10.1186/s40104-015-0029-9
- 34. Geraert PA, Padilha JCF, Guillaumin S:** Metabolic and endocrine changes induced by chronic heat exposure in broiler chickens: Biological and endocrinological variables. *Br J Nutr*, 75, 205-216, 1996. DOI: 10.1017/BJN19960125
- 35. Lei KY, Slinger SJ:** Energy utilization in the chick in relation to certain environmental stresses. *Can J Anim Sci*, 50, 285-292, 1970. DOI: 10.4141/cjas70-043
- 36. Yamauchi K, Yamamoto K, Ishiki Y:** Morphological alterations of the intestinal villi and absorptive epithelial cells in each intestinal part in fasted chickens. *Jpn J Poult Sci*, 32, 241-251, 1995. DOI: 10.2141/jpsa.32.241
- 37. Hu XF, Guo YM:** Corticosterone administration alters small intestinal morphology and function of broiler chickens. *Asian-Australas J Anim Sci*, 21, 1773-1778, 2008. DOI: 10.5713/ajas.2008.80167
- 38. Gartner LP, Hiatt JL:** Color Textbook of Histology. 2<sup>nd</sup> edn., WB Saunders, Baltimore, MD, USA, 2001.
- 39. Griffiths L, Leeson S, Summers JD:** Influence of energy system and level of various fat sources on performance and carcass composition of broiler. *Poult Sci*, 56, 1018-1026, 1977. DOI: 10.3382/ps.0561018
- 40. Nitsan Z, Dvora A, Zoref Z, Mokady S:** Effect of added soybean oil and dietary energy on metabolisable and net energy of broiler diets. *Br Poult Sci*, 38, 101-106, 1997. DOI: 10.1080/00071669708417948
- 41. Markovic R, Sefer D, Krstic M, Petrujkic B:** Effect of different growth promoters on broiler performance and gut morphology. *Archiv Med Vet*, 60 (2): 442-448, 2009.
- 42. Freitas KC, Gabriel JE, Leite LC, de Armas RD, Lanna DPD, Madeira HMF:** Molecular characterization of ruminal bacterial diversity. *Acta Scientiific Anim Sci*, 30, 187-192, 2008. DOI: 10.4025/actascianimsci.v30i2.4699
- 43. Choct M, Hughes RJ, Wang J, Bedford M, Morgan AJ, Annison G:** Increased small intestinal fermentation is partly responsible for the anti-nutritive activity of non-starch polysaccharides in chickens. *Br Poult Sci*, 37, 609-621, 1996. DOI: 10.1080/00071669608417891
- 44. Innis SM, Pinsk V, Jacobson K:** Dietary lipids and intestinal inflammatory diseases. *J Peds*, 58, 89-95, 2006. DOI: 10.1016/j.jpeds.2006.06.058
- 45. Laflamme DR, Xu H, Long GM:** Effect of diets differing in fat content on chronic diarrhea in cats. *J Vet Intern Med*, 25, 230-235, 2011. DOI: 10.1111/j.1939-1676.2010.06655.x
- 46. Adibi S:** Regulation of expression of the intestinal oligopeptide transporter (PepT-1) in health and disease. *Am J Physiol-Gastrointestinal Liver Physiol*, 285, 779-788, 2003. DOI: 10.1152/ajpgi.00056.2003
- 47. Sun X, Zhang H, Sheikahmadi A, Wang Y, Jiao H, Lin H, Song Z:** Effects of heat stress on the gene expression of nutrient transporters in the jejunum of broiler chickens (*Gallus gallus domesticus*). *Intl J Biometeorol*, 59, 127-135, 2015. DOI: 10.1007/S00484-014-0829-1
- 48. Shepherd EJ, Helliwell PA, Mace OJ, Morgan EL, Patel N, Kellett GL:** Stress and glucocorticoid inhibit apical Glut2-trafficking and intestinal glucose absorption in rat small intestine. *J Physiol*, 560, 281-290, 2004. DOI: 10.1113/jphysiol.2004.072447
- 49. Hu XF, Guo YM, Huang BY, Bun S, Zhang LB, Li JH, Liu D, Long FY, Yang X, Jiao P:** The effect of glucagon-like peptide 2 injection on performance, small intestinal morphology, and nutrient transporter expression of stressed broiler chickens. *Poult Sci*, 89, 1967-1974, 2010. DOI: 10.3382/ps.2009-00547
- 50. Hurwitz S, Weiselberg M, Eisner M, Bartov I, Riesenfeld G, Sharvit M, Niv A, Bornstein S:** The energy requirements and performance of growing chickens and turkeys as affected by environmental temperature. *Poult Sci*, 59, 2209-2299, 1980. DOI: 10.3382/ps.0592290
- 51. Min YN, Shi JS, Wei FX, Wang HY, Hou XF, Niu ZY, Liu FZ:** Effects of dietary energy and protein on growth performance and carcass quality of broilers during finishing phase. *J Anim Vet Advances*, 11, 3652-3657, 2012. DOI: 10.3923/javaa.2012.3652.3657
- 52. Dairo F, Adeshinwa ASAOK, Oluwasola TA, Oluyemi JA:** High and low dietary energy and protein levels for broiler chickens. *Afr J Agri Res*, 5 (15): 2030-2038, 2010.
- 53. Das GB, Hossain ME, Akbar MA:** Effects of different oils on productive performance of broiler. *Iranian J Appl Anim Sci*, 4 (1): 111-116, 2014.
- 54. Monfaredi A, Rezaei M, Sayyahzadeh H:** Effect of supplemental fat in low energy diets on some blood parameters and carcass characteristics of broiler chicks. *S Afr J Anim Sci*, 41, 24-32, 2011.