

Effects of *GH-AluI* and *MYF5-TaqI* Polymorphisms on Weaning Weight and Body Measurements in Holstein Young Bulls ^[1]

Mehmet Ulaş ÇINAR ¹ Bilal AKYÜZ ²  Jale METİN KIYICI ¹
Korhan ARSLAN ² Mahmut KALİBER ¹ Esmâ Gamze AKSEL ²

¹ Department of Animal Science, Faculty of Agriculture, Erciyes University, TR-38000 Kayseri - TURKEY

² Department of Genetics, Faculty of Veterinary Medicine, Erciyes University, TR-38000 Kayseri - TURKEY

Article Code: KVFD-2018-20193 Received: 21.01.2018 Accepted: 23.08.2018 Published Online: 05.09.2018

How to Cite This Article

Çınar MU, Akyüz B, Metin Kıyıcı J, Arslan K, Kaliber M, Aksel EG: Effects of *GH-AluI* and *MYF5-TaqI* polymorphisms on weaning weight and body measurements in Holstein young bulls. *Kafkas Univ Vet Fak Derg*, 24 (6): 873-880, 2018. DOI: 10.9775/kvfd.2018.20193

Abstract

Live weight and body measurements are economically important quantitative traits that affect carcass yield and calf survival in cattle. Four genes, growth hormone gene (*GH*), myogenic factor 5 (*MYF5*), fatty acid binding protein 4 (*FABP4*) and signal transducers and activators of transcription 5A (*STAT5A*) were chosen as candidate genes for live weight and body measurements due to their important role in growth. The aims of this study were to genotype *GH-AluI*, *MYF5-TaqI*, *FABP4-HinII* and *STAT5A-AvaI* polymorphisms and to investigate their associations with live weights and body measurements in Holstein young bulls. Genotyping of the single nucleotide polymorphism (SNP) markers in these candidate genes was carried out using the restriction fragment length polymorphism (RFLP) analysis. Frequencies of L allele for *GH*, A allele for *MYF5*, G allele for *FABP4* and C allele for *STAT5A* were, 0.96, 0.61, 0.79 and 0.74, respectively in the examined animals. The regression analysis indicated that the *GH-AluI* polymorphism showed an association with weaning weight (WW), 180th day weight and hearth girth from birth to 180 days of age. The *MYF5-TaqI* polymorphism was found to influence body length at birth and birth weight (BW). However, no significant association was detected between the *FABP4-HinII* genotypes and measured traits. The *GH-AluI* and *MYF5-TaqI* polymorphisms may be useful for selection on live weight and body measurement traits in Holstein young bulls.

Keywords: *GH*, Live weight, *MYF5*, RFLP, Weaning weight

GH-AluI ve *MYF5-TaqI* Polimorfizmlerinin Erkek Holstein Buzağlarında Sütten Kesim Ağırlığı ve Vücut Ölçüleri Üzerine Etkileri

Öz

Canlı ağırlık ve vücut ölçüleri gibi önemli kantitatif özellikler, karkas verimi ve buzağı yaşama gücünü etkilemektedirler. Büyümedeki rolleri nedeniyle büyüme hormonu geni (*GH*), miyojenik faktör 5 (*MYF5*), yağ asidi bağlayıcı protein 4 (*FABP4*) ve sinyal dönüştürücü ve transkripsiyonu aktive edici faktör 5A (*STAT5A*) canlı ağırlık ve vücut ölçüleri için aday gen olarak önerilmiştir. Bu çalışmada erkek Holstein buzağlarında *GH-AluI*, *MYF5-TaqI*, *FABP4-HinII* ve *STAT5A-AvaI* polimorfizmleri belirlenerek, elde edilen genotip verileriyle canlı ağırlık artışı ve vücut ölçüleri arasındaki ilişkilerin araştırılması amaçlanmıştır. İncelenen örneklerin bu aday genlerdeki SNP markırları yönünden genotipleri restriksiyon parçacık uzunluk polimorfizmi (RFLP) analiziyle belirlenmiştir. İncelenen örneklerde *GH* için L allel, *MYF5* için A allel, *FABP4* için G allel ve *STAT5A* için C allel frekansı diğer allelden yüksek bulunmuştur (sırasıyla 0.96, 0.61, 0.79 ve 0.74). Elde edilen genotip verileri kullanılarak yapılan regresyon analizi sonunda *GH-AluI* polimorfizmi ile sütten kesimdeki (WW) ve 6. ay canlı ağırlıkları ile tüm ölçüm dönemlerindeki göğüs çevresi uzunluğu ile ilişkili olduğu görülmüştür. *MYF5-TaqI* polimorfizmi ise doğumdaki vücut uzunluğu ve doğum ağırlığı (BW) ile ilişkili bulunmuştur. İncelenen örneklerde *FABP4-HinII* polimorfizmi ölçülen özelliklerin hiç biri ile ilişkili bulunmamıştır. Çalışma sonunda *GH-AluI* ve *MYF5-TaqI* polimorfizmlerinin canlı ağırlık ve vücut ölçüleri yönünden yapılacak seleksiyon çalışmalarında kullanılabilmesi düşünülmüştür.

Anahtar sözcükler: *GH*, Canlı ağırlık, *MYF5*, RFLP, Sütten kesim ağırlığı

INTRODUCTION

Beef is an excellent source of protein for human nutrition but, the production costs is higher compared to poultry

and pork ^[1]. Therefore, increasing the meat yield is one of the most important issue in cattle breeding specially in countries in which their consumer preferences depending on mainly ruminant meat. Genomic selection



İletişim (Correspondence)



+90 352 2076666/29721



bakyuz@erciyes.edu.tr

is an encouraging development in livestock, proposing improved production by deciphering molecular genetic markers to design novel breeding programs and to develop new markers-based models for selecting favorable genotypes [2].

One of the best known genetic marker for beef yield is the growth hormone gene (*GH*) [3,4] which is encoding for growth hormone. *GH* is located on cattle chromosome 19 (BTA19) [5], and consists of five exons and four introns [6]. It is playing role in numerous physiological processes such as growth traits, mammary gland development and lactation [7]. Although, studies have mostly focused on association of *GH* genotype with milk yield traits, association of *GH* genotypes with live weight and body measurement traits have been less investigated [8]. Several polymorphisms were detected in the bovine *GH* gene, however, the best known of these polymorphisms is the leucine (L) to valine (V) substitution at position 127 in exon 5 of *GH* gene which creates a cut point for *AluI* restriction enzyme [4].

Skeletal muscles occur result of a series of physiological processes which is named as myogenesis, and it includes cell specification, proliferation and differentiation, in which multi-potential mesodermal cells are differentiated into myoblasts [9]. Myogenesis begins at embryonic stage and continues to postnatal maturation [10] and controlled mainly by myogenic determination (*MyoD*) gene family [11]. Myogenic factor 5 (*MYF5*) gene is one of the four members of *MyoD* family [12] and located on BTA5 in bovine [13]. It is evaluated as a candidate gene for growth traits in beef cattle breeding because of its tasks in muscle development and growth traits [13,14].

The fatty acid binding protein 4 (*FABP4*) has got important tasks in lipid hydrolysis and intracellular fatty acid uptake in different tissues [15,16]. This protein is encoded by *FABP4* gene and its mRNA expresses firstly in adipocytes [17] from a locus located on BTA14 in cattle [18]. In literature, few reports were shown the relationship between *FABP4* and carcass weight in native Korean cattle breed [19,20]. However, no study was devoted to association *FABP4* variants with live weight and body measurements in Holstein cattle.

Signal transducers and activators of transcription protein (STAT) is a family of transcription factors which is mediated the effects of some peptide hormones and cytokines. STAT family has seven members [21], of them *STAT5A* is an important mediator of growth hormone which is located on BTA19 in bovine [22]. Therefore, it was thought that *STAT5A* gene may be used as a marker for important yield traits such as growth and live weight gain in livestock [23].

Although functional relation of these genes in growth metabolism has been reviewed in literature, association of polymorphisms in these genes with live weight traits and body measurements have not been strongly taken into consideration in Holstein cattle breed. Therefore, aims of

this study were to investigate SNPs in *GH*, *MYF5*, *FABP4* and *STAT5A* genes, and investigate their association with live weight traits and some body measurements in Holstein young bulls reared in Turkey.

MATERIAL and METHODS

The project was approved by the relevant Animal Ethics committee of Erciyes University (#13/72 10.04.2013). A total of 59 male Holstein calves, born between March 2013 and November 2014 were used in this study. Animals were not applied to a special feeding program. Calves were stay with their mother after birth and consume *ad-libitum* colostrum for three days. At the end of three days, calves were fed with milk 10% of their birth weight until weaning. At seven days old, calf starter feed which contains calve growing feed and forages were provided *ad-libitum* until weaning. After weaning, calves were fed with milk replacer for 40 days until milk replacer weaning day. Young bulls were fed with forage and beef cattle feed (concentrated protein) mixture until 180th day. As live weight traits; birth weight (BW), weaning weight (30. day) (WW), milk replacer weaning weight (70. day) (RW) and weight at 180th day after birth (SW) were measured. The animals were weighed to the nearest kilogram using an electronic weighing scale (EziWeigh 5i, Tru-Test, New Zealand) mounted on a concrete platform. For body measurements; body length, wither height and hearth girth were measured. Body measurements were taken by two observers using an ordinary measuring tape and recorded in centimeters. Body length was measured as the distance from *Atriculus huneri* to *Tuber ichii*; wither height was measured as the distance from the ground to the highest point of wither. Hearth girth was measured as behind the front shoulder at the fourth ribs, posterior to the front leg [24].

All blood samples were collected post-natal period, and genomic DNA was extracted by the phenol:chloroform:i soamylalcohol method. The *GH*, *MYF5*, *FABP4* and *STAT5A* gene polymorphisms were genotyped by using PCR-RFLP. The PCR reactions mixtures of all genes were prepared as total volume of 25 μ L, including 1.5 mM $MgCl_2$, 200 μ M dNTP, 5 pmol of forward and reverse primer of each gene, 1 \times PCR buffer, 1U Taq polymerase and approximately 100 ng DNA. The PCR protocols for investigated genes were as follow: for *GH* gene, 4 min at 94°C for, then 40 cycles of 94°C for 40 s, 60°C for 40 s, 72°C for 40 s and the final extension at 72°C 10 min; for *MYF5* gene, initial denaturation at 94°C for 4 min, then 38 cycles of 94°C for 1 min, 64°C for 30 s, 72°C for 1 min and final extension at 72°C for 4 min; for *FABP4* gene, initial denaturation at 95°C for 4 min, then 32 cycles of 94°C for 1 min, 62°C for 1 min, 72°C for 1 min and final extension at 72°C for 10 min; for *STAT5A* gene, pre-denaturation at 95°C for 4 min, then 34 cycles of 94°C for 1 min, 64°C for 1 min, 72°C for 1 min and final extension at 72°C for 10 min. The obtained PCR products were digested by 5 U of restriction endonuclease enzymes according to

their own protocols (Fermentas, Thermo Fisher Scientific Inc., Waltham, MA, USA). After digestion procedures, the genotypes were detected on 3% (for *GH* and *STAT5A* genes) and 2% (for *FABP4* and *MYF5* genes) agarose (Prona Agarose; Basica Le, Burgos, Spain) gel electrophoresis. The nucleotide sequences of PCR primers and restriction enzymes used for genotyped amplification and RFLP process are presented in *Table 1*.

Allele and genotype frequencies of genes, investigated in the present study were calculated by using OEGE - Online Encyclopedia for Genetic Epidemiology studies online tool [29]. Statistical analyses were performed by using IBM SPSS Statistics 22.0 software. For *GH*, *MYF5*, *FABP4* and *STAT5A* genes genotype and phenotype associations were investigated by using independent general linear model (GLM) procedure and Duncan test for significance levels; $P < 0.1$ and $P < 0.05$. Sire ($n=11$) were assumed random factor in the GLM model. The statistical model used as follows:

$$Y_{ij} = \mu + S_j + G_i + e_{ij}$$

Where Y_{ij} is the observation of the weaning weight and body measurements traits; μ is the overall mean for each

trait, S_j is the random effect of j^{th} sire, G_i is the fixed effect of i^{th} genotype for the relevant polymorphism and e_{ij} is the random residual error.

RESULTS

The 223 bp products were obtained after PCR process and PCR products were digested by *AluI* restriction enzyme for *GH* gene. At the end of digestion, it was expected one band (223 bp) for VV genotype, three bands (223, 171 and 52 bp) for LV genotype, two bands (171 and 52 bp) for LL genotype. The band of 52 bp could not be observed on agarose gel electrophoresis. However, two bands (223 and 171 bp) were enough for genotyping (*Fig. 1a*). The LL genotype was found to be the highest frequency, and the VV genotype was found to be the lowest frequency in our investigated Holstein population. The L allele frequency was higher than V allele (*Table 2*).

The 490 bp products were obtained after PCR process and they were digested by *TaqI* restriction enzyme for *MYF5* gene. At the end of digestion, it was observed one band (490 bp) for AA genotype, three bands (490, 367 and 123

Table 1. Primer sequence, accession number, PCR product size

Gene	Accession Number	Sequence	Product Size	Restriction Enzyme	Reference
GH	EF592534.1	GCTGCTCCTGAGGGCCCTTCG GCGGCGGCACTTCATGACCCT	223 bp	<i>AluI</i>	[25]
MYF5	M95684.1	AGAGCAGCAGTTTTGACAGC GCAATCCAAGCTGGATAAGG	512 bp	<i>TaqI</i>	[26]
FABP4	NC007312	ATTATCCCCACAGAGCATCG ACAAGACTTGGCCTCAAGGA	399 bp	<i>HinII</i>	[27]
STAT5A	AJ237937	CTGCAGGGCTGTCTGAGAG TGGTACCAGGACTGTAGCACAT	215 bp	<i>AvaI</i>	[28]




Fig 1. Agarose electrophoresis pattern of GH (a), MYF5 (b), FABP4 (c) and STAT5A (d). M: molecular marker (100 bp)

Table 2. Genotype and allele frequencies of the GH, MYF5, FABP4 and STAT5A genes in beef Holstein cattle

Gene	Genotype						Allele Frequency		Chi-squared P=0.734 (df=1)
	LL		LV		VV		L	V	
GH	Obs (Exp)	F	Obs (Exp)	F	Obs (Exp)	F	0.96	0.04	
	54 (54.11)	0.915	5 (4.79)	0.085	0 (0.11)	-			
MYF5	AA		AG		GG		A	G	$\chi^2=1.16$ P=0.2819 (df=1)
	Obs (Exp)	F	Obs (Exp)	F	Obs (Exp)	F	0.61	0.39	
	20 (21.97)	0.339	32 (28.07)	0.542	7 (8.97)	0.119			
FABP4	AA		AG		GG		A	G	$\chi^2=4.26$ P=0.0389 (df=1)
	Obs (Exp)	F	Obs (Exp)	F	Obs (Exp)	F	0.21	0.79	
	0 (2.65)	-	25 (19.7)	0.424	34 (36.25)	0.576			
STAT5A	CC		CT		TT		C	T	$\chi^2=7.49^{**}$ P=0.0062 (df=1)
	Obs (Exp)	F	Obs (Exp)	F	Obs (Exp)	F	0.74	0.26	
	28 (32.07)	0.475	31 (22.86)	0.525	0 (4.07)	-			

Obs: Observed genotype; Exp: Expected genotype; F: Frequency; df: degree of freedom

Table 3. Least squares means and standard errors for BW, WW, RW and SW in male calves according to GH, MYF5, FABP4 and STAT5A genotypes

Gene	Genotype	N	Traits			
			BW (kg) Mean (\pm SE)	WW (kg) Mean (\pm SE)	RW (kg) Mean (\pm SE)	SW (kg) Mean (\pm SE)
GH	LL	54	42.38 \pm 1.13	53.37 \pm 1.25 ^a	73.21 \pm 1.71	188.57 \pm 4.78
	LV	5	39.80 \pm 2.70	46.89 \pm 2.98 ^b	72.43 \pm 4.09	166.30 \pm 11.41
	P		0.360	0.041	0.856	0.065
MYF5	AA	20	45.13 \pm 6.57	55.61 \pm 6.47	72.88 \pm 8.21	185.05 \pm 24.63
	AG	32	41.88 \pm 4.17	53.55 \pm 6.13	76.71 \pm 8.87	180.22 \pm 19.07
	GG	7	40.71 \pm 2.30	54.20 \pm 4.54	76.34 \pm 5.72	185.14 \pm 28.44
P			0.078	0.719	0.303	0.930
FABP4	AG	25	42.47 \pm 1.31	52.72 \pm 1.51	72.87 \pm 1.98	185.25 \pm 5.71
	GG	34	41.69 \pm 1.33	52.54 \pm 1.53	73.38 \pm 2.00	186.83 \pm 5.78
	P		0.603	0.918	0.819	0.807
STAT5A	CC	28	42.23 \pm 1.33	52.68 \pm 1.53	75.59 \pm 1.90 ^a	184.38 \pm 5.78
	CT	31	41.95 \pm 1.34	52.59 \pm 1.54	70.61 \pm 1.91 ^b	187.69 \pm 5.81
	P		0.858	0.959	0.028	0.622

Statistical differences among genotypes were shown as: ^a and ^b P<0.05

bp) for AG genotype, two bands (367 and 123 bp) for GG genotype (Fig. 1b). The AG genotype was found to be the highest frequency whereas GG genotype was found to be the lowest frequency in our animals. The A allele frequency was found higher than G allele (Table 2).

The 399 bp products were obtained after PCR process and they were digested by *HinIII* restriction enzyme for *FABP4* gene. At the end of digestion, it was observed one band (399 bp) for GG genotype, three bands (399, 302 and 97 bp) for AG genotype, two bands (302 and 97 bp) for AA genotype (Fig. 1c). The GG genotype was found to be the highest frequency; however, the AA genotype was not found. The G allele frequency was higher than A allele (Table 2).

The 215 bp products were obtained after PCR process and PCR products were digested by *AvaI* restriction enzyme for *STAT5A* gene. At the end of digestion, it was observed one band (215 bp) for TT genotype, three bands (215, 181 and 34 bp) for CT genotype, two bands (181 and 34 bp) for CC genotype (Fig. 1d). The CT genotype was found to be the highest frequency, and the TT genotype was not detected. The C allele frequency was higher than T allele (Table 2).

Association was found between *GH* and *WW*. The calf with *GH*-LL genotype had higher *WW* than other genotype (P<0.05) (Table 3). *STAT5A* was found associated with *RW*. For *STAT5A* genotype CC had highest *RW* compared to other genotype CT (P<0.05) (Table 3). Additionally, prospective associations were identified between *GH*-LL and higher *SW*

Table 4. Least squares means and standard errors for body lengths in calves according to *GH*, *MYF5*, *FABP4* and *STAT5A* genotypes

Gene	Genotype	N	Traits			
			Body Length at Birth (cm) Mean (±SE)	Body Length at Weaning Old (cm) Mean (±SE)	Body Length at Milk Replacer Feed Old (cm) Mean (±SE)	Body Length at 180 th Day (cm) Mean (±SE)
GH	LL	54	67.39±0.797	74.35±0.68	78.91±0.67	106.74±1.46
	LV	5	69.43±1.903	71.76±1.62	76.56±1.60	99.90±3.48
P			0.306	0.128	0.161	0.063
MYF5	AA	20	68.70±4.13 ^a	74.15±4.11	78.35±3.36	106.65±6.72
	AG	32	67.03±2.82 ^{ab}	74.31±2.40	78.59±3.36	106.34±6.37
	GG	7	65.29±3.45 ^b	74.86±3.08	79.57±3.36	108.86±6.41
P			0.035	0.744	0.623	0.531
FABP4	AG	25	67.37±0.93	74.11±0.80	78.44±0.79	106.14±1.74
	GG	34	67.89±0.94	74.01±0.81	78.86±0.80	105.77±1.77
P			0.620	0.914	0.641	0.850
STAT5A	CC	28	67.98±0.94	73.97±0.81	78.62±0.80	104.28±1.72
	CT	31	67.25±0.94	74.15±0.82	78.66±0.80	107.66±1.73
P			0.505	0.850	0.966	0.095

Statistical differences among genotypes were shown as: ^a and ^b P<0.05

Table 5. Least squares means and standard errors for HG in calves according to *GH*, *MYF5*, *FABP4* and *STAT5A* genotypes

Gene	Genotype	N	Traits			
			HG at Birth (cm) Mean (±SE)	HG at Weaning Old (cm) Mean (±SE)	HG at Milk Replacer Feed Old (cm) Mean (±SE)	HG at 180 th Day (cm) Mean (±SE)
GH	LL	54	80.56±0.71	87.22±1.04	92.01±1.11	131.81±1.29
	LV	5	77.43±1.70	82.38±2.48	86.53±2.64	125.98±3.07
P			0.081	0.065	0.050	0.072
MYF5	AA	20	81.70±4.40	88.60±4.74	92.30±4.54	131.65±8.280
	AG	32	80.63±2.85	87.00±4.72	90.78±5.01	130.84±5.036
	GG	7	80.14±2.73	88.00±5.57	93.43±6.99	131.86±7.105
P			0.541	0.613	0.483	0.935
FABP4	AG	25	80.74±0.84	86.82±1.24	91.53±1.33	131.16±1.54
	GG	34	79.65±0.85	86.51±1.26	91.24±1.35	131.13±1.55
P			0.253	0.826	0.847	0.988
STAT5A	CC	28	80.12±0.86	85.77±1.24	90.32±1.32	130.39±1.55
	CT	31	80.29±0.86	87.57±1.24	92.47±1.33	131.91±1.55
P			0.863	0.213	0.165	0.396

(P=0.065) and *MYF5*-AA and higher BW (P=0.078) (Table 3). Calves with *MYF5*-AA genotype had highest body length at birth than other genotypes (P<0.05) (Table 4). Additionally, prospective association was identified between *GH*-LL (P=0.063) and *STAT5A*-CT (P=0.095) and longer body length at 180th day (Table 4). Prospective association was identified between *GH* genotypes and hearth girth (HG) at different ages (Table 5). *GH*-LL genotype was found associated with longer HG in all measured ages (Table 5). No association was found between *FABP4* genotypes and investigated traits in this study.

DISCUSSION

To the best of our knowledge there is only few data about association between *GH*, *MYF5*, *FABP4*, and *STAT5A* genes polymorphisms with live weight and morphological measurements in Holstein cattle breed. In the present study genotype frequency of *GH*-LL was found most abundant compared to other genotypes (Table 2). Similar results were observed in other Holstein populations from different countries. Frequency of genotype VV was found low or none in different Holstein populations [30-33].

These findings are consistent with our results, obtained in this study. Among investigated genes in our study, most promising association results were obtained from *GH* gene regarding live weight and body measurement traits in Holstein young bulls (Table 3, 4 and 5). *GH*-LL genotype was associated with higher weaning weight compared to *GH*-LV (Table 3). To the best of our knowledge, no association study was found regarding *GH*-*Alul* polymorphism and WW in Holstein cattle breed. However, association between *GH*-*Alul* polymorphism and weaning weight investigated in Canchim cattle [34] and Charolais and Zebu hybrid together with Charolais, Nelore and Canchim hybrid [35] and in all investigated genotypes no association was observed between *GH*-*Alul* polymorphism and WW. Additionally, *GH*-*Alul* polymorphism was found prospectively associated with SW (Table 3) and 180th day body length (Table 4). In literature it was stated that *GH*-*Alul* genotypes effects the growth and feed intake levels in cattle [36,37]. Additionally, it was shown that GH concentration is affected from *GH*-*Alul* genotypes and age in cattle [36]. In the present study *GH*-*Alul* genotypes found only associated with WW, SW and body length at 180th day but not with the other traits (Table 3 and 4). We thought that this may be due to effects of environmental factors such as feed intake on *GH* genotypes. According to our literature search no study was found investigating effects of *GH*-*Alul* genotypes with heart girth measurement. Heinrichs et al. [38] reported estimation of dairy heifer body weight from heart-girth measurements by using equations or tables. In the present study, heart girth was found prospectively associated with *GH*-*Alul* genotypes in all periods (Table 5), therefore it is thought that *GH*-*Alul* polymorphisms might be used in selecting cattle with longer heart girth which may cause higher body weights.

In the current study all three genotypes for *MYF5*-*TaqI* polymorphism were observed in the investigated Holstein population (Table 2). *MYF5*-*TaqI* AG genotype frequency was found most abundant compared to other two genotypes (Table 2). Similar results were observed in different cattle breeds in Korean [39] and Turkish native cattle breeds [26] that genotype AG was found more abundant than two other genotype. There are not so many researches about association of *MYF5* genotypes with body weight in Holstein cattle. Nasr et al. [40] found an association between body weight and another *MYF5*-*TaqI* site. However, in the current study we only found association with birth weight in young Holstein bulls. As a member of myogenic regulatory factors, *MYF5* is an important transcription factor for skeletal myogenesis in mammalian embryos [41]. Because of its crucial role in embryonic growth, we thought that *MYF5* variants may affect birth weight and body measurements at birth. Supporting this idea above, in the present study, *MYF5*-*TaqI* polymorphism was found associated with body length at birth (Table 4) and prospectively associated with BW (Table 3). Similarly, an association between *MYF5*-*TaqI* polymorphism and birth weight was reported in a Canadian

commercial beef cattle population developed from various cattle breeds [13]. However, no association found in terms of same polymorphism in Korean and Chinese native cattle populations [39,42]. Our anticipation on inconsistent association results may resulted from breed differences. Because distinct genetic difference between European and Asian cattle breeds has been shown in literature [43].

In our study *MYF5*-*TaqI* GG genotype calves had lower birth weight and body length compared to other genotypes, however, animals those have low birth weight and body length are compensating these traits and reaching to similar body weight and length of animals with AA and AG genotypes. Probability of perinatal mortality was found higher in calves with heavier birth weight and birth weights above 42 kg shown as at high risk of perinatal mortality [44]. Therefore, selecting animals with GG genotype may lead us to select calves with lower risk of perinatal mortality without any economic loss at the slaughter age.

For *FABP4*-*HinII* polymorphism genotype GG was the most abundant and genotype AA was none in the Holstein population (Table 2). Similarly, in the Korean Hanwoo cattle genotype GG was also found as common genotype and frequency of AA was found as lowest (0.07) [18]. And no association was observed between *FABP4*-*HinII* polymorphism and investigated phenotypes in Holstein young bulls.

In the present study, no *STAT5A*-*Aval* TT genotype was observed in Holstein male calves. Similarly, this genotype was not found also in Polish Black-and-White [23] and Holstein cattle breeds [37]. However, TT genotype was observed in Podolica bulls [28] and Polish native cattle [23]. *STAT5A*-*Aval* polymorphism was found associated with RW in our investigated animals (Table 3), CC genotype had shown higher RW compared to CT genotype. In consistent with our results, *STAT5A*-*Aval* CC genotype was found favorable for body weight in different ages from different cattle breeds [23,28,37].

Taken together, results in our study provide evidence that interaction between *GH*-*Alul* and *MYF5*-*TaqI* polymorphisms have potential effects for growth and morphological traits which are correlated with economical traits in Holstein cattle. Further studies are ultimately needed to use the SNPs of these two candidate genes in larger populations for genomic selection and investigate other polymorphisms those are linked with *GH*-*Alul* and *MYF5*-*TaqI* SNPs for growth related traits in cattle.

CONFLICTS OF INTEREST

The authors declare no conflict of interest

ACKNOWLEDGMENTS

This work was supported by Erciyes University, Scientific Research Project Coordination Unit (Grant number FCD-

2013-4640). Authors are indebted to Saray Çiftliği-Kayseri farm management for providing animals and helps during collection of phenotypes.

AUTHORS' CONTRIBUTIONS

MUC: Made experimental design and wrote the manuscript. JMK: Collected phenotype and samples and performed statistical analysis. KA: Performed laboratory experiments, responsible for chemical and reagents. EGA: Performed laboratory experiments. MK: Collected phenotype and samples. BA: Made experimental design and wrote the manuscript

REFERENCES

- Henchion M, McCarthy M, Resconi VC, Troy D:** Meat consumption: Trends and quality matters. *Meat Sci*, 98, 561-568, 2014. DOI: 10.1016/j.meatsci.2014.06.007
- Jonas E, de Koning DJ:** Genomic selection needs to be carefully assessed to meet specific requirements in livestock breeding programs. *Front Genet*, 6, 49, 2015. DOI: 10.3389/fgene.2015.00049
- Ardiyanti A, Abe F, Kobashikawa H, Hirayama T, Sugino T, Suzuki K, Katoh K:** Plasma hormone and metabolite concentrations involved in the somatotropic axis of Japanese Black heifers in association with growth hormone gene polymorphism. *Domest Anim Endocrinol*, 37, 243-249, 2009. DOI: 10.1016/j.domaniend.2009.07.002
- Hradecká E, Čitek J, Panicke L, Řehout V, Hanusová L:** The relation of *GH1*, *GHR* and *DGAT1* polymorphisms with estimated breeding values for milk production traits of German Holstein sires. *Czech J Anim Sci*, 53, 238-246, 2008. DOI: 10.17221/362-CJAS
- Hediger R, Johnson SE, Barendse W, Drinkwater RD, Moore SS, Hetzel J:** Assignment of the growth hormone gene locus to 19q26-qter in cattle and to 11q25-qter in sheep by in situ hybridization. *Genomics*, 8, 171-174, 1990. DOI: 10.1016/0888-7543(90)90241-L
- Woychik RP, Camper SA, Lyons RH, Horowitz S, Goodwin EC, Rottman FM:** Cloning and nucleotide sequencing of the bovine growth hormone gene. *Nucleic Acids Res*, 10, 7197-7210, 1982. DOI: 10.1093/nar/10.22.7197
- Sodhi M, Mukesh M, Prakash B, Mishra BP, Sobti RC, Singh KP, Ahlawat SPS:** *MspI* allelic pattern of bovine growth hormone gene in Indian Zebu cattle (*Bos indicus*) breeds. *Biochem Genet*, 45, 145-153, 2007. DOI: 10.1007/s10528-006-9068-4
- Tatsuda K, Oka A, Iwamoto E, Kuroda Y, Takeshita H, Kataoka H, Kouno S:** Relationship of the bovine growth hormone gene to carcass traits in Japanese Black cattle. *J Anim Breed Genet*, 125, 45-49, 2008. DOI: 10.1111/j.1439-0388.2007.00688.x
- Buckingham M:** Skeletal muscle formation in vertebrates. *Curr Opin Genetics Dev*, 11, 440-448, 2001. DOI: 10.1016/S0959-437X(00)00215-X
- te Pas MFW, Soumilion A, Harders FL, Verburg FJ, van den Bosch TJ, Galesloot P, Meuwissen THE:** Influences of myogenin genotypes on birth weight, growth rate, carcass weight, backfat thickness, and lean weight of pigs. *J Anim Sci*, 77, 2352-2356, 1999. DOI: 10.2527/1999.7792352x
- Soumilion A, Erkens JHF, Lenstra JA, Rettenberger G, te Pas MFW:** Genetic variation in the porcine myogenin gene locus. *Mamm Genome*, 8, 564-568, 1997. DOI: 10.1007/s003359900504
- Robakowska-Hyżorek D, Oprządek J, Żelazowska B, Olbromski R, Zwierzchowski L:** Effect of the g.-723G→T Polymorphism in the bovine myogenin factor 5 (MYF5) gene promoter region on gene transcript level in the longissimus dorsi muscle and on meat traits of Polish Holstein-Friesian cattle. *Biochem Genet*, 48, 450-464, 2010. DOI: 10.1007/s10528-009-9328-1
- Li C, Basarab J, Snelling WM, Benkel B, Murdoch B, Hansen C, Moore SS:** Assessment of positional candidate genes MYF5 and IGF1 for growth on bovine chromosome 5 in commercial lines of *Bos taurus*. *J Anim Sci*, 82, 1-7, 2004. DOI: 10.2527/2004.8211
- Verner J, Humpolíček P, Knoll A:** Impact of *MYOD* family genes on pork traits in Large White and Landrace pigs. *J Anim Breed Genet*, 124, 81-85, 2007. DOI: 10.1111/j.1439-0388.2007.00639.x
- Chmurzyńska A:** The multigene family of fatty acid-binding proteins (FABPs): Function, structure and polymorphism. *J Appl Genet*, 47, 39-48, 2006. DOI: 10.1007/BF03194597
- Damcott CM, Moffett SP, Feingold E, Barmada MM, Marshall JA, Hamman RF, Ferrell RE:** Genetic variation in fatty acid-binding protein-4 and peroxisome proliferator-activated receptor gamma interactively influence insulin sensitivity and body composition in males. *Metabolism*, 53, 303-309, 2004. DOI: 10.1016/j.metabol.2003.10.010
- Matarese V, Bernlohr DA:** Purification of murine adipocyte lipid-binding protein. Characterization as a fatty acid- and retinoic acid-binding protein. *J Biol Chem*, 263, 14544-14551, 1988.
- Shin SC, Heo JP, Chung ER:** Genetic variants of the FABP4 gene are associated with marbling scores and meat quality grades in Hanwoo (Korean cattle). *Mol Biol Rep*, 39, 5323-5330, 2012. DOI: 10.1007/s11033-011-1331-z
- Cho S, Park TS, Yoon D-H, Cheong HS, Namgoong S, Park BL, Lee HW, Han CS, Kim EM, Cheong IC, Kim H, Shin HD:** Identification of genetic polymorphisms in FABP3 and FABP4 and putative association with back fat thickness in Korean native cattle. *BMB Rep*, 41, 29-34, 2008. DOI: 10.5483/BMBRep.2008.41.1.029
- Lee SH, van der Werf JHH, Lee SH, Park EW, Oh SJ, Gibson JP, Thompson JM:** Genetic polymorphisms of the bovine fatty acid binding protein 4 gene are significantly associated with marbling and carcass weight in Hanwoo (Korean Cattle). *Anim Genet*, 41, 442-444, 2010. DOI: 10.1111/j.1365-2052.2010.02024.x
- Darnell JE, Kerr IM, Stark GR:** Jak-STAT pathways and transcriptional activation in response to IFNs and other extracellular signaling proteins. *Science*, 264, 1415-1421, 1994. DOI: 10.1126/science.8197455
- Seyfert H-M, Pitra C, Meyer L, Brunner RM, Wheeler TT, Molenaar A, McCracken JY, Herrman J, Thiesen HJ, Schwerin M:** Molecular characterization of STAT5A- and STAT5B-encoding genes reveals extended intragenic sequence homogeneity in cattle and mouse and different degrees of divergent evolution of various domains. *J Mol Evol*, 50, 550-561, 2000. DOI: 10.1007/s002390010058
- Flisikowski K, Oprządek J, Dymnicki E, Zwierzchowski L:** New polymorphism in the bovine *STAT5A* gene and its association with meat production traits in beef cattle. *Anim Sci Pap Rep*, 21, 147-157, 2003.
- Brown JE, Brown CJ, Butts WT:** Evaluating relationship among immature measures of size, shape and performance on beef bulls. I. Principle component as measure of size and shape in young Hereford and Angus bulls. *J Anim Sci*, 36, 1010-1020, 1973. DOI: 10.2527/jas1973.3661010x
- Schlee P, Grami R, Rottmann O, Pirchner F:** Influence of growth hormone genotype on breeding value of Simmental bulls. *J Anim Breed Genet*, 111, 253-256, 1994. DOI: 10.1111/j.1439-0388.1994.tb00465.x
- Sahin C, Akyuz B:** Detection of MYF5 gene polymorphism with PCR-RFLP method in five cattle breeds breeding in Turkey. *Mediterr Agric Sci*, 30, 35-38, 2017.
- Maharani D, Jung Y, Jung WY, Jo C, Ryoo SH, Lee SH, Yeon SH, Lee JH:** Association of five candidate genes with fatty acid composition in Korean cattle. *Mol Biol Rep*, 39, 6113-6121, 2012. DOI: 10.1007/s11033-011-1426-6
- Dario C, Selvaggi M, Carnicella D, Bufano G:** STAT5A/Aval polymorphism in Podolica bulls and its effect on growth performance traits. *Livest Sci*, 123, 83-87, 2009. DOI: 10.1016/j.livsci.2008.10.011
- Rodriguez S, Gaunt TR, Day INM:** Hardy-Weinberg equilibrium testing of biological ascertainment for Mendelian randomization studies. *Am J Epidemiol*, 169, 505-514, 2009. DOI: 10.1093/aje/kwn359
- Grochowska R, Sørensen P, Zwierzchowski L, Snochowski M, Løvendahl P:** Genetic variation in stimulated GH release and in IGF-I of young dairy cattle and their associations with the leucine/valine

polymorphism in the GH gene. *J Anim Sci*, 79, 470-476, 2001. DOI: 10.2527/2001.792470x

- 31. Heidari M, Azari MA, Hasani S, Khanahmadi A, Zerehdaran S:** Effect of polymorphic variants of GH, PIT-1, and β -LG genes on milk production of Holstein cows. *Russ J Genet*, 48, 417-421, 2012. DOI: 10.1134/S1022795412040060
- 32. Kovács K, Völgyi-Csík J, Zsolnai A, Györkös I, Fésüs L:** Associations between the Alul polymorphism of growth hormone gene and production and reproduction traits in a Hungarian Holstein-Friesian bull dam population. *Arch Tierz*, 49, 236-249, 2006. DOI: 10.5194/aab-49-236-2006
- 33. Misrianti R, Anggraeni A, Andreas E, Sumantri C:** Growth hormone gene polymorphism and its association with partial cumulative milk yields of Holstein Friesian dairy cattle. *Media Peternakan*, 35, 145-151, 2012. DOI: 10.5398/medpet.2012.35.3.145
- 34. Grossi Ddo A, Buzanskas ME, Grupioni NV, de Paz CCP, Regitano LC, de Alencar MM, Schenkel FS, Munari DP:** Effect of IGF1, GH, and PIT1 markers on the genetic parameters of growth and reproduction traits in Canchim cattle. *Mol Biol Rep*, 42, 245-251, 2015. DOI: 10.1007/s11033-014-3767-4
- 35. Pereira AP, de Alencar MM, de Oliveira HN, de A Regitano LC:** Association of GH and IGF-1 polymorphisms with growth traits in a synthetic beef cattle breed. *Genet Mol Biol*, 28, 230-236, 2005. DOI: 10.1590/S1415-47572005000200009
- 36. Oka A, Iwaki F, Iwamoto E, Tatsuda K:** Effects of growth rate during the early fattening period on growth, carcass characteristics and circulating hormones in the different growth hormone genotypes of Japanese Black steers. *Anim Sci J*, 78, 142-150, 2007. DOI: 10.1111/j.1740-0929.2007.00418.x
- 37. Oprządek J, Flisikowski K, Zwierzchowski L, Juszcuk-Kubiak E, Rosochacki S, Dymnicki E:** Associations between polymorphism of some candidate genes and growth rates, feed intake and utilisation, slaughter indicators and meat quality in cattle. *Arch Tierz*, 48, 81-87, 2005.
- 38. Heinrichs AJ, Erb HN, Rogers GW, Cooper JB, Jones CM:** Variability in Holstein heifer heart-girth measurements and comparison of prediction equations for live weight. *Prev Vet Med*, 78, 333-338, 2007. DOI: 10.1016/j.prevetmed.2006.11.002
- 39. Chung ER, Kim WT:** Association of SNP marker in IGF-I and MYF5 candidate genes with growth traits in Korean cattle. *Asian-Australas J Anim Sci*, 18, 1061-1065, 2005. DOI: 10.5713/ajas.2005.1061
- 40. Nasr SM, Ateya AI, Sadek KM, Radwan HA:** *TaqI* polymorphism in *MYF5* gene and its association with body weight in Friesian bull calves. *Asian J Anim Vet Adv*, 11 (7): 429-433, 2016. DOI: 10.3923/ajava.2016.429.433
- 41. Francetic T, Li Q:** Skeletal myogenesis and *MYF5* activation. *Transcription*, 2, 109-114. 2011. DOI: 10.4161/trns.2.3.15829
- 42. Zhang RF, Chen H, Lei CZ, Zhang CL, Lan XY, Zhang YD, Zhang HJ, Bao B, Niu H, Wang XZ:** Association between polymorphisms of MSTN and MYF5 genes and growth traits in three Chinese cattle breeds. *Asian-Australas J Anim Sci*, 20, 1798-1804, 2007. DOI: 10.5713/ajas.2007.1798
- 43. Yonesaka R, Sasazaki S, Yasue H, Niwata S, Inayoshi Y, Mukai F, Mannen H:** Genetic structure and relationships of 16 Asian and European cattle populations using DigiTag2 assay. *Anim Sci J*, 87, 190-196, 2016. DOI: 10.1111/asj.12416
- 44. Johanson JM, Berger PJ:** Birth weight as a predictor of calving ease and perinatal mortality in Holstein cattle. *J Dairy Sci*, 86, 3745-3755, 2003. DOI: 10.3168/jds.S0022-0302(03)73981-2