

The Impact of an Essential Oil Mixture on Growth Performance and Intestinal Histology in Native Turkish Geese (*Anser anser*)

Mükremin ÖLMEZ^{1,a} Tarkan ŞAHİN^{1,b} Özlem KARADAĞOĞLU^{2,c} Ebru KARADAĞ SARI^{3,d}
Serpil ADIGÜZEL IŞIK^{4,e} Turgut KIRMIZIBAYRAK^{4,f} Mehmet Akif YÖRÜK^{5,g}

¹ Department of Animal Nutrition and Nutritional Diseases, Faculty of Veterinary Medicine, Kafkas University, TR-36100 Kars - TURKEY

² Department of Agricultural and Animal Production, Kars Vocational High School, Kafkas University, TR-36100 Kars - TURKEY

³ Department of Histology and Embryology, Faculty of Veterinary Medicine, Kafkas University, TR-36100 Kars - TURKEY

⁴ Department of Animal Breeding and Husbandry, Faculty of Veterinary Medicine, Kafkas University, TR-36100 Kars - TURKEY

⁵ Department of Animal Nutrition and Nutritional Diseases Faculty of Veterinary Medicine, Atatürk University, TR-25240 Erzurum - TURKEY

ORCID: ^a 0000-0002-5003-3383; ^b 0000-0003-0155-2707; ^c 0000-0002-5917-9565; ^d 0000-0001-7581-6109; ^e 0000-0002-4456-8779

^f 0000-0003-3626-6137; ^g 0000-0002-5833-9803

Article ID: KVFD-2020-24070 Received: 12.02.2020 Accepted: 23.06.2020 Published Online: 04.07.2020

How to Cite This Article

Ölmez M, Şahin T, Karadağoğlu Ö, Karadağ Sarı E, Adigüzel Işık S, Kırmızıbayrak T, Yörük MA: The impact of an essential oil mixture on growth performance and intestinal histology in native Turkish geese (*Anser anser*). *Kafkas Univ Vet Fak Derg*, 26 (5): 625-631, 2020. DOI: 10.9775/kvfd.2020.24070

Abstract

This study investigated the impact of an essential oil (EO) mixture (Biomin) on growth performance, carcass yield, visceral organ weights, and duodenal crypt depth in native Turkish geese (*Anser anser*). Sixty 1-day-old goslings constituted the animal material of the trial. Goslings were divided into three treatment groups, each consisting of four subgroups having five animals each. Animals were homogeneously distributed among the groups after being weighed and identified. The three treatment groups received the essential oil mixture at rates of 0.0% (without EO), 0.1% and 0.2%, respectively. The animals were given concentrate feed (21.93% CP and 3010 kcal/kg ME) for the first 2 weeks, and were both grazed and provided with barley meal for the next 8 weeks. The trial was continued for a period of 10 weeks. Results showed that the essential oil mixture had no significant effect on live weight gain, feed intake, feed conversion rate, carcass yield, and heart-liver-gizzard weights ($P>0.05$). On the other hand, the crypt depths of the EO 0.1% and EO 0.2% groups were greater than that of the EO 0.0% group ($P<0.05$). This study demonstrated that EO supplementation had no impact on performance, but increased duodenal crypt depth in native Turkish geese.

Keywords: Crypt depth, Essential oil, Native Turkish goose, Performance

Yerli Türk Kazlarında (*Anser anser*) Esansiyel Yağ Karışımının Büyüme Performansı ve Bağırsak Histolojisi Üzerine Etkisi

Öz

Bu çalışmada esansiyel yağ (EY) karışımının (Biomin) yerli Türk kazlarında besi performansı, karkas randımanı, bazı organ ağırlıkları ve duodenum kript derinliği üzerine etkisi incelenmiştir. Hayvan materyali olarak 60 adet 1 günlük Yerli Türk kazı (*Anser anser*) kullanılmıştır. Kazlar üç deneme grubuna ve her birinde beş hayvan bulunan dört alt gruba ayrılmıştır. Hayvanlar tartılarak ve ayaklarından numaralandırılarak homojen dağıtılmıştır. Esansiyel yağ, gruplara sırasıyla; %0.0 (EY'siz), %0.1 ve %0.2 düzeylerinde verilmiştir. Hayvanlar iki hafta konsantre yemle (%21.93 HP ve 3010 kcal/kg ME) sekiz hafta mera+arpa karmasıyla beslenmiştir. Deneme 10 hafta sürmüştür. Araştırmada esansiyel yağ karışımının canlı ağırlık artışı, yem tüketimi, yem dönüşüm oranı, karkas randımanı, kalp, karaciğer, taşlık ağırlıkları üzerine anlamlı bir etkisi olmamıştır ($P>0.05$). Diğer yandan EY %0.1 ve %0.2 gruplarındaki hayvanlara ait kript derinlikleri, EY %0.0 grubuna göre daha yüksek çıkmıştır ($P<0.05$). Yapılan bu araştırma yerli Türk kazlarında EY karışımının hayvanların performansına herhangi bir etki yapmazken, duodenum kript derinliğini artırdığını göstermiştir.

Anahtar sözcükler: Esansiyel yağ, Kript derinliği, Performans, Yerli Türk kazı



Correspondence



+90 474 2426807-5116



mukremin.olmez@hotmail.com

INTRODUCTION

The intensive and indiscriminate use of antibiotics in the poultry sector has led to antibiotic residues in food products of animal origin, and the emergence of cross-resistant bacteria. These risks, which are both directly and indirectly linked to human health, have impelled researchers to investigate alternative growth factors [1]. Since the ban of the use of antibiotic growth promoters (AGPs) in food animals in the European Union (EU) complying with EU law, aromatic herbs and herbal products (oil, powder) have attracted the attention of the poultry sector as feed additives alternative to antibiotics [2-5]. Essential oils (EOs) are natural and non-residual alternative feed additives, which are obtained from aromatic herbs by various methods and are used as flavourings, appetite-stimulants, digestive stimulants and performance enhancers [6]. The type and level of phenolic compounds found in essential oils vary with the aromatic herbs the oils are derived from, and elicit various activities (i.e. antioxidant, antimicrobial, antifungal) [7-9]. Dietary supplementation with essential oils increases the productive capacity of poultry species by increasing the activity of digestive enzymes and eliminating pathogenic microorganisms [10,11]. Previous research has shown that carvacrol essential oils increase villus height and the villus height/crypt depth ratio, which in return improves digestive capacity [12,13]. Furthermore, rosemary oil essential oils have also been reported to increase the digestion of fat and fat-soluble vitamins by enabling the binding of bile acids to substances, and thereby, to improve the feed conversion rate [14]. Thus, it is considered that the combined use of synergistic essential oils, which are derived from different aromatic herbs and show different effects, even at low but appropriate doses, could improve animal production [15].

People have increasingly demanded poultry meat as an alternative source of animal protein [16]. Furthermore, goose meat is healthy, wholesome, and contains low cholesterol levels. Therefore, compared to other poultry species, geese are well adapted to cold climate conditions, more resistant to diseases, and do not require to be housed in costly pens. Although geese are classified as waterfowl, they are capable of living in drylands. Geese make good use of pasture plants and can be fed by grazing alone [17].

This study was aimed at the investigation of the impact of two different doses of an EO mixture on performance and intestinal histomorphology in native Turkish geese.

MATERIAL and METHODS

Ethical Approval

This study was conducted according to the approval (KAU-HADYEK/2019-118) granted by the Local Ethics Board for Experimental Animals of Kafkas University.

Animals and Treatment Design

The study was carried out at the premises of the Research Farm of Kafkas University. Sixty 1-day-old native Turkish goslings (*Anser anser*) of both sexes constituted the animal material. After hatching, the goslings were weighed, identified by unique numbers, and assigned to three groups. Each 20 animals in groups were divided into four subgroups containing five animals. The animals were homogeneously distributed among the groups by weight (32.76 ± 0.01 g). The essential oil mixture was given to the three treatment groups at concentrations of 0%, 0.1% and 0.2%, respectively, in drinking water. During the first two weeks of the study, the animals were housed in cages under favourable environmental conditions and were fed on commercial goose starter ration containing 21.93% of crude protein (CP) and 3010 kcal/kg of metabolizable energy (ME) [18]. As from the third week, the animals were raised on the floor and acclimatized to grazing. The minimum area provided to each animal raised on the floor was 0.5 m². As of the third week, the animals were not only grazed but also provided with a barley meal (BM). Feed and water were provided *ad libitum*. The study was continued for ten weeks. The nutrient and chemical compositions of the starter ration, barley meal and pasture plants that the geese were fed on are presented in [Table 1](#). The flora of the pasture consists of 64% Gramineae, 23% Leguminosae, and 13% other species. The treatment groups received different doses of essential oil and aromatic herb mixture containing peppermint oil, garlic oil, aniseed oil, fennel oil, cinnamon and cumin. The mixture was a commercial product supplied from Austria (BIOMIN GmbH).

Performance

Starting from the day of hatching to the day of slaughter, the animals were weighed individually on a weekly basis to determine their live weight gain. The feed intake (first two weeks concentrate feed-next eight weeks barley meal) of the animals was also calculated on a weekly basis. Feed conversion rates (FCR) were determined by calculating the ratio of feed intake to live weight gain.

Carcass Traits

Ten-week-old geese were fasted for 12 h prior to slaughter. Then, six animals, representative of the live weight of the groups, were selected from each group and were sacrificed by slaughtering the head at the occipital-atlantoaxial articulation. The birds were bled for 10-15 min and defeathered with machines. Following evisceration, the liver, gizzard and heart were weighed individually. The hot carcass weights of the unchilled carcasses were determined.

Histological Procedure

The duodenal tissue samples taken from the geese were firstly fixed in 10% formalin solution for 24 h. Then routine

Table 1. Composition and nutrient ingredients of the feeds		
Ingredients (starter ration)	%	
Corn, yellow	64.40	
Soybean meal, 48% CP	21.46	
Fish meal, 64% CP	7.00	
Sunflower meal, 32% CP	4.80	
Vegetable oil	0.55	
Limestone	0.65	
Dicalcium phosphate	0.35	
DL-methionine	0.08	
L-Lysine HCl	0.15	
L-Threonine	0.06	
Salt	0.25	
Vit-Min Mix*	0.25	
Nutrient Values		
Dry matter (%)	89.72	
Crude protein (%)	21.93	
Ca (%)	0.87	
Available P (%)	0.44	
Na (%)	0.16	
Met+Cys (%)	0.90	
Lysine (%)	1.34	
Threonine (%)	0.91	
Tryptophan (%)	0.26	
Metabolic Energy, (kcal/kg)	3010	
Chemical Analysis		
Items	Barley Meal	Pasture
Dry matter (%)	88.11	94.55
Crude protein (%)	12.28	7.96
Crude fat (%)	2.18	1.65
Crude ash (%)	3.73	7.43
Crude fiber (%)	5.41	42.88
Neutral detergent fiber (%)	23.23	60.30
Acid detergent fiber (%)	6.35	48.27
Lignin (%)	0.57	9.62
Non-protein nitrogen (%)	76.41	40.09
Non-fiber carbohydrate (%)	58.59	22.67
Hemicellulose (%)	16.88	12.04
Starch (%)	55.23	-
Metabolic Energy Poultry kcal/kg	3097.05	-

* Vit-Min mix: Vit A: 10.000IU, Vit D₃: 4.000IU, Fe (iron sulfate monohydrate): 30 mg; I (calcium iodine anhydride): 1.5 mg, Co (cobalt carbonate monohydrate): 0.5 mg, Cu (copper sulfate pentahydrate): 5 mg, Mn (manganese oxide): 80 mg, Zn (zinc oxide): 80 mg, Se (selenium selenite): 0.3 mg

tissue processing was applied and embedded in paraffin. Five-micron-thick sections were cut from the paraffin blocks and applied Mallory's modified triple staining (Triple) with a view to demonstrate the general structure

Table 2. Effects of EO on performance parameters of geese (Mean±SEM; n=20)				
Periods	EO Groups	LWG (g)	FI (g)	FCR
0-2. weeks	0.0%	33.91±0.98	49.93±0.27	1.48±0.04
	0.1%	33.18±1.53	49.59±0.24	1.50±0.07
	0.2%	32.32±0.17	49.53±0.28	1.53±0.01
P		0.584	0.531	0.680
2-4. weeks	0.0%	51.51±0.79	121.33±0.28	2.36±0.04
	0.1%	51.16±0.26	121.53±0.39	2.38±0.02
	0.2%	51.90±0.86	121.88±0.38	2.35±0.04
P		0.755	0.559	0.688
4-6. weeks	0.0%	45.47±0.78	171.05±0.48	3.76±0.06
	0.1%	45.81±0.67	168.55±0.57	3.68±0.05
	0.2%	49.49±1.17	169.18±0.45	3.67±0.05
P		0.787	0.403	0.423
6-8. weeks	0.0%	49.10±0.69	201.24±0.50	4.27±0.09
	0.1%	50.03±0.71	201.21±0.45	4.13±0.11
	0.2%	48.55±0.66	201.58±0.61	4.08±0.11
P		0.342	0.858	0.424
8-10. weeks	0.0%	48.87±0.32	228.60±0.34	4.66±0.03
	0.1%	47.52±0.85	228.61±0.64	4.82±0.08
	0.2%	48.05±0.70	228.28±0.44	4.75±0.07
P		0.385	0.376	0.279
0-10. weeks	0.0%	45.36±0.25	154.53±0.11	3.40±0.02
	0.1%	45.09±0.40	153.90±0.08	3.41±0.03
	0.2%	45.59±0.17	154.09±0.05	3.29±0.04
P		0.534	0.059	0.635

EO: Essential oil; LWG: Live weight gain; FI: Feed intake; FCR: Feed conversion rate. SEM: Standart error of the mean

of the duodenal tissue. The sections were examined with a light microscope (Olympus BX51, Japan). The crypt depths of the geese in all groups were measured using the ImageJ (LOCI, University of Wisconsin) software.

Statistical Analysis

The differences between the groups for performance, carcass traits, and histological parameters were analysed using the Statistical Package (SPSS portable PASW 18) software by one-way analysis of variance (ANOVA). The paired comparison of the data was performed with Duncan's test. The significance level was accepted as $P < 0.05$.

RESULTS

The effects of the EO mixture on growth performance are shown in Table 2. The results obtained in the present study demonstrated that the EO mixture had no effect on live weight gain, feed intake and feed conversion rate ($P > 0.05$).

Values pertaining to the slaughter and carcass traits of the study groups are presented in Table 3. Accordingly, it was

Table 3. Effects of EO on carcass traits of geese (Mean±SEM; n=6)

EO Groups	Slaughter Weight (g)	Carcass Yield (%)	Liver (g)	Gizzard (g)	Heart (g)
0.0%	3263.4±22.64	71.66±0.51	161.60±2.25	187.60±10.69	24.80±1.36
0.1%	3236.2±52.66	71.22±0.33	161.60±2.94	178.40±6.55	25.80±1.74
0.2%	3173.3±34.84	71.38±0.30	161.40±2.11	170.80±4.90	25.00±1.97
P	0.279	0.724	0.998	0.343	0.909

EO: Essential oil; SEM: Standart error of the mean



**Fig 1.** Goose duodenum tissue. **A:** 0.1% EO group, **B:** 0.2% EO group. **V:** Intestinal villi, **C:** Crypts, **TM:** Tunica muscularis, **TS:** Tunica serosa; Triple staining

Fig 2. Effects of EO on duodenal crypt depths (μm), (Mean±SEM; n=6); ^{a,b} Values in the same column with no common super-script differ significantly ($P<0.05$); EO: Essential oil; SEM: Standart error of the mean



determined that the groups did not differ for the weight at slaughter and carcass yield of the animals ($P>0.05$). Similarly, no significant difference was determined between the groups for liver, gizzard and heart weights ($P>0.05$). The average weight of the liver, which is one of the most valuable products of geese, was determined to be 161.60 g, 161.60 g and 161.40 g, respectively, in the groups that were given 0%, 0.1% and 0.2% EO.

In all groups, the duodenal tissue of the geese was observed to be composed of the innermost tunica mucosa, tunica muscularis consist of smooth muscle cells and the outermost tunica serosa consist of connective tissue layers. The tunica mucosa formed the intestinal villi and crypts (Fig. 1, Fig. 2).

The statistical analysis of the intestinal crypt depths of all groups demonstrated that the 0.0% EO group significantly differed from the 0.1% and 0.2% EO groups ($P<0.05$).

DISCUSSION

No statistically significant difference existed between the EO treatment groups (0.0%, 0.1% and 0.2%) for the live weight gain values between weeks 0-10 of the study. Generally, it is considered that herbal extracts produce favourable results in poultry. However, there are very few studies conducted on the use of aromatic herbs in geese. In these limited number of studies, it has been suggested that, in geese, performance values are not affected by supplementation with aromatic herbs [19,20]. The results of

the present study are in agreement with these previous poultry studies reporting unaffected performance [21-23]. Researchers have also reported that improved environmental conditions and reduced pathogen burden decrease the effects expected from the use of essential oils. Jamroz et al. [24] contrary to reported that 100 mg of an essential oil mixture containing carvacrol, cinnamaldehyde and capsicum increased live weight parameters in broiler chickens. This increase was attributed to cinnamaldehyde, which is the main active substance of cinnamon, increasing the secretion of pancreatic and intestinal enzymes, and capsicum increasing the production of hepatic enzymes, bile, pancreatic enzymes and intestinal lipase, resulting in an increased absorption of nutrients [24,25]. The results of the present study do not agree with some literature reports [26-28]. Some studies suggest that essential oil supplementation shows adverse effects on the performance of poultry [29]. Differences between the results of the present study and previous research have been attributed to different essential oil components having been used and tested, the essential oils found in the composition of mixtures interacting with each other, and EOs having been used at different levels. Furthermore, it is considered that geographical conditions, the timing of harvest, as well as animal- and environment-related factors may also contribute to discrepancies in study results.

Throughout the study period, no significant difference was observed between the groups for weekly feed intake and feed conversion rate values. In previous studies conducted by Baowei et al. [19] and Yaman et al. [20] in geese, it was reported that neither feed intake nor feed conversion rate was affected. Similarly, while Fascina et al. [30] indicated that essential oil mixtures had no effect on growth performance, feed intake and feed conversion rate, Nobakht and Mehmanavaz [31] reported that different fat resources did not cause any change in average feed intake values in broiler chickens. Furthermore, Khaksar et al. [32] reported that thymol essential oil had no effect on feed intake and feed conversion rate in quails. These findings are in agreement with previous studies, which were conducted by Mathlouthi et al. [33], Küçükylmaz et al. [34], and suggested that essential oil mixtures did not affect feed intake and feed conversion rate.

The effects of essential oils arise from the active molecules found in their structure. The results of the present study are similar to those reported by Lee et al. [35] and Günel et al. [36], who suggested that herbal extracts do not affect feed intake in broiler chickens, but contradict with the report of Tekeli et al. [37], who suggested that herbal extracts increase feed intake in broiler chickens. Jamroz et al. [24] reported that the supplementation of maize- and wheat-based rations with an essential oil mixture containing carvacrol, cinnamon oil and black pepper oil increased the feed conversion rate of broiler chickens by 4%. On the other hand, Al-Kasie [38] indicated that supplementation with an

essential oil mixture containing carvacrol and cinnamon oil decreased the feed conversion rate, when given at a dose of 200 ppm ($P < 0.05$).

The end-study carcass traits and visceral organ weights of the groups are presented in Table 3. Accordingly, no statistically significant difference was observed between the groups for slaughter weight, carcass yield and heart, liver and gizzard weights ($P > 0.05$).

The findings obtained in the present study comply with those reported by Muhl and Liebert [39], who suggested that the supplementation of broiler grower rations with phytogetic extracts (5% carvacrol, 3% cinnamaldehyde and 2% *Capsicum oleoresin*) had no effect on carcass yield, and also agree with the report of Şimsek et al. [40], who indicated that mixtures of carvacrol, clove oil and aniseed oil do not have any effect on carcass yield and characteristics in broiler chickens. Furthermore, similar to the present study, upon investigating the effects of different protein sources in geese, Şahin et al. [41] reported to have not detected any difference between the study groups for carcass traits. The similarities between the results of these different studies are attributed to the similarity of the management conditions the animals were exposed to.

The results obtained for carcass yields in the present study differ not only from those reported by Fascina et al. [30] who indicated a significant increase in carcass yield with the supplementation of broiler rations with phytogetic extracts, but also from those reported by Al-Kasie et al. [38] who pointed out to significantly increased carcass yields with the supplementation of broiler rations with essential oil mixtures containing carvacrol and cinnamon oil ($P < 0.01$).

Literature reports have been published, which suggest that herbal feed additives improve slaughter and carcass traits [40,42]. Researchers attribute this positive effect to the appetizing, enzyme-secretagogue and antimicrobial activity of essential oils [43].

The results of the present study are similar to those reported by Criste et al. [44] for gizzard and hearth weights, by Kim et al. [45] for liver weight, by Yeganeparast et al. [22] for hearth weights, and by Salehifar et al. [46] for liver and gizzard weights. Çelik and Şahin [47] determined that an essential oil mixture, which was administered in drinking water and contained peppermint oil, carvacrol, cade oil and rosemary oil did not affect heart, liver and gizzard weights. The visceral organ weight parameters determined in the present study differ from the results of some other previous studies conducted on essential oil supplementation [42,48].

Low crypt depth is one of the main indicators of intestinal health. Peng et al. [49] reported that the supplementation of broiler rations with oregano essential oil reduced the

depth of intestinal crypts, and thereby, increased growth performance. Some other studies have reported that essential oil supplementation does not affect duodenal crypt depth in broiler chickens^[50,51]. In agreement with the present study, Yang et al.^[52] reported that cinnamon oil, when supplemented at a high dose (800 mg/kg), increased duodenal crypt depth, but had no effect on performance parameters. In fact, increased crypt depth was observed not to show any adverse effect on performance parameters in geese.

In conclusion, it has been determined that, in geese, supplementation with different doses of essential oil mixtures increases intestinal crypt depth, but this increase does not affect live weight gain. In view of the limited number of studies in geese, it is considered that further studies need to be carried out in poultry, and particularly in geese, such that different doses of various essential oils are investigated in different slaughtered age.

CONFLICT OF INTEREST

All of the authors declare that they have all participated in the design, execution, and analysis of the paper and that they have approved the final version.

AUTHOR CONTRIBUTIONS

Experimental design, feeding trial and article writing; MÖ and TŞ, Performance parameters calculation; ÖK, Histological analysis; EKS, Statistical Analysis; SAI and TK, Interpretation and editing of results; MAY.

REFERENCES

- Bingöl NT, Karslı MA, Aldemir R, Yılmaz O, Türel İ:** Etçi piliçlerin yemlerine katılan *Plantago major* ekstraktının performans ve karkas özellikleri üzerine etkisi. *YYU Vet Fak Derg*, 21 (1): 49-53, 2010.
- Brenes A, Roura E:** Essential oils in poultry nutrition: Main effects and modes of action. *Anim Feed Sci Technol*, 158 (1-2): 1-14, 2010. DOI: 10.1016/j.anifeeds.2010.03.007
- Karásková K, Suchý P, Straková E:** Current use of phytogetic feed additives in animal nutrition: A review. *Czech J Anim Sci*, 60, 521-530, 2015. DOI: 10.17221/8594-CJAS
- Malekzadeh M, Shakouri MD, Benamar HA:** Effect of thyme species extracts on performance, intestinal morphometry, nutrient digestibility and immune response of broilers. *Kafkas Univ Vet Fak Derg*, 24 (6): 783-790, 2018. DOI: 10.9775/kvfd.2018.19719
- Ahmad S, Khalique A, Pasha TN, Mehmood S, Hussain K, Ahmad S, Rasheed B, Awais MM, Bhatti SA:** Influence of feeding *Moringa oleifera* pods as phytogetic feed additive on performance, blood metabolites, chemical composition and bioactive compounds of breast meat in broiler. *Kafkas Univ Vet Fak Derg*, 24 (2): 195-202, 2018. DOI: 10.9775/kvfd.2017.18616
- Köksal BH, Küçükersan MK:** Effects of humate and vegetable extract mixture supplementation to diets on growth performance, some immunity and serum biochemistry parameters in broiler chickens. *Kafkas Univ Vet Fak Derg*, 18 (1): 103-108, 2012. DOI:10.9775/kvfd.2011.5114
- Inci H, Ozdemir G, Sengul AY, Sogut B, Nursoy H, Sengul T:** Using juniper berry (*Juniperus communis*) as a supplement in Japanese quail diets. *R Bras Zootec*, 45 (5): 230-235, 2016. DOI: 10.1590/S1806-92902016000500004
- Karakullukçu MZ, Kocaoğlu Güçlü B, Kara K, Tugrulay S:** Yumurta tavuğu karma yemlerine ilave edilen bazı esansiyel yağların performans ve yumurta kalitesine etkisi. *İstanbul Univ Vet Fak Derg*, 42, 31-37, 2016. DOI: 10.16988/iuvfd.2016.70557
- Lipiński K, Mazur M, Antoszkiewicz Z, Purwin C:** Polyphenols in monogastric nutrition-A review. *Ann Anim Sci*, 17 (1): 41-58, 2017. DOI: 10.1515/aoas-2016-0042
- Masood A, Qureshi AS, Shahid RU, Jamil H:** Effects of oral administration of essential oil (mix oil) on growth performance and intestinal morphometry of japanese quails (*Coturnix coturnix japonica*). *Pak Vet J*, 2020 (Article in Press). DOI: 10.29261/pakvetj/2020.018
- Lee KW, Kim JS, Oh ST, Kang CW, An BK:** Effects of dietary sanguinarine on growth performance, relative organ weight, cecal microflora, serum cholesterol level and meat quality in broiler chickens. *J Poult Sci*, 52 (1): 15-22, 2015. DOI: 10.2141/jpsa.0140073
- Jerzsele A, Szeker K, Cszizinsky R, Gere E, Jakab C, Mallo JJ, Galfi P:** Efficacy of protected sodium butyrate, a protected blend of essential oils, their combination, and *Bacillus amyloliquefaciens* spore suspension against artificially induced necrotic enteritis in broilers. *Poult Sci*, 91 (4): 837-843, 2012. DOI: 10.3382/ps.2011-01853
- Mehdi Y, Létourneau-Montminy MP, Gaucher ML, Chorfi Y, Suresh G, Rouissi T, Brar SK, Côté C, Ramirez AA, Godbout S:** Use of antibiotics in broiler production: Global impacts and alternatives. *Anim Nutr*, 4 (2): 170-178, 2018. DOI: 10.1016/j.aninu.2018.03.002
- Hashemipour H, Kermanshahi H, Golian A, Veldkamp T:** Effect of thymol and carvacrol feed supplementation on performance, antioxidant enzyme activities, fatty acid composition, digestive enzyme activities, and immune response in broiler chickens. *Poult Sci*, 92 (8): 2059-2069, 2013. DOI: 10.3382/ps.2012-02685
- Sevim B, Cufadar Y:** Etlik piliçlerde karma yeme farklı esansiyel yağlar ve karışımlarının ilavesinin performans ve karkas özellikleri üzerine etkisi. *TURJAF*, 5, 964-968, 2017. DOI: 10.24925/turjaf.v5i8.964-968.1261
- Yıldız G:** Kaz besleme. *Yem Mag Derg*, 26, 60-62, 2000.
- Anonim:** Geese. <http://www.waterfowl.org.uk>; Accessed: 12.03.2020.
- NRC:** Nutrient Requirements of Poultry. Ninth Revised ed., National Academies Press, Washington, DC, 1994.
- Baowei W, Guoqing H, Qiaoli W, Bin Y:** Effects of yeast selenium supplementation on the growth performance, meat quality, immunity, and antioxidant capacity of goose. *J Anim Physiol Anim Nutr*, 95 (4): 440-448, 2011. DOI: 10.1111/j.1439-0396.2010.01070.x
- Yaman H, Ulukanlı Z, Elmali M, Unal Y:** The effect of a fermented probiotic, the kefir, on intestinal flora of poultry domesticated geese (*Anser anser*). *Reveu Med Vet*, 157 (7): 379-386, 2006.
- Bidar N, Hassanabadi A, Moghaddam H, Varidi M, Mohsenzadeh M:** The effect of *Lavandula angustifolia* essential oil on performance, blood metabolites and nutrient digestibility in broiler chickens. *Iran J Appl Anim Sci*, 9 (3): 328-339, 2017.
- Yeganeparast M, Khalili M, Salari J:** Effect of different levels of *Satureja hortensis* essential oil on performance, carcass characteristics, acidity and intestinal microflora population in broilers. *J Cell Anim Biol*, 10 (1): 1-8, 2016. DOI: 10.5897/JCAB2015.0440
- Gharejanloo M, Mehri M, Shirmohammad F:** Effect of different levels of turmeric and rosemary essential oils on performance and oxidative stability of broiler meat. *Iran J Appl Anim Sci*, 7 (4): 655-662, 2017.
- Jamroz D, Wiliczekiewicz A, Wartelecki T, Orda J, Skorupińska J:** Use of active substances of plant origin in chicken diets based on maize and locally grown cereals. *Br Poult Sci*, 46 (4): 485-493, 2005. DOI: 10.1080/00071660500191056
- Platel K, Srinivasan K:** Digestive stimulant action of spices: A myth or reality? *Indian J Med Res*, 119 (5): 167-179, 2004.
- Gumus R, Ercan N, Imik H:** The effect of thyme essential oil (*Thymus vulgaris*) added to quail diets on performance, some blood parameters, and the antioxidative metabolism of the serum and liver tissues. *Braz J Poult Sci*, 19 (2): 297-304, 2017. DOI: 10.1590/1806-9061-2016-0403
- Küçükyılmaz K, Kiyama Z, Akdağ A, Çetinkaya M, Atalay H, Ateş A,**

- Gürsel FE, Bozkurt M:** Effect of lavender (*Lavandula stoechas*) essential oil on growth performance, carcass characteristics, meat quality and antioxidant status of broilers. *S Afr J Anim Sci*, 47 (2): 178-186, 2017.
- 28. Masouri L, Salari S, Sari M, Tabatabaei S, Masouri B:** Effect of feed supplementation with *Satureja khuzistanica* essential oil on performance and physiological parameters of broilers fed on wheat- or maize-based diets. *Br Poult Sci*, 58 (4): 425-434, 2017. DOI: 10.1080/00071668.2017.1327701
- 29. Windisch W, Schedle K, Plitzner C, Kroismayr A:** Use of phytogetic products as feed additives for swine and poultry. *J Anim Sci*, 86 (Suppl. 14): E140-E148, 2008. DOI: 10.2527/jas.2007-0459
- 30. Fascina VB, Sartori JR, Gonzales E, Carvalho FBd, Souza IMGpd, Polycarpo GdV, Stradiotti AC, Pelícia VC:** Phytogetic additives and organic acids in broiler chicken diets. *R Bras Zootec*, 41 (10): 2189-2197, 2012. DOI: 10.1590/S1516-35982012001000008
- 31. Nobakht A, Mehmannaavaz Y:** Investigation the effects of using of *Thymus vulgaris*, *Lamiaceae menthapiperita*, *Oreganum valgare* medicinal plants on performance, egg quality, blood and immunity parameters of laying hens. *Iran J Anim Sci*, 41, 129-136, 2010.
- 32. Khaksar V, Van Krimpen M, Hashemipour H, Pilevar M:** Effects of thyme essential oil on performance, some blood parameters and ileal microflora of Japanese quail. *J Poult Sci*, 49 (2): 106-110, 2012. DOI: 10.2141/jpsa.011089
- 33. Mathlouthi N, Ballet N, Larbier M:** Influence of beta-glucanase supplementation on growth performances and digestive organs weights of broiler chickens fed corn, wheat and barley-based diet. *Int J Poult Sci*, 10, 157-159, 2011. DOI: 10.3923/ijps.2011.157.159
- 34. Küçükylmaz K, Bozkurt M, Çatlı AU, Çınar M, Bintaş E:** Yeme dehidre maya ve humat bileşik katılmasının etlik piliçlerin performans ve bazı kesim özellikleri üzerine etkileri. *Ziraat Fak Derg*, 7 (1): 83-92, 2012.
- 35. Lee KW, Everts H, Kappert HJ, Frehner M, Losa R, Beynen AC:** Effects of dietary essential oil components on growth performance, digestive enzymes and lipid metabolism in female broiler chickens. *Br Poult Sci*, 44 (3): 450-457, 2003. DOI: 10.1080/0007166031000085508
- 36. Gunal M, Yaylı G, Kaya O, Karahan N, Sulak O:** The effects of antibiotic growth promoter, probiotic or organic acid supplementation on performance, intestinal microflora and tissue of broilers. *Int J Poult Sci*, 5 (2): 149-155, 2006. DOI: 10.3923/ijps.2006.149.155
- 37. Tekeli A, Çelik L, Kutlu HR:** Plant extracts; a new rumen moderator in ruminant diets. *Tekirdag Ziraat Fak Derg*, 4, 71-79, 2007.
- 38. Al-Kassie GAM:** Influence of two plant extracts derived from thyme and cinnamon on broiler performance. *Pak Vet J*, 29, 169-173, 2009.
- 39. Muhl A, Liebert F:** Growth, nutrient utilization and threonine requirement of growing chicken fed threonine limiting diets with commercial blends of phytogetic feed additives. *J Poult Sci*, 44, 297-304, 2007. DOI: 10.2141/jpsa.44.297
- 40. Simsek UG, Güler T, Çiftçi M, Ertas ON, Dalkılıç B:** Esans yağ karışımının (kekik, karanfil ve anason) broylerlerde canlı ağırlık, karkas ve etlerin duyuşal özellikleri üzerine etkisi. *YYÜ Vet Fak Derg*, 16 (2): 1-5, 2005.
- 41. Sahin T, Tilki M, Kaya İ, Unal Y, Aksu Elmali D:** Effect of different protein levels for finishing period on fattening performance and carcass traits in native Turkish geese. *J Anim Vet Adv*, 7 (11): 1364-1369, 2008.
- 42. Toghyani M, Toghyani M, Gheisari A, Ghalamkari G, Mohammadrezaei M:** Growth performance, serum biochemistry and blood hematology of broiler chicks fed different levels of black seed (*Nigella sativa*) and peppermint (*Mentha piperita*). *Livest Sci*, 129 (1-3): 173-178, 2010. DOI: 10.1016/j.livsci.2010.01.021
- 43. Tekce E, Gül M:** Effects of *Origanum syriacum* essential oil added in different levels to the diet of broilers under heat stress on performance and intestinal histology. *Eur Poult Sci*, 80, 1-11, 2016. DOI: 10.1399/eps.2016.157
- 44. Criste RD, Panaite TD, Tabuc C, Saracila M, Soica C, Olteanu M:** Effect of oregano and rosehip supplements on broiler (14-35 days) performance, carcass and internal organs development and gut health. *AgroLife Sci J*, 6 (1): 75-83, 2017.
- 45. Kim SJ, Lee KW, Kang CW, An BK:** Growth performance, relative meat and organ weights, cecal microflora, and blood characteristics in broiler chickens fed diets containing different nutrient density with or without essential oils. *Asian Australas J Anim Sci*, 29 (4): 549-554, 2016. DOI: 10.5713/ajas.15.0426
- 46. Salehifar E, Abbasi M, Bahari-Kashani R:** Effects of Myrtle (*Myrtus communis*) essential oil on growth performance, carcass characteristics, intestinal morphology, immune response and blood parameters in broiler chickens. *Livest Sci*, 8, 63-71, 2017.
- 47. Çelik R, Şahin T:** İçme suyuna farklı düzeylerde ilave edilen esansiyel yağ karışımlarının (nane + kekik + ardıç + biberiye) broylerlerde besi performansı, kesim ve karkas özellikleri üzerine etkisi. *Vet Hekim Der Derg*, 86, 22-35, 2015.
- 48. Mansoub NH:** Comparitive effect of butyric acid and probiotic on performance and serum composition of broiler chickens. *Adv Environ Biol*, 5, 1188-1191, 2011.
- 49. Peng QY, Li JD, Li Z, Duan ZY, Wu YP:** Effects of dietary supplementation with oregano essential oil on growth performance, carcass traits and jejunal morphology in broiler chickens. *Anim Feed Sci Technol*, 214, 148-153, 2016. DOI: 10.1016/j.anifeeds.2016.02.010
- 50. Chowdhury FN, Hossain D, Hosen M, Rahman S:** Comparative study on chemical composition of five varieties of groundnut (*Arachis hypogaea*). *World J Agric Sci*, 11 (5): 247-254, 2015.
- 51. Gao YY, Zhang XL, Xu LH, Peng H, Wang CK, Bi YZ:** Encapsulated blends of essential oils and organic acids improved performance, intestinal morphology, cecal microflora, and jejunal enzyme activity of broilers. *Czech J Anim Sci*, 64 (5): 189-198, 2019. DOI: 10.17221/172/2018-CJAS
- 52. Yang YF, Zhao LL, Shao YX, Liao XD, Zhang LY, Lu L, Luo XG:** Effects of dietary graded levels of cinnamon essential oil and its combination with bamboo leaf flavonoid on immune function, antioxidative ability and intestinal microbiota of broilers. *J Integr Agric*, 18 (9): 2123-2132, 2019. DOI: 10.1016/S2095-3119(19)62566-9