

## RESEARCH ARTICLE

# Identification of LncRNA Expression in the Estrous Cycle of Qira Black Sheep and Its Combination with miRNA Analysis

Xi CHEN<sup>1,†,a</sup> Hanying CHEN<sup>2,†,b</sup> Song JIANG<sup>1,c</sup> Hong SHEN<sup>1,d</sup> Xiancun ZENG<sup>1,e(\*)</sup><sup>†</sup> These authors contributed equally to this study<sup>1</sup> College of Animal Science and Technology, Shihezi University, 832003 Shihezi, Xinjiang, CHINA<sup>2</sup> School of Pharmacy, Shihezi University, 832002 Shihezi, Xinjiang, CHINAORCID: <sup>a</sup> 0000-0003-0457-1823; <sup>b</sup> 0000-0002-3792-0316; <sup>c</sup> 0000-0002-1690-3939; <sup>d</sup> 0000-0002-6672-5480; <sup>e</sup> 0000-0002-4250-7817

Article ID: KVFD-2021-26203 Received: 09.04.2021 Accepted: 19.10.2021 Published Online: 27.10.2021

## Abstract

In order to investigate the expression of lncRNAs in the ovaries of Qira black sheep at different stages of the estrous cycle, Qira black sheep were used as experimental materials in this experiment, and after estrus synchronization, ovarian tissues at four different stages of the estrous cycle were collected to extract total RNA. The samples at different stages were first subjected to genome-wide analysis using RNA-seq technology; target genes of lncRNAs were predicted using co-expression methods; and then GO and KEGG analysis of target genes was performed. Genes related to the estrous cycle of Qira black sheep were also selected to study their transcriptional differences. Finally, lncRNA-miRNA and mRNA-lncRNA-miRNA interaction networks were established to further analyze the effect of lncRNA alignment on Qira black sheep reproduction. The results showed that in the differentially expressed part, 14 lncRNAs were differentially expressed in Estrous VS Diestrus; the differential expression levels of lncRNAs in the two comparison groups of Estrous VS Metestrus and Estrus VS Proestrus were 18 and 24, respectively. The results of GO and KEGG functional enrichment analysis showed that differentially expressed lncRNAs and their target genes were mainly involved in reproduction-related pathways such as retinol metabolism, ovarian sterol production and endosterol biosynthesis. In the combined analysis of lncRNA-miRNA and mRNA-lncRNA-miRNA, genes related to reproduction, such as LNC011583, LNC003443 and bta-miR-202, were found, thus it can be seen that lncRNAs have some effect in the reproduction of Qira black sheep.

**Keywords:** Estrous, Ovary, LncRNA, Qira black sheep

## Qira Kara Koyunlarında Östrus Siklusunda LncRNA Ekspresyonunun Belirlenmesi ve miRNA Analizi İle Kombinasyonu

### Öz

Bu çalışmada, Qira kara koyunlarının ovaryumlarında lncRNA'ların ekspresyonunun araştırılması için materyal olarak östrus siklusunun farklı aşamalarında olan Qira kara koyunları kullanıldı. Total RNA eldesi amacıyla, östrus senkronizasyonunu takiben siklusun dört farklı aşamasından ovaryum dokuları toplandı. Farklı aşamalarındaki örnekler, önce RNA-seq teknolojisi kullanılarak genom çapında analize tabi tutuldu, lncRNA'ların hedef genleri birlikte ifade edilme yöntemleri kullanılarak tahmin edildi ve takiben hedef genlerin analizi GO ve KEGG ile gerçekleştirildi. Qira kara koyunlarında östrus siklusu ile ilgili genler, transkripsiyonel farklılıklarının araştırılması amacıyla seçildi. Son olarak, lncRNA hizalamasının Qira kara koyunlarında üreme üzerine etkisini daha fazla analiz etmek için lncRNA-miRNA ve mRNA-lncRNA-miRNA etkileşim ağları kuruldu. Sonuçlar, çeşitli ekspresyon bölgelerinde, 14 lncRNA'nın diöstrusa göre östrusta farklı şekillerde eksprese edildiğini gösterdi ve metöstrusa karşı östrus ve proöstrusa karşı östrus gruplarında lncRNA'ların diferansiyel ekspresyon seviyeleri sırasıyla 18 ve 24 saptandı. GO ve KEGG fonksiyonel zenginleştirme analizleri, farklı şekilde eksprese edilen lncRNA'ların ve bunların hedef genlerinin, esas olarak retinol metabolizması, yumurtalık sterol üretimi ve endosterol biyosentezi gibi üreme ile ilgili fonksiyonlarda yer aldığını gösterdi. lncRNA-miRNA ve mRNA-lncRNA-miRNA'nın kombine analizi sonucu, LNC011583, LNC003443 ve bta-miR-202 gibi üreme ile ilgili genler saptandı, dolayısıyla lncRNA'ların Qira kara koyunlarının üremesinde bir miktar etkisinin olduğu görülebilir.

**Anahtar sözcükler:** Östrus, Ovaryum, LncRNA, Qira kara koyunu

## INTRODUCTION

In general, the boundary of mammalian singleton and

multiple birth traits is relatively obvious, but there is diversity in this trait in sheep, and it is of great scientific value to study the mechanism of multiple birth in sheep<sup>[1]</sup>.

### How to cite this article?

Chen X, Chen H, Jiang S, Shen H, Zeng X: Identification of lncRNA expression in the estrous cycle of Qira black sheep and its combination with miRNA analysis. *Kafkas Univ Vet Fak Derg*, 27 (6): 733-740, 2021.  
DOI: 10.9775/kvfd.2021.26203

### (\*) Corresponding Author

Tel: +86 137 79204376

E-mail: zengxiancun@163.com (X. Zeng)



This article is licensed under a Creative Commons Attribution-NonCommercial 4.0 International License (CC BY-NC 4.0)



## Data Analysis

**Quality Control and Screening of lncRNAs:** Raw data in fastq format (raw reads) were first processed through an in-house perl script to obtain clean data, then bowtie2 (v2.2.8) (<http://bowtie-bio.sourceforge.net/bowtie2/manual.shtml>) was used to build the index of the reference genome, and finally HISAT2(v2.0.4) [12] (<http://daehwankimlab.github.io/hisat2/>) was used to align the clean reads at the paired ends to the reference genome (reference genome and gene model annotation files were downloaded directly from the genome website). StringTie (v1.3.1) [13] (<http://ccb.jhu.edu/software/stringtie/#contact>) was used, the mapped reads for each sample were assembled. Finally, we predicted target genes by co-expression.

**Differential Expression and Functional Enrichment:** First, by observing the distribution of differential transcripts on the chromosome, and the linked expression of their surrounding transcripts, we can pick out the target genes related to the study. Then, hierarchical cluster analysis was performed by FPKM expression levels of differential transcripts under different experimental conditions. Finally, through the volcano plot, the overall distribution of differential transcripts or genes can be visually seen. In this experiment, we selected candidate lncRNAs for subsequent analysis based on the transcriptional assembly results of lncRNAs, combined with the functional characteristics of lncRNA non-coding proteins as well as their structural characteristics. Differential expression in gene expression data was determined using Cuffdiff (v2.1.1) (<http://cole-trapnell-lab.github.io/cufflinks/manual/>) software and the Ballgown suite, where transcripts with  $P < 0.05$  were designated as differentially expressed.

Gene Ontology (GO) enrichment analysis of differentially expressed genes or lncRNA target genes was implemented by the Goseq R software package. We used KOBAS software to test the statistical enrichment of differential expression genes or lncRNA target genes in KEGG pathways [14]. KEGG is a database resource for understanding high-level functions and utilities of the biological system [15] (<http://www.genome.jp/kegg/>).

**lncRNA-miRNA-mRNA Association Analysis:** lncRNAs, as a type of ceRNAs (competing endogenous RNAs) [16,17], can competitively bind miRNAs with genes, so we searched

for lncRNA-miRNA-gene pairs that also possess miRNA binding sites for combined analysis.

**qRT-PCR:** QRT-PCR was used in this experiment to validate gene expression levels. The amplification procedure was performed using a two-step method: 95°C 30 s; 95°C 5s, 60°C 30 s, for 45 cycles in total. Each sample was repeated three times for a total of 12 samples. Information about primers for qRT-PCR using GAPDH as the reference gene is shown in Table 2. Finally, the statistical analysis of the test data was performed using the  $2^{-\Delta\Delta ct}$  method for relative quantitative analysis, and the statistical software SPSS 17.0 was used for significance test.

## RESULT

### Overview of Sequencing Data

In total, there were 12 libraries in this study, as shown in Table 1, resulting in 1,694,019,460 raw reads. In order to ensure the quality of the data and remove the adapter-bearing, low-quality reads inside, 1,647,629,538 clean reads were obtained after screening. Among them, clean reads accounted for 99.99% of raw reads. 91.73% ~ 93.99% of clean reads in all libraries were successfully mapped to the reference genome (Table 3). To further validate the data, we selected 2 differentially expressed lncRNAs targeting reproduction-related genes and determined their expression levels by qRT-PCR, and the results were verified to be consistent with those obtained by sequencing (Fig. 1).

### Screening of lncRNAs

A total of 11981 lncRNAs, 13,831 TUCPs and 22,824 mRNAs were identified in all samples for further analysis (Fig. 2-A). We obtained Pearson correlation coefficients between 0.875 and 0.949 among the samples, of which the highest correlation with QD1 is QM3 (0.949) (Fig. 2-B). The results showed that the expression levels of ovaries were almost consistent during the four stages of the estrous cycle (Fig. 3-A), while the expression levels of lncRNAs and TUCP were much lower than those of mRNAs (Fig. 3-B).

### Differential Expression Analysis


In this experiment, we identified 11981 lncRNAs and 22,824 mRNAs from the ovaries of Qira Black sheep for further

**Table 2.** lnc-010801, lnc-008562 and internal reference amplification primers

Primer	Gene Sequence	Fragment Size/bp
GAPDH-F	CCTGCCAAGTATGATGAGAT	119
GAPDH-R	TGAGTGTCTGCTGTTGAAGT	
LNC-010801-F	GCGGGAAGTCTGTCTCT	124
LNC-010801-R	CGAAAAGTCCGAAACACCAG	
LNC_008562-F	CGCCAAATCGGAGTAAACA	100
LNC_008562-R	AATTCATCCAGGCAGGGTC	

**Table 3.** The summary of sample data quality


Sample Name	Raw Reads	Clean Reads	Clean Bases	Error Rate(%)	Q20 (%)	Q30 (%)	GC Content (%)
He1	90840870	88180586	13.23G	0.01	97.79	94.2	49.09
He2	120787312	117362696	17.6G	0.01	97.7	93.94	48.18
He3	114504810	111147030	16.67G	0.01	97.88	94.37	48.6
QD1	121671830	118548894	17.78G	0.01	97.73	94.03	48.26
QD2	111730482	108979944	16.35G	0.01	97.8	94.18	47.82
QD3	123522800	121471712	18.22G	0.01	97.15	93.23	48.47
QE1	109926982	106702328	16.01G	0.01	97.66	93.95	48.42
QE2	115505728	112006036	16.8G	0.01	97.75	94.15	48.63
QE3	121606340	118252002	17.74G	0.01	97.72	94.08	47.17
QM1	113856982	110990072	16.65G	0.01	97.78	94.15	49.28
QM2	121209526	117616398	17.64G	0.01	97.62	93.8	47.37
QM3	116530402	113689300	17.05G	0.01	97.75	94.1	47.42
QP1	110697120	107454358	16.12G	0.01	97.85	94.35	50.25
QP2	103318274	100061160	15.01G	0.01	97.82	94.27	49.01
QP3	98310002	95167022	14.28G	0.01	97.44	93.39	47.96

**Fig 1.** QRT-PCR validation of differentially expressed genes

was the most differentially expressed, with 15 up-regulated and 9 down-regulated. The remaining two comparison groups also had up- and down-regulated genes, respectively. From the lncRNA differential expression clustering results plot (Fig. 4-D), we can see that the three proestrus groups (QP1/2/3) and the three estrus groups (QE1/2/3) were clustered, indicating that the differences between these two groups were quite different.

### GO and KEGG Enrichment Analysis


In this experiment, target genes were predicted by means of co-expression. A total of 13,0152 potential target genes were identified by co-expression, and GO and KEGG analyses were performed on these potential target genes.

**Fig 2.** Screening results (A) and correlation expression (B)


analysis. Differential expression analysis was performed using Ballgown. In this experiment, we will elaborate the overall differential expression of lncRNAs from three comparison groups: QE\_VS\_QD, QE\_VS\_QM and QE\_VS\_QP.

We obtained differential expression results for the three comparison groups (Fig. 4-A,B,C). Among them, QE\_VS\_QP

After obtaining the 56 differentially expressed lncRNAs, the target genes of these lncRNAs were subjected to GO enrichment analysis and their functions were described. GO analysis showed that all were divided into BP (biological process), CC (cellular component) and MF (molecular function). From the GO enrichment histogram (Fig. 5-A), the GO terms of these differentially expressed lncRNA




**Fig 3.** Comparison of expression levels of each group (A) and each gene (B)



**Fig 4.** Differential expression in each comparison group (A,B,C) and differential expression clustering results (D)

target genes are enriched for several processes, such as: transcription, DNA-dependent, RNA biosynthetic process, neutral lipid catabolic process, etc.

The scatter plot is a graphical presentation of the KEGG enrichment analysis results, and we picked the 20 pathway entries with the most significant enrichment for




**Fig 5.** GO (A) and KEGG (B) analysis results of differentially expressed lncRNA target genes

presentation in this plot. KEGG analysis of heterologously expressed co-expressed lncRNA target genes showed that they were rich in pathways related to reproduction: Jak-STAT signaling pathway, retinol metabolism, Ovarian steroidogenesis, etc. (Fig. 5-B).

### ***LncRNA-miRNA-mRNA Association Analysis***

By constructing an interactive network diagram of miRNA-lncRNAs (miRNAs have been proposed in other literature), we can better understand the involvement and function of lncRNAs in the estrous cycle of Qira Black Sheep. We

constructed the interaction network of miRNA-lncRNAs separately for the three comparison groups (Fig. 6-A,B,C). We obtained 248 miRNA-lncRNA interaction combinations with potential relationship compared by QE\_VS\_QP, of which LNC001917, LNC011056, LNC003367, LNC009395, LNC003443, etc. could interact with multiple different miRNAs, so they were in the middle position, and it can be speculated that these genes play a role in the regulation of estrous cycle in Qira black sheep. However, in the QE\_VS\_QD and QE\_VS\_QM comparison groups, we also obtained 175 and 22 interaction network graphs with potential relationships, respectively.



**Fig 6.** Target lncRNA-miRNA interaction net (A,B,C) and Target miRNA-lncRNA-mRNA interaction net (D)

We also constructed an interaction network map of miRNA-lncRNA-mRNA (Fig. 6-D). We obtained that in QE\_VS\_QP, bta-miR-1343-5p, bta-miR-2411-5p, bta-miR-2387, etc. are at the comparative core position because they are able to interact with the most genes. Among them, LNC001917, LNC003443 and LNC001763 can interact with multiple different miRNAs, so we speculated that these genes may play a role in the regulation of the estrous cycle in Qira Black sheep.

## DISCUSSION

From the results of GO and KEGG enrichment analysis in this experiment, it can be seen that the most significantly different GO term is the term of the cofactor. However, it has been shown that <sup>[18]</sup>, the cofactors may catalyze the synthesis of steroid hormones by increasing enzyme activity. In the results of KEGG enrichment analysis of lncRNAs, there is also a pathway of cytochrome P450 enzymes, which are expressed in both ovarian testes and play an important role in the synthesis of anabolic steroids and maintenance of sex hormones <sup>[19]</sup>. In KEGG enrichment analysis of lncRNA target genes, we obtained pathways related to fecundity as: retinol metabolism, ovarian endosterol production and endosterol biosynthesis. Studies have found that retinol and its derivatives play an important role in ovarian steroidogenesis, oocyte maturation and

early embryonic development, are abundant in cumulus granulosa cells, and are involved in signaling related to ovarian development <sup>[20-22]</sup>.

In the combined analysis of lncRNA-miRNAs, we obtained that bta-miR-26b was present in the gene interacting with LNC011583 in the QE\_VS\_QD comparison, which was pointed out by <sup>[23]</sup> that it belongs to the miR-26 family and plays a key role in estrogen stimulation <sup>[24]</sup>. So we speculated that LNC011583 gene, which interacts with bta-miR-26b, may play a role in estrogen stimulation, thereby affecting the content of estrogen. We also obtained oar-miR-432 and bta-miR-202 genes interacting with LNC003443 gene in QE\_VS\_QP comparison. Among them, the oar-miR-432 gene is likely to function by directly or indirectly affecting gonadotropin-releasing hormone (GnRH) activity associated with reproductive hormone release <sup>[25]</sup>. It has been proposed <sup>[26]</sup> that the bta-miR-202 gene is expressed only in the gonads. On the one hand, we can speculate that when LNC003443 gene interacts with oar-miR-432, it may have an effect on the reproduction of Qira Black sheep by affecting the activity of gonadotropin-releasing hormone (GnRH); on the other hand, when LNC003443 gene interacts with bta-miR-202 gene, it may have an effect on the ovary and thus affect the reproduction of Qira Black sheep.

In this experiment, the target genes of lncRNA-controlled

reproduction-related pathways were obtained by GO and KEGG enrichment analysis. The results showed that the target genes of lncRNAs had an effect on reproductive hormone synthesis such as steroids and retinol, thus we speculated that lncRNAs had some effect on Qira black sheep reproduction. Finally, we also performed a combined analysis combining mRNA and miRNA, speculating on genes that may be associated with reproduction, such as LNC011583 and LNC003443. It further illustrates the expression of lncRNAs during the estrous cycle in Qira black sheep. This experiment provides some reference for the study of fertility in Qira black sheep.

### AVAILABILITY OF DATA AND MATERIALS

The datasets during and/or analyzed during the current study available from the corresponding author on reasonable request.

### ACKNOWLEDGEMENTS AND FUNDING INFORMATION

This research was supported by the National Natural Science Foundation of China (31660643 and 31260536). The authors have no competing interests.

### CONFLICT OF INTEREST

The authors declared that there is no conflict of interest.

### AUTHOR CONTRIBUTIONS

XC Zeng designed the study, conducted the experiments, analysed the data, and drafted the manuscript. HY Chen and X Chen designed the study and drafted the manuscript. S Jiang, H Shen conducted parts of the experiments and collected samples.

### REFERENCES

- Bai J:** Candidate Genes of Multiparity in Xinjiang Qira Black Sheep and Duolang Sheep. Shihezi University, 2007. DOI: 10.7666/d.y1165085
- Yu J, Dong H, Ji Y, Kong Y, Guligena, Meizi B, Yang F, Yassen K, Wen Q, Mizinisha:** Current situation of sheep breed resources and protection and utilization measures in Hotan. *Gra-fee Livest*, 2, 7-9, 2011. DOI: 10.3969/j.issn.1003-6377.2011.02.003
- Tan R, Cong LY, Yang QY, Xu CS:** Study on protection and utilization scheme of Qira Black Sheep. *Gra-fee Livest*, 3, 31-33, 2011. DOI: 10.3969/j.issn.1003-6377.2011.03.011
- Shao YY, Megenisha K, Yan MY, Yan MY, Fan QX, Yuan XF, Zeng XC:** Expression of LNC-010801 and LNC-008562 in the ovary of Qira Black Sheep at different stages of estrous cycle. *Chin Herb Sci*, 38 (6): 9-12, 2018. DOI: 10.3969/j.issn.2095-3887.2018.06.003
- Gong Y, Ma HM:** Progress in transcriptional lncRNA in pigs. *Swine Ind Sci*, 296 (2): 100-106, 2020.
- Birney E, Stamatoyannopoulos JA, Dutta A, Roderic Guigó, Gingeras TR, Margulies EH, Weng Z, Snyder M, Dermitzakis ET:** Identification and analysis of functional elements in 1% of the human genome by the ENCODE pilot project. *Nature*, 447, 799-816, 2007. DOI: 10.1038/nature05874
- Chen LL, Feng SS, Fan ZS, Gong C, Liu BY, Liu ZH, Li CW, Song EW, Sun SH, Wu GZ, Wu H, Wu M, Xu G, Yuan JH, Zeng CY, Zhu YM:** Progress in non-coding RNA research. *Sci Sin Vita*, 49 (12): 1573-1605, 2019. DOI: 10.1360/SSV-2019-0179
- Niu CY, Xue LL, Ji H, Li SZ:** Progress in the biological function of lncRNAs. *Chin J Biol Prod*, 32 (2): 228-232, 2019.
- Feng X, Li F, Wang F, Zhang G, Pang J, Ren C, Zhang T, Yang H, Wang Z, Zhang Y:** Genome-wide differential expression profiling of mRNAs and lncRNAs associated with prolificacy in Hu sheep. *Biosci Rep*, 38 (2): BSR20171350, 2018. DOI: 10.1042/BSR20171350
- Han DX, Sun XL, Wang CJ, Yu ZW, Zheng Y, Huang YJ, Wang WH, Jiang H, Gao Y, Yuan B:** Differentially expressed lncRNA-m433s1 regulates FSH secretion by functioning as a miRNA sponge in male rat anterior pituitary cells. *Biol Reprod*, 101, 416-425, 2019. DOI: 10.1093/biolre/ioz100
- Zheng LP, Luo RC, Su T, Hu LL, Gao FX, Zhang XN:** Differentially expressed lncRNAs after the activation of primordial follicles in mouse. *Reprod Sci*, 26 (8): 1094-1104, 2019. DOI: 10.1177/1933719118805869
- Langmead B, Salzberg SL:** Fast gapped-read alignment with Bowtie 2. *Nat Methods*, 9 (4): 357-359, 2012. DOI: 10.1038/nmeth.1923
- Pertea M, Daehwan K, Geo MP, Jeffrey TL, Salzberg SL:** Transcript-level expression analysis of RNA-seq experiments with HISAT, StringTie and Ballgown. *Nat Protoc*, 11 (9): 1650-1667, 2016. DOI: 10.1038/nprot.2016.095
- Mao XZ, Cai T, Olyarchuk JG, Wei LP:** Automated genome annotation and pathway identification using the KEGG Orthology (KO) as a controlled vocabulary. *Bioinformatics*, 21 (19): 3787-3793, 2005. DOI: 10.1093/bioinformatics/bti430
- Minoru K, Michihiro A, Susumu G, Masahiro H, Mika H, Masumi I, Toshiaki K, Shuichi K, Shujiro O, Toshiaki T:** KEGG for linking genomes to life and the environment. *Nucleic Acids Res*, 36, D480-D484, 2018. DOI: 10.1093/nar/gkm882
- Salmena L, Poliseno L, Tay Y, Kats L, Pandolf PP:** A ceRNA hypothesis: The rosetta stone of a hidden RNA language. *Cell*, 146, 353-358, 2011. DOI: 10.1016/j.cell.2011.07.014
- Tay Y, Rinn J, Pandolf PP:** The multilayered complexity of ceRNA crosstalk and competition. *Nature*, 505, 344-352, 2014. DOI: 10.1038/nature12986
- Xie J:** Analysis of lncRNA and mRNA in Goat Ovary at Different Stages of Estrous Cycle. *Fuyang Normal University*, 2019.
- Hakki T, Silvia Z, Dragan CA, Bureik M, Bernhardt R:** Coexpression of redox partners increases the hydrocortisone (cortisol) production efficiency in CYP11B1 expressing fission yeast *Schizosaccharomyces pombe*. *J Biotechnol*, 133, 351-359, 2008. DOI: 10.1016/j.jbiotec.2007.06.022
- Brown JA, Eberhardt DM, Schrick FN, Roberts MP, Godkin JD:** Expression of retinol-binding protein and cellular retinol-binding protein in the bovine ovary. *Mol Reprod Dev*, 64 (3): 261-269, 2003. DOI: 10.1002/mrd.10225
- Mohan M, Thirumalapura NR, Malayer J:** Bovine cumulus granulosa cells contain biologically active retinoid receptors that can respond to retinoic acid. *Reprod Biol Endocrin*, 1:104, 2003. DOI: 10.1186/1477-7827-1-104
- Zhao YY, Zhang YM, Liu Y, Li WT, Yang YL:** Progress in the role of retinol in ovarian development. *Chin J Anim Husbandry*, 55 (5): 1-5, 2019.
- Gu B, Liu H, Han Y, Chen Y, Jiang HZ:** Integrated analysis of miRNA and mRNA expression profiles in 2-, 6-, and 12-month-old Small Tail Han Sheep ovaries reveals that oar-miR-432 downregulates RPS6KA1 expression. *Gene*, 710, 76-90, 2019. DOI: 10.1016/j.gene.2019.02.095
- Maillot G, Lacroix-Triki M, Pierredon S, Gratadou L, Schmidt S, Bénès V, Roché H, Dalenc F, Auboeuf D, Millevoi S, Vagner S:** Widespread estrogen-dependent repression of microRNAs involved in breast tumor cell growth. *Cancer Res*, 69, 8332-8340, 2009. DOI: 10.1158/0008-5472.can-09-2206
- Zhang ZB, Tang JS, Di R, Liu QY, Hu W:** Integrated hypothalamic transcriptome profiling reveals the reproductive roles of mRNAs and miRNAs in sheep. *Front Genet*, 10:1296, 2020. DOI: 10.3389/fgene.2019.01296
- Sontakke SD, Mohammed BT, McNeilly AS, Donadeu FX:** Characterization of microRNAs differentially expressed during bovine follicle development. *Reproduction*, 148 (3): 271-283, 2014. DOI: 10.1530/rep-14-0140